



C 6. 8

R41276



Digitized by the Internet Archive
in 2015

https://archive.org/details/b21901272_0001



FLORA BRITANNICA.

V O L. I.



Ex libris

FLORA BRITANNICA,

Coll. Rey

AUCTORE

Med. Edin.

JACOBO EDKARDO SMITH, M.D.

H. L.

SOCIETATIS LINNÆANÆ PRÆSIDE,

REGIÆ LONDINENSIS, HOLM. UPSAL. TAURIN. OLYSSIP.

PHILADELPH.—PHYSIOGR. LUND. BEROLIN. PARIS.

ALIARUMQUE SOCIETATUM SOCIO.

“Nullius addictus jurare in verba magistri.”

HOR.

VOL. I.

LONDINI:

TYPIS J. DAVIS, CHANCERY-LANE:

VENEUNT APUD J. WHITE, FLEET-STREET.

MDCCC.

VIRO ILLUSTRISSIMO

J O S E P H O B A N K S,

REGI A CONSILIIS INTIMIS,

BARONETO, BALNEI EQUITI,

REGIÆ SOCIETATIS PRÆSIDI,

SOCIETATIS LINNÆANÆ SOCIO HONORARIO,

HOC OPUS,

IN SUMMÆ OBSERVANTIÆ PIGNUS,

DAT DICAT

AUCTOR.

P R Æ F A T I O.

EX cœli nostri beatissimâ temperie, tum ex variis Dei Optimi Maximi muneribus, quæ ad solum situmque nostrum insularem spectant, Flora Britannica semper inter copiosissimas Florarum Europæarum habita est. Præ ceteris quidem terris accuratiùs indagata fuit Britannia. Quæ autem regio unquam hominem Raio nostro peritiorem jactaret? Vix ulla æquè accuratum. Fatendum est, quod pauci quidam botanici apud nos ante illum rei herbariæ aliquod auxilii, studio at certè partiali, irregulari, ideoque imperfecto, contulissent. Proculdubiò autem Raius jure summo pater Floræ Britannicæ dici meritus est. Scilicet extemplò practici omnes apud nos, illius opera sedulò versabant. Ex illius etiam operibus, quas plantas nostra tulerit patria omninò peregrini didicerunt. Hinc evenit, quod cum Linnæi Systema novum, botanices scientiæ accommodatissimum, ideoque citiùs animos modo non omnium facilè occupaturum, prodit,
non

P R Æ F A T I O.

non nisi ad Raii opera redactum collatumque apud botanicos nostros intelligi potuit ; itaque inter antiqua et nova imperia quasi medius stetit Raius. Linnæus profectò Raii opera studiosus citavit ; atque illi qui sub novis Linnæi legibus tempora feliciora sperarent, imprimis Raii dicta sollicitè perpendebant. Quoniam autem inter auctores commercium multis de causis imperfectum fuit, pluri mi errores locum habuerunt. Aliquando Raius ipse, et sæpissimè editor ejus Synopsis Dillenius, synonyma auctorum quos citaverunt perperàm intellexerunt. Sæpè etiam celeberrimus Linnæus parili modo hallucinatus est. Et quamvis Hudsonus, primus nostratum qui Linnæi Systema interpres exposuit, discipulique ejus hisce obscuritatibus lucis quoddam dederunt, tamen fontes, ex quibus veram integramque scientiam haurirent, nimis pauci illis innotuerunt ; ne dicam, quòd timor, ne ex nimia audentiâ rei botanicæ vim inferrent, plus æquo corripuit quam quo finem propositum rectâ irent viâ.

Auctor hujusce operis causas, uti sperat, horum errorum probè perspexit. Melioribus igitur auspiciis, quam quæ illis dignissimis viris concessa fuerunt, usus ; præsertim cum ex possessione Musæi Linnæani sibi facultas data est specimina Linnæana adeundi, conferendi, comprobandique, Floram Britannicam haud timuit de novo investigare. Nihil pro comperto habuit, quod non ipse scrutatus est.

P R Æ F A T I O.

est. Omnem specificum, omnem genericum characterem ad incudem cautus retulit, et quâcunque res postularet, reformavit. Synonyma auctorum perpensè exquisivit : si denique animus alicubi in dubio hæreret, ad herbaria originalia sese contulit. Descriptiones suas ex sylvestribus plantis, quâcunque liceret, fidus desumpsit : brevi, omnia tam laboriosè investigavit, quam si nemo hætenus aut tractavisset rem, aut etiam in animo habuisset. Non verò audet sperare quod ipse solus omninò errorum fuerit immunis, atque opus immaculatum oculis lectorum subjecerit. Hoc autem liceat profiteri, quod nullibi arte affectatâ, aut ambagibus verborum, quæ ignoraret studuit cclarc. Hæc solummodo, si ullam forsân sibi fas sit vindicare, laus detur, quòd summoperè conatus est errores vitare, et quæ non potuit accuratè determinare, palam sapientioribus proponere, eorundemque opem poscere.

Quod ad classificationem attinet, scèrè omninò Linnæi vestigiis inniti voluit ; neque enim adhuc sensit quid commodi hodiernæ Systematis Linnæani, planè artificialis, mutationes rei botanicæ contulerint. Omissio igitur ordinis Syngenesiæ Monogamiæ ; tum amotio generum, in quibus flores quantumvis organis sexualibus, at non aliter structurâ differre visa sint, ex classe Polygamiâ ; atque quædam aliæ generum muscorum mutationes, quas necessitas imperiosa jussit (verùm enimverò et in his tractandis quasi medio ire studuit) et dispositio alia

P R Æ F A T I O.

rerum quarundam Cryptogamicarum, sunt præcipuæ innovationes quæ in hoc opere obviàm erunt.

In habitationibus plantarum rariorum notandis Raium secutus est, et nomina locorum, quò nostrates faciliùs intelligant, linguâ scripsit vernaculâ. Si ex vivorum quorundam indiciis habitationes acceperit, horum nomina (nisi si sibi ea loca priùs innotuissent) candidus impertivit. In descriptionum structurâ, ad illa quæ ipse vidit et pro certo comperit confirmanda, ullam scriptorum aut studiosorum auctoritatem adducere supervacaneum fore duxit : haud tamen, e contra, aliquid sibi ignotum protulit, nisi auctore, quo fretus est, priùs memorato.

Cum auctores citat, eorum verba aut figuras citat, minimè verò synonyma ibi collata : non quòd si synonymon aliquod apud illos omiserit, vellet fidem accurationi ejus detrahere. Quæcunque synonyma ad opus elucidandum necessaria fore putaverit, et sine dubio locis citatis accommodata, illa studiosè contulit.

Plurimorum amicorum auxilium, quorum nomina suis locis inscribentur, lubens gratusque admodùm agnoscit. Neque etiam non potest quin pectus gaudium tacitum et dulcissimum pertentet de expectatione operis hujusce, quæ undique ex ore omnium diutiùs expressa est. Hanc expectationem quàm minimè frustrari voluit ; dumque ideò obsequi festinanter studet, erroribus, in quos in-

P R Æ F A T I O.

cautus incederit, sperat et obsecrat lectores ignoscere.

Pauculæ quædam plantæ hanc notam (†) præse ferunt. Hæ sunt quas plantas auctor non verè indigenas esse suspicatur; aut jamdudum periisse; vel alias quasdam malè pro his plantis, sub quarum nomine inclaruerunt, acceptas fuisse. Hæc autem nota eradenda est ante Poam bulbosam, p. 102, quam Dom. Stone nuper prope Yarmouth spontaneam invenit. Asteriscus (*) præfixus innuit hanc plantam olim ex hortis evasisse, et demùm habitationem apud nos liberam et verè suam elegisse.

Dum omnia hæc auctor memorat, dolet sanè quòd opus prodit imperfectum. At pretium papyri, hodiè omni ratione alienum, cogit labores in præsens interrumpere. Amicis ergo suadentibus, visum est duo prima volumina, cum indice suo, jam in lucem mittere. Intra anni spatium spes est impedimentum istud rei literariæ, satis mirum, ferè inauditum, cessaturum fore. Aliquis tamen usus ex hoc malo eveniat: nempe illæ hallucinationes aut omissiones quæ in his duobus voluminibus detegantur, (et has ad se bono animo transmitti orat,) in sequentibus forsàn corrigantur. Index copiosissimus totius operis, et catalogus auctorum citatorum, ad calcem adjicientur.

Dabam NORVICI,

Pridie Iduum Martii, MDCCC.

Classis I.

MONANDRIA.

Stamen 1.

MONOGYNIA.

Stylus 1.

* *Gymnospermæ.*

1. SALICORNIA. *Cal.* monophyllus. *Cor.* nulla.
2. HIPPURIS. *Cal.* obsoletu *Cor.* nulla.
Valeriana rubra.

** *Angiospermæ.*

3. CHARA. *Bacca* polysperma.
4. ZOSTERA. *Spadix* multiflorus. *Caps.* monosperma.

DIGYNIA.

Styli 2.

5. CALLITRICHE. *Cal.* nullus. *Petala* 2:
Semina 4, nuda.

I. MONANDRIA.

I. MONOGYNIA.

1. SALICORNIA. *G. Pl.* 5. *Juss.* 86.

Calyx ventricosus, integer. *Petala* nulla. *Stamina* 1 vel 2. *Semen* 1, calyce tectum.

herbacea. 1. SALICORNIA geniculis compressis emarginatis, internodiis obconicis, spicis pedunculatis apicem versus attenuatis.

Salicornia herbacea. *Sp. Pl.* 5. *With.* 4. *Relb. Suppl.* 2, 1. *Fl. Dan.* t. 303.

S. annua. *Afzel. Mfs.* 1. *Bast. Opusc.* v. 2. 105, t. 10. *Engl. Bot.* t. 415.

S. europæa α. *Huds.* 1.

Salicornia. *Raii Syn.* 136.

S. five Kali geniculatum. *Ger. em.* 535.

Small jointed Glasswort. *Pet. H. Brit.* t. 9. f. 3.

β. *Salicornia erecta*, foliis brevibus, cupressiformis. *Dill. in Raii Syn.* 137.

S. biennis. *Afzel. Mfs.* 2.

S. europæa β. *Huds.* 1.

Woody jointed Glasswort. *Pet. H. Brit.* t. 9. f. 4.

γ. *Salicornia myosuroides procumbens*, furculis longissimis. *Dill. in Raii Syn.* 137.

S. europæa δ. *Huds.* 1.

δ. *S. ramosior procumbens*, foliis brevibus purascentibus. *Dill. in Raii Syn.* 137.

S. europæa η. *Huds.* 1.

ε. *Kali geniculatum majus sempervirens humilius.* *Schola Bot.* 37. *Dill. in Herb. Sberard.*

Angl. Marsh Samphire or Jointed Glasswort.

In

In littoribus maritimis inundatis cœnosis vel arenosis frequens.

β, γ, δ, found by Mr. Stonestreet in a little salt-marsh on the east side of Pool, Dorsetshire; ε was also gathered by the same gentleman, in what place is not mentioned.

Annua vel biennis. Fl. Augusto, Septembri.

Radix fibrosa, parva. Caulis plerumque erectus, basi subdivisus, supernè ramosus, ramis oppositis, simplicibus, erectis, valdè succulentis, aphyllis, geniculatis, geniculis compressis, apice dilatatis, emarginatis. Spicæ oppositæ cum impari majore terminali, pedunculatæ, teretes, apicem versus sensim attenuatæ, acutiusculæ, articulatæ. Flores oppositi, utrinque subterni.

β gaudet caule ramosiori, spicis longioribus et crassioribus; γ caule diffusio, spicis longissimis; δ caule ramosissimo, spicis parvis, purpurascentibus; ε caule ramosissimo firmitiori, spicis brevibus, viridibus.

2. SALICORNIA geniculis teretibus integris, infruticosa. ternodiis æqualibus, spicis subsessilibus cylindraceis obtusis.

Salicornia fruticosa. *Sp. Pl.* 5. *With.* 3. *Afzel. Mss.* 3.

S. europæa γ. *Huds.* 1.

Kali geniculatum perenne fruticosius procumbens. *Raii Syn.* 136.

Kali. *Camer. Epit.* 246.

Angl. Shrubby Samphire or Jointed Glasswort.

In palustribus marinis.

Near the Isle of Shepey. *Sir Hans Sloane, Bart.*

In the Isle of Grain in the Thames. *Dillenius.*

On the sea-shore all the way from Weymouth turnpike to Rhodipole; Mr. Yalden. *Lightfoot Mss.*

Perennis. Fl. Augusto, Septembri.

Radix lignosa, perennis. Caulis suffruticosus, adscendens,

ramosissimus, ramis ramulisque oppositis, minusque carnosus. Internodia omnia ferè cylindracea. Spicæ plerumque sessiles, omnino cylindraceæ, obtusæ, multifloræ, articulis brevissimis.

2. HIPPURIS. *G. Pl.* 5. *Juss.* 18. *Gærtn. t.* 84.
Cal. obsoletus, integer. *Cor.* nulla. *Stigma*
 simplex. *Semen* 1, inferum.

vulgaris. 1. HIPPURIS foliis verticillatis linearibus.

Hippuris vulgaris. *Sp. Pl.* 6. *Suppl.* 81. *Huds.* 2.
With. 5. *Relb.* 1. *Sibth.* 1. *Abbot.* 1. *Curt.*
Lond. fasc. 4. *t.* 1. *Dicks. H. Sicc. fasc.* 2. 1.
Fl. Dan. *t.* 87.

Linnopeuce. *Raii Syn.* 136.

Cauda equina fœmina. *Ger. em.* 1114.

Equisetum palustre, *Linariæ scopariæ* folio. *Dill.*
in Raii Syn. 136. *Herb. Buddl. Gen.* 5. *p.* 7.
Angl. *Mare's-tail.*

In fossis et stagnis limosis, haud vulgaris.

Near Hornsey. *Huds.* Near Lynn, and in other parts of Norfolk.

Perennis. *Fl.* Maio.

Radix e fibris longis, verticillatis. *Caulis* simplicissimus, erectus, crebrè geniculatus; ad medium usque immersus atque defoliatus; supernè foliosus. *Folia* verticillata, 8—12, patentia, linearia, acuta, integerrima, glabra. *Flores* axillares, solitarii, sessiles.

Hyberno tempore, sub aquâ latens, gaudet foliis crebrioribus, longioribus, pallidis, tenuioribus et subpellucidis, ut in Herbario Buddleiano videas.

3. CHARA. *G. Pl.* 567. *Juss.* 18. *Gærtn. t.* 84.
Cal. nullus. *Cor.* nulla. *Anthera* sessilis.
Stylus nullus. *Bacca* polysperma.

vulgaris. 1. CHARA incrinis striata, foliis subulatis articulatis.

Chara vulgaris. *Sp. Pl.* 1624. *Huds.* 397.

With. 2. *Relb.* 345. *Sibth.* 1. *Engl. Bot.*

t. 336. *Hedw. Theor.* *t.* 32, 33. *Abbot.* 197.

C. vulgaris foetida. *Raii Syn.* 132. *Vaill. Act.*

Paris. Ann. 1719. 17. *t.* 3. *f.* 1.

C. n. 1681. *Hall. Hist.* v. 3. 4.

Equisetum foetidum sub aquâ repens. *Baub.*

Prodr. 25. *Ger. em.* 1115.

β. *Chara minor*, caulibus et foliis tenuissimis. *Raii*

Syn. 133. *Vaill. Act. Paris. Ann.* 1719. 18.

Angl. Common *Chara*.

In aquis cœnosis frequens ; β in purioribus.

Annua. Fl. Julio.

Herba immerfa, foetida, fragilis, plerumque terrâ calcareâ seruposâ albidâ undique obducta. Caulis pedalis, filiformis, ramosa, foliosa, striata, torta. Folia verticillata, suboctona, erecto-patentia, subulata, acuta, articulata, striata ; supernè canaliculata, florifera, foliolosa. Anthera nuda, sessilis, depressa, carnea, demum rimosa. Germen foliolis quaternis circumdatum, antheræ proximum, ovatum, spiraliter striatum, apice quinquedentatum. Bacea corticata, polysperma. Stigma atque impregnationis ratio ignota sunt.

Foliorum septa terreâ plerumque larvâ obtecta sunt, unde error in horum descriptionem apud *Engl. Bot.* accidit.

Var. β nequaquam me iudice differt, nisi quod, præ aquis purioribus, nuda est et lætè virens, nec crustâ calcareâ obvoluta.

2. *CHARA* *fulcata*, foliis subulatis articulatis, foliolis verticillatis, aculeis caulinis setaceis deflexis.

Chara hispida. *Sp. Pl.* 1624. *Huds.* 398. *With.* 3.

Relb. 345. *Engl. Bot.* *t.* 463.

C. major caulibus spinosis. *Raii Syn.* 132. *Vaill.*

Act. Paris. Ann. 1719. 18. *t.* 3. *f.* 3.

C. n. 1682. *Hall. Hist.* v. 3. 4.

β. *C. major subcinerca fragilis.* *Raii Syn.* 132.
Vaill. Act. Paris. Ann. 1719. 18.

C. tomentosa. *Huds.* 398. *Sibth.* 2. nec Linn.
Equisetum majus subcinereum aquis immersum.
Morif. Sect. 15. t. 4. f. 9. *Herb. Bobart.*
Angl. Prickly Chara.

In fossis et stagnis.

North of England. *Huds.* In a clay-pit at Wif-
 beach. *Mr. W. Skrimshire.*

β. In Oxfordshire, Kent, and near Bath. *Ray.*
 North of England. *Huds.*

Annua. Fl. Julio, Augusto.

Habitus prioris sed major, caule quintuplo crassiore, ful-
 cato potius quam striato, supernè præcipuè aculeato,
 aculeis superioribus omnibus deflexis, inferioribus di-
 rectione variis. Foliola verticillata. Flores ferè ut in
C. vulgari.

Var. β vix differt, nisi caule sub verticillis tantum supe-
 rioribus aculeato. Hujus synonyma ad Charam suam
 tomentosam perperam retulit Linnæus.

flexilis.

3. *CHARA inermis lævis diaphana, foliis cylindra-
 ceis obtusis mucronulatis.*

Chara flexilis. *Sp. Pl.* 1624. *Huds.* 398. *With.* 3.

C. translucens minor flexilis. *Dill. in Raii Syn.*
 133. *Vaill. Act. Paris. Ann.* 1719. 18. t. 3.
 f. 9.

Conserva nidifica. *Fl. Dan. t.* 761.

Angl. Smooth Chara.

In fossis et lacubus.

Frequent in the ponds about Henley not far
 from Ipswich; Buddle. *Dill.* In maritime
 ditches, and near Hornsey. *Huds.*

Annua.

Annua. Fl. Julio, Augusto.

Virens, pellucida, glabra, exstriata. Caulis teres, ramosus, æqualis, fistulosus. Folia verticillata, cylindracea, filiformia, obtusa, mucronulata, sæpe articulata et etiam ramosa, inde potius ramuli dicenda; florifera aggregata, terminalia, breviora.

4. ZOSTERA. *G. Pl.* 472. *Juss.* 24. *Gærtn.* t. 19.

Spadix linearis, intra foliorum vaginam, hinc florifer. *Cal.* nullus. *Cor.* nulla. *Anthera* sessilis, germiui opposita. *Stigmata* duo, linearia. *Caps.* monosperma.

1. ZOSTERA pericarpis sessilibus.

marina.

Zostera marina. *Sp. Pl.* 1374. *Huds.* 395.

With. 496. *Engl. Bot.* t. 467. *Fl. Dan.* t. 15.

Alga. *Raii Syn.* 52.

β. Fucus marinus, seu Alga marina graminea, minor. *Raii Syn.* 52.

Algoides. *Mich. Ic. ined.* t. 60. f. 2.

γ. Fucus sive Alga marina graminea angustifolia feminifera ramosior. *Raii Syn.* 52.

Algoides. *Mich. Ic. ined.* t. 59?

δ. Potamogeton marinum in utriculis epiphyllispermon minus. *Raii Syn.* 53.

ε. Alga angustifolia vitrariorum. *Raii Syn.* 53.

Angl. Grass-wrack.

In littoribus marinis, æstuariis, vel fossis salis et cœnosis.

Perennis. Fl. Augusto, Sèptembri.

Radix fibrosa. Caules teretes, glabri, geniculati; basi decumbentes, radicales; supernè natantes, foliosi, subramosi, compressiusculi. Folia alterna, petiolata, linearia, obtusiuscula, integerrima, glabra, parum supra basin rimâ longitudinali dehiscentia, atque spadiceum planum,

planum, linearem, exsertentia, uno latere floriferum. Anthera oblonga, sessilis ad latus germinis, sed cum proximâ superiore alternans. Germen sessile, oblongum. Stigmata duo, linearia, acuta, patula. Capsula elliptica, pendula, monosperma.

Varietates ex fide Synopses Raianæ receptas, donec ulterius doceat experientia, ad unam speciem reduxi, collatis iconibus ineditis Michelianis in bibliothecâ meâ, ut et Bankianâ, asservatis. Zosteræ oceanica Linnæi, sui generis planta, nostras nunquam oras attigit. Hujus fructificationem illustravit acutissimus vir Philippus Cavolinus, Neapolitanus, apud Usteri Annal. Botan. fasc. 9. ut etiam Michelius in Ic. ined. t. 58.

II. DIGYNIA.

5. CALLITRICHE. *G. Pl.* 6. *Juss.* 19. *Gærtn.* t. 68.

Cal. nullus. *Petala* duo. *Stigmata* acuta. *Semina* quatuor, compressa, nuda, hinc marginato-alata.

Flores aliquot monoici.

aquatica. 1. CALLITRICHE.

Callitriche aquatica. *Huds.* 439. *Sibth.* 2.

C. verna. *Sp. Pl.* 6. *Witb.* 5. *Relb.* 2. *Abbot.* 1. *Fl. Dan.* t. 129.

Stellaria. *Raii Syn.* 289. *Dill. Giff. App.* 119. t. 6.

Alfine aquis innatans, foliis longiusculis. *Baub. Hist.* v. 3. 777. 2.

Water Star-wort. *Pet. H. Brit.* t. 6. f. 3.

β. *Stel-*


- β. *Stellaria minor et repens.* *Raii Syn.* 289.
Callitriche Plinii. *Column. Ecphr.* 316.
Alfine palustris serpyllifolia. *Ger. em.* 614.
 γ. *Callitriche autumnalis.* *Sp. Pl.* 6. *With.* 6.
Relb. 2. *Abbot.* 2.
Stellaria aquatica foliis longis tenuissimis. *Raii Syn.* 290.
Lenticula palustris angustifolia, folio in apice dissecto. *Loef. Pruff.* 140. *t.* 38.
Long Water Star-wort. *Pet. H. Brit. t.* 6. *f.* 4.
Angl. Water Star-wort.

In fossis, lacubus et stagnis vulgaris.

Annua. Fl. Aprili usque ad Octobrem.

Herba aquis innatans, peractâ impregnatione, submersa. Radices fibrosæ, simplices. Caules filiformes, ramosi, foliosi, glabri. Folia opposita, spatulata, obtusa, integerrima, triplinervia, pellucida, glabra; inferiora linearia, uninervia, emarginata. Flores axillares, solitarii, sessiles, exigui, albi, hermaphroditici masculis et foemineis interdum immixtis. Fructus parvus, sessilis, quadricarinatus.

Planta in γ tota submersa, foliis omnibus uniformibus, linearibus, emarginatis, floribus omnibus hermaphroditicis. Hæc etiam primâ æstate floret.



Classis II.

DIANDRIA.

Stamina 2.

MONOGYNIA.

Stylus 1.

* *Flores inferi, monopetali, regulares.*

6. **LIGUSTRUM.** *Cor. quadrifida. Bacca tetrasperma.*

7. **FRAXINUS.** *Cor. nulla, aut quadripartita. Caps. compressa, disperma.*

** *Fl. inferi; monopetali, irregulares, angiospermi.*

9. **VERONICA.** *Cor. quadripartita, rotata.*

10. **PINGUICULA.** *Cor. ringens, calcarata. Cal. quinquefidus.*

11. **UTRICULARIA.** *Cor. ringens, calcarata. Cal. diphyllus.*

*** *Fl. inferi, monopetali, irregulares, gymnospermi.*

12. **LYCOPUS.** *Cor. subæqualis. Stam. distantia.*

13. **SALVIA.** *Cor. ringens. Filamenta transversè pedicellata.*

**** *Fl. superi.*

8. **CIRCÆA.** *Cor. dipetala. Cal. diphyllus.*

***** *Fl. apetalì, vel tetrapetalì.*

Salicornia. Lepidium ruderaie. Coronopus didyma.

DIGYNIA.

Styli 2.

14. **ANTHOXANTHUM.** *Cal. Gluma uniflora, bivalvis. Cor. Gluma bivalvis, aristata.*

II. DIANDRIA.

I. MONOGYNIA.

6. LIGUSTRUM. *G. Pl.* 9. *Juss.* 106. *Gærtn.*
t. 92.

Cor. quadrifida. *Bacca* supera, bilocularis, tetrasperma.

vulgare. 1. LIGUSTRUM foliis elliptico-lanceolatis obtusis mucronulatis.

Ligustrum vulgare. *Sp. Pl.* 10. *Syst. Veg. Ed.* 14. 56. *Huds.* 3. *With.* 10. *Relb.* 3. *Sibth.* 4. *Abbot.* 2. *Curt. Lond. fasc.* 5. *t.* 1.

Ligustrum. *Raii Syn.* 465. *Ger. em.* 1394. *Fuchsf. Hist.* 480. *Matth. Vulgr.* v. 1. 154. *Camer. Epit.* 89. *Trag. Hist.* 1005. *Swert. Flor. p.* 2. *t.* 38. *f.* 4.

Angl. Privet, Print, or Prim-print.

In sylvis dumetis et sepibus, præcipuè solo glareoso, humidiusculo.

Frutex. Fl. Maio, Junio.

Rami virgati. Folia opposita, subsessilia, atro-virentia, glabra, lanceolata, (subinde elliptica,) integerrima, mucronulata nec acuminata, quandoque per hyemem persistentia. Paniculæ terminales, densæ. Flores lactei, graveolentes. Baccæ nigræ, valdè amaræ, sicut folia et cortex. Aërem fumosum urbium non timet.

7. FRAXINUS. *G. Pl.* 550. *Juss.* 105. *Gærtn.*
t. 49.

Cal. nullus, aut quadripartitus. *Cor.* nulla, aut quadripartita. *Capsf.* supera, bilocularis, superne foliacea, compressa. *Sem.* solitaria, pendula.

Flores aliquot sæminci.

1. FRAX-

1. FRAXINUS foliolis ferratis, floribus calyce *excelsior*.
corollâque destitutis.

Fraxinus excelsior. *Sp. Pl.* 1509. *Huds.* 446.

With. 57. *Relb.* 381. *Sibth.* 18. *Fl. Dan.*

t. 969. *Abbot.* 220.

Fraxinus. *Raii Syn.* 469. *Ger. em.* 1472.

Camer. Epit. 64.

Common Ash Tree. *Hunt. Evel. Sylv. tab. ad*
p. 150.

- β. Ramis pendulis.

Angl. Common Ash. β. Weeping Ash.

In sylvis et sepibus; montosis calcareis gaudet.

- β. Near Gamlingay, Cambridgeshire. *Prof.*
Martyn.

Arbor. Fl. Aprili, Maio.

Excelsa valdeque formosa est arbor, cortice cinereo, gemmis magnis nigris. Flores in paniculis laxis, hermaphroditici, foemineis (subinde numerosissimis) intermixtis. Antheræ magnæ, atro-purpureæ, quibus delapsis hæc arbor, nostratum ultima, frondescit. Folia impari-pinnata, serrata, quandoque simplicia, (conf. *Huds. et With.*) autumnno primis cedens pruinis. Capsulæ citò deciduæ sunt. Semina compressa, pulvere ferrugineo micantia, more Amygdalorum.

8. CIRCÆA. *G. Pl.* 11. *Juss.* 319. *Gærtn. t.* 24.

Cor. dipetala. *Cal.* diphyllus, superus. *Caps.*
bilocularis. *Sem.* solitaria.

1. CIRCÆA caule erecto, foliis ovatis denticu-*lutetiana*.
latis opacis pubescentibus.

Circæa lutetiana. *Sp. Pl.* 12. *Huds.* 10. *With.*

10. *Relb.* 11. *Sibth.* 9. *Curt. Lond. fasc.* 3.

t. 3. *Raii Syn.* 289. *Ger. em.* 351. *Fl. Dan.*

t. 210. *Dicks. H. Sicc. fasc.* 8. 1. *Retz. Obs.*

fasc. 1. 9. *Lob. Ic.* 266.

Angl.

Angl. Common Enchanter's Nightshade.

In locis umbrosis humidis, areis, cœmeteriis.

Perennis. Fl. Junio, Julio.

Radix valdè repens, vix extirpanda. Caulis erectus, parum ramosus, racemo plerumque solitario, terminali. Folia adulta ovata, obscurè viridia, pubescentia, opaca, margine subrepanda, denticulata nec ferrata. Calyx haud membranaceus.

alpina. 2. CIRCÆA caule adscendente, foliis cordatis ferratis nitidis, calyce membranaceo.

Circæa alpina. *Sp. Pl.* 12. *Huds.* 10. *With.* 11. *Dickf. H. Sicc. fasc.* 8. 2.

C. minima. *Column. Ecphr.* v. 2. 79. t. 80.

C. alpina omnium minima. *Ambrosin. Phyt.* 162.

Solanifolia Circæa alpina. *Bauh. Pin.* 168. *Morif. Sect.* 5. t. 34. f. ult.

β. *Circæa alpina.* *Fl. Dan.* t. 256.

Magnitudo *C. lutetianæ*; caulis erectus, racemo subsolitario.

Angl. Mountain Enchanter's Nightshade.

In umbrosis roridis, subalpinis, saxosis, Angliæ borealis. β in nemorosis ubi exuberat umbra.

In Lancashire, Westmoreland and Cumberland.

Mr. Dawson Turner. In several places by the road side from Ulverstone to Hawkeshead, and at Rydall, Westmoreland. *James Crowe, Esq.* At Lowdore water-fall. *Sir T. Frankland, Bart.* In Hamilton Wood, Scotland. *Dr. Hope.* At the foot of mountains about Loch Broom, Ross-shire. *Lightf.*

β. In shady groves by the river side at Matlock bath, Derbyshire. This seems to be what *Mr. Wood* found near Leeds; see *Withering Ed.* 2.

Perennis.

Perennis. Fl. Julio, Augusto.

Radix repens, ut in priori specie. Tota planta minùs pubescit. Caules humiles, basi diffusi, geniculis rubris. Folia tenera, lætè pallidèque virentia, nitida, cordata, dentato-ferrata; petiolis membranaceo-angulatis. Racemi plerumque plurimi, breves, terminales et axillares. Calyx membranaceus, albus, apice rubro.

Mira confusio orta est de hisce speciebus apud Linnæum, cujus characteres omnino lubrici, ne dicam falsi, sunt. Differentias veras in foliis detexere Scopoli et Hallerius consentiente tandem Linnæo.

§ facie et magnitudine gaudet *C. lutetianæ*, etiam racemo subfolitario; at differt colore pallido, foliisque ferratis vel subserratis, petiolis membranaceo-angulatis. Herba ob solum fertilius valde luxurians, forsitan minùs abundè floret, quod et aliis sæpe accidit.

9. VERONICA. *G. Pl.* 12. *Juss.* 99. *Gærtn. t.* 54.
Cor. quadrifida, rotata, laciniâ infimâ angustiore.
Caps. superâ, bilocularis.

* *Spicatæ.*

1. VERONICA spicâ terminali, foliis oppositis *spicata*; obtusiusculis crenato-ferrulatis apice integerimis, caule adscendente simplicissimo.

Veronica spicata. Sp. Pl. 14. *Huds.* 3. *With.* 12.
Relb. 3. *Engl. Bot. t.* 2.

V. spicata recta minor. Raii Syn. 279. *Baub. Hist. v.* 3. 274.

V. recta minima. Ger. em. 627.

V. spicata minor. Vaill. Par. t. 33. *f.* 4.
Angl. Spiked Speedwell.

In pratis et pascuis editioribus calcareis, nec humidis.

Plentiful on Newmarket heath, and Gogmagog hills. On Cavenham heath near Bury. Near Penny bridge, Lancashire. *Mr. Woodward.*

Perennis. Fl. Julio; sæpe ad Septembrem usque.

Caules spithamæi, adscendentes, simplices, teretes, foliosi, sublignosi, pubescentes. Folia opposita, plerumque obtusa, subinde acutiuscula, erenato-ferrulata, apice integerrima, pilosa. Spica erecta, multiflora. Flores saturatè cærulei, fauce barbati.

Variat foliis linearibus et ferè integerrimis. *D. Woodward.*

hybrida. 2. *VERONICA* spicis terminalibus, foliis oppositis ellipticis obtusis inæqualiter crenato-ferratis, caule cretiusculo.

Veronica hybrida. *Sp. Pl.* 14. *With.* 12.

V. spicata β *Huds.* 3.

V. spicata Cambrobritannica, Bugulæ subhirsuto folio. *Raii Syn.* 278. t. 11.

Angl. Welsh Speedwell.

Ad latera montium, rarissimè.

On the side of Craig Wreiddin, Montgomeryshire; Mr. Lhwyd. *Ray.* At Cartmel Wells, Lancashire. *Mr. Crowe and Mr. Woodward.*

Perennis. Fl. Julio.

Præcedenti nimis affinis, sed omni parte duplo major, spicis plerumque plurimis. Caulis et folia insuper magis scabra sunt, profundiusque ferrata; at petiolus alatus etiam in priori et sæpiùs invenitur.

officinalis. 3. *VERONICA* spicis lateralibus pedunculatis, foliis oppositis scabris, caule procumbente.

Veronica officinalis. *Sp. Pl.* 14. *Huds.* 4. *With.*

13. *Relb.* 4. *Sibth.* 4. *Abbot.* 2. *Curt. Lond.*

fasc. 3. t. 1. *Fl. Dan.* t. 248. *Riv. Monop.*

Irr. t. 93. *Dicks.* II. *Sicc. fasc.* 13. 3.

V. mas, supina et vulgatissima. *Raii Syn.* 281.

V. vera et major. *Ger. em.* 626.

V. mas. *Fuchs. Hist.* 166.

Angl. Male, or Common Speedwell.

Vulgaris in pascuis sterilibus arenosis; et ericetis.

Perennis. Fl. Maio, Junio.

Caules latè diffusi. Folia pallida, rigidula, hirsuto-scabra. Spicæ axillares, solitariae, pedunculatae, multifloræ, pilosæ; caulem superantes. Flores dilutè cærulei, venis saturationibus picti.

* * *Corymbo-racemosæ.*

4. VERONICA corymbo terminali pauciflora, *saxatilis*, foliis ellipticis, caulibus diffusis, capsulâ ovatâ quadrivalvi.

Veronica saxatilis. *Suppl.* 83. *Dicks. Crypt. fasc.* 2. 29. *Tr. of Linn. Soc.* v. 2. 288. *With.* 14. *Bauh. Hist.* v. 3. 284. *Scop. Carn.* v. 1. 11.

V. fruticulosa. *Huds.* 4? *Fl. Dan. t.* 342. *Sm.* in *Tr. of Linn. Soc.* v. 1. 191.

V. n. 545 β. *Hall. Hist.* v. 1. 236.

V. tertia fruticans. *Chus. Hist.* v. 1. 347.

V. alpina frutescens. *Morif. sect.* 3. t. 22. f. 5.

V. alpina con foglia di Serpillo. *Pon. Bald.* 181.

V. fruticans serpillifolia. *Ger. em.* 628.

Angl. Blue Rock Speedwell.

In alpinis Scoticis.

On Ben Lawers. *Mr. Dickson.*

Perennis. Fl. Julio.

Caules basi decumbentes, implexi, fruticulosi, ramosi; ramis adscendentibus, simplicibus, foliosis. Folia opposita, parva, elliptica, vel elliptico-lanceolata, obtusa, integerrima, rariùs medio ferrata, obscurè viridia, sub-carnosa, glabra; ad basin ramorum congesta, supernè magis distantia. Flores 3 vel 5 in corymbo terminali, pedunculis, propriâ bractea duplo vel triplo longioribus, insidentes, magni, speciosi, cæruleo-violacei, fauce rubrâ. Calyx quadrifidus, ferè æqualis, obtusus, pubescens. Capsula pubescens, ovata, quadrivalvis.

A V. fruticulosa differt foliis plerumque latioribus, can-
libus magis diffusis, corollâ magnâ, cœruleâ, nec car-
neâ; at præcipuè pedunculis longioribus, paucioribus,
corymbosis, nunquam spicatis.

*fruticu-
losa.*

5. VERONICA corymbo terminali multifloro spi-
cato, foliis elliptico-lanceolatis, caulibus erec-
tis, capsulâ ovatâ quadrivalvi.

Veronica fruticulosa. *Sp. Pl.* 15. *Mant.* 316.

V. frutescens. *Scop. Carn. v. 1.* 19. excluso syn.
Morisoni.

V. n. 545. *Hall. Hist. v. 1.* 235. *t.* 16. *f.* 1.

Angl. Flesh-coloured Shrubby Speedwell.

In humidis subalpinis Scotiis.

On Cruachan, Argylshire. *Rev. Dr. Walker.* Ben
Lawers. *Mr. Brown.*

Perennis. Fl. Julio.

Caules basi ramosi, tortuosi, prostrati, fruticulosi, dein
erecti, striati, spithamæi, simplices, foliosi, teretes, sub-
pubescentes. Folia opposita, elliptico-lanceolata, ple-
rumque integerrima, interdum rarè ferrata, pallidè vi-
ridia, concaviuscula, margine subhirsuta. Corymbus
terminalis, pubescens, subspicatus, et demum in race-
mum longissimum, foliolosum, productus. Pedunculi
floriferi longitudine bractearum, fructiferi sesquilon-
giores. Calyx quadrifidus, ferè æqualis, obtusus, hir-
sutus. Corolla pallidè incarnata lineis purpureis, præ-
cedente minor. Capsula ovata, hirsuta, quadrivalvis.

alpina.

6. VERONICA corymbo terminali subspicato,
foliis ovatis glabris subserratis, calyce ciliato,
caule adscendente simplici.

Veronica alpina. *Sp. Pl.* 15. *Fl. Lapp. Ed.* 2. 7.
t. 9. *f.* 4. *Dicks. Crypt. fasc.* 2. 29. *Tr. of*
Linn. Soc. v. 2. 287. *Fl. Dan. t.* 16. *With.* 14.
Engl. Bot. t. 484.

V. n. 544. *Hall. Hist. v. 1.* 235. *t.* 15. *f.* 2.

Angl. Alpine Speedwell.

In spongiosis irriguis alpium Scoticarum.

On mountains about Garway-moor, and on Ben-Nevis, first found by Mr. Dickson. On the mountains of Badenoch. *Mr. J. Mackay.*

Perennis. Fl. Julio, Augusto.

Caules digitales, adscendentes, simplices, foliosi, basi radicantes. Folia opposita, elliptico-ovata, tenuia, levitèr hirsuta, sæpiùs integerrima, subinde crenata, subtrinervia, obscurè viridia, lucida. Flores exigui, cærulei, in corymbo parvo, tandem in brevem racemum producto. Pedunculi fructiferi etiam elongantur, ut in præcedentibus. Calyx sub-æqualis, obtusiusculus, ciliatus et sæpiùs hirsutus; pili articulati, sicut pubescentia foliorum omnis. Capsula elliptica, emarginata, bivalvis, pubescens, stylo persistente brevi terminata.

7. VERONICA racemo terminali subspicato, foliis *serpyllifolia* ovatis subcrenatis trinerviis glabris, capsulâ *folia* obcordatâ stylo brevior.

Veronica serpyllifolia. *Sp. Pl.* 15. *Huds.* 4.

With. 14. *Relb.* 4. *Sibth.* 5. *Abbot.* 3. *Curt.*

Lond. fasc. 1. t. 3.

V. pratensis minor. *Raii Syn.* 279.

V. minor. *Ger. em.* 627.

V. minima repens. *Riv. Monop. Irr.* t. 99. f. 1.

β. *V. humifusa.* *Dicks. in Tr. of Linn. Soc.* v. 2. 288.

With. 15. *Sym. Syn.* 4.

V. alpina. *Lightf.* 72 (descr. falsa) et 1138.

Angl. Smooth Speedwell. *Paul's Betony.*

In pratis et pascuis. β in alpihus Scoticis.

β on the highest mountains only, under wet shady rocks. *Dicks.*

Perennis. Fl. Maio, Junio.

Herba lucida, subcarnosa, in humidis glabra; in montosis hirta evadit. Caules basi decumbentes et radicantes, ramosi, foliosi, ramis plerumque erectis. Folia petiolata, ovato-subrotunda, magis vel minus crenata,

trinervia. Racemus terminalis, elongatus, laxis, multiflorus, sæpiùs pubescens. Bractæ ellipticæ. Calycis lacinia obovato-ellipticæ, glabriusculæ, æquales. Corolla cæruleo- vel incarnato-albida, striis violaceis. Capsula obcordata, didyma, longitudine styli vel parùm brevior.

Caulis in β prostratus, racemo brevior.

Becabunga.

8. VERONICA racemis lateralibus, foliis ellipticis planis, caule repente.

Veronica Becabunga. *Sp. Pl.* 16. *Huds.* 5. *With.* 15. *Relb.* 5. *Sibth.* 5. *Abbot.* 3. *Curt. Lond. fasc.* 2. t. 3. *Woodw. Med. Bot.* t. 7. *Fl. Dan.* t. 511.

V. aquatica rotundifolia, Becabunga dicta, minor. *Raii Syn.* 280.

Anagallis seu Becabunga. *Ger. em.* 620.

Beccabunga. *Riv. Monop. Irr.* t. 100. f. 1.

Sium. *Fuchs. Hist.* 725.

Sion non odoratum. *Trag. Hist.* 188.

Beccabunga, herba. *Pharmac. Lond.*

Angl. Brooklime.

In rivulis et fossis limpidis frequens.

Perennis. Fl. Julio.

Herba glaberrima, lucida. Caules decumbentes, vel nates, teretes. Folia brevius petiolata, elliptica, obtusa, obsolete ferrata, lætè viridia, subcarnosa. Racemi axillares, oppositi, pedunculati, multiflori. Bractæ lineari-lanceolatae, pedicellis breviores. Calycis lacinia acutæ. Corolla cærulea. Capsula ferè didyma.

Anagallis.

9. VERONICA racemis lateralibus oppositis, foliis lanceolatis ferratis, caule erecto.

Veronica Anagallis. *Sp. Pl.* 16. *Huds.* 5. *With.* 15. *Relb.* 5. *Sibth.* 5. *Abbot.* 3. *Curt. Lond. fasc.* 5. t. 2. *Fl. Dan.* t. 903.

V. aqua-

V. aquatica longifolia media. *Raii Syn.* 280.

Anagallis aquatica major. *Ger. em.* 620.

Angl. Water Speedwell, Long-leaved Brooklime.

In fossis et palustribus.

Perennis. Fl. Julio.

A præcedente discrepat primâ facie foliis lanceolatis, longioribus, acutis, minime ovatis vel ellipticis, cauleque erecto. Racemi insuper longiores et acutiores sunt.

10. *VERONICA* racemis lateralibus alternis : *pe-scutel-*
dicellis divaricatis, foliis linearibus denticu- *lata.*
latis.

Veronica scutellata. *Sp. Pl.* 16. *Syst. Veg.*

Ed. 14. 59. *Huds.* 5. *With.* 16. *Relb.* 6.

Sibth. 5. *Abbot.* 3. *Curt. Lond. fasc.* 5. t. 3.

Fl. Dan. t. 209.

V. aquatica angustifolia minor. *Raii Syn.* 280.

V. palustris angustifolia. *Riv. Monop. Irr.*
t. 96. f. 1.

Angl. Narrow-leaved Marsh Speedwell.

In palustribus et inundatis, solo fabuloso.

Perennis. Fl. Julio, Augusto.

Caules debiles, diffusi. Folia lineari-lanceolata, rariùs integerrima, sæpiùs denticulata, subinde ferrata. Racemi axillares, solitarii, alterni, nunquam oppositi, laxi, divaricati, longitudine varii. Flores parvi, cæruleo-incarnati, striati. Capsula subdidyma.

Variat caule foliisque pubescentibus. *D. C. Sutton.*

11. *VERONICA* racemis lateralibus elongatis fili- *montana.*
formibus paucifloris, foliis ovatis petiolatis
ferratis, caule undique piloso.

Veronica montana. *Sp. Pl.* 17. *Suppl.* 83.

Huds. 6. *With.* 16. *Relb.* 7. *Sibth.* 6. *Ab-*

bot. 4. *Curt. Lond. fasc.* 4. *t.* 2. *Jacq. Austr*
t. 109. *Hoffm. Germ.* *t.* 1.

V. chamædryoides, foliis pediculis oblongis insi-
dentibus. *Raii Syn.* 281.

Alyssum Dioscoridis montanum. *Column. Ecphr.*
v. 1. 286. *t.* 288. bona.

Angl. Mountain Speedwell. Mountain Mad-
wort.

In sylvis humidis opacis, et ad sepes, imprimis solo
calcareo, humidiusculo.

In Charlton Wood, Kent. *Sherard.* In the woody
part of the Devil's ditch on Newmarket heath;
also at Linton, in Gamlingay park and Kingston
wood. *Relb.* Near Worcester. *Dr. Stokes.* Near
Virginia water. *Rev. Dr. Goodenough.* About
Kirkstall Abbey near Leeds, Mr. Aikin; and at
Shortwood, Pucklechurch, Gloucestershire, plen-
tifully, Mr. Swayne; *With.* Very common in
Yorkshire. *Sir T. Frankland, Bart.*

Perennis. Fl. Maio, Junio.

Radix fibrosa. Caules decumbentes, debiles, undique pi-
loso, foliosi. Folia petiolata, ovata, inciso-ferrata, tenuia,
lucida, subpilosa. Racemi axillares, alterni, laxi, pauci-
flori. Flores pallidè cærulei, purpureo picti. Capsula
orbiculato-didyma, compressa, acutè marginata, ciliata.

Chamæ- 12. *VERONICA* racemis lateralibus, foliis ovatis
drys. sessilibus rugosis inciso-ferratis, caule bifa-
riam piloso.

Veronica Chamædrys. *Sp. Pl.* 17. *Syst. Veg.*
Ed. 14. 50. *Huds.* 6. *With.* 16. *Relb.* 6.
Sibth. 5. *Curt. Lond. fasc.* 1. *t.* 2. *Mart. Fl.*
Russ. 1. 66. *Fl. Dan.* 1. 448.

V. Chamædrys sylvestris dicta. *Raii Syn.* 281.
Chamædrys. *Brunf. Herb.* *v.* 1. 125.

C. syl-

C. sylvestris. *Ger. em.* 657.

Angl. Germander Speedwell. Wild Germander.

In pratis, pascuis et ad sepes vulgaris.

Percennis. Fl. Maio.

Caules patuli, basi decumbentes, utrinque lineâ longitudinali pilosâ. Folia sessilia, ovata, inciso-ferrata, margine præcipuè pilosa. Racemi axillares, oppositi, multiflori, caulem superantes. Flores magni, formosi, lætè cærulei, extûs incarnati. Capsula obcordata, parva.

*** *Floribus solitariis.*

13. VERONICA floribus solitariis, foliis ovatis *agrestis*. inciso-ferratis pedunculo brevioribus, caulibus procumbentibus, feminibus urceolatis.

Veronica agrestis. *Sp. Pl.* 18. *Huds.* 7. *With.* 17. *Relb.* 7. *Sibth.* 6. *Curt. Lond. fasc.* 1. t. 1. *Fl. Dan.* t. 449.

V. floribus singularibus, in oblongis pediculis, chamædryfolia. *Raii Syn.* 279.

V. folio chamædryos. *Riv. Monop. Irr.* t. 99. f. 2.

Alfine foliis Triflaginis. *Ger. em.* 616.

A. media. *Fuchs. Hist.* 22.

Angl. Procumbent Speedwell. Germander Chickweed.

In arvis, ruderatis, cultis ubique.

Annua. Fl. ab Aprili usque ad Septembrem.

Caules plures, procumbentes, supernè simplices, teretes, pilosi. Folia sparsa, subsessilia, cordato-ovata, lætè ferrata. Pedunculi axillares, solitarii, uniflori, longitudine foliorum vel longiores, fructiferi recurvi. Calycis lacinia ovato-lanceolata, obtusæ, ciliatæ. Corolla parva, lætè cyanea. Capsula didyma, ventricosa, hirta. Semina utrinque fena, umbilicato-concava, extûs corrugata.

arvensis. 14. VERONICA floribus solitariis, foliis ovatis inciso-ferratis; floralibus lanceolatis pedunculo longioribus, caule erecto.

Veronica arvensis. *Sp. Pl.* 18. *Huds.* 6. *With.* 12.

Relb. 8. *Sibth.* 6. *Curt. Lond. fasc.* 2. *t.* 2.

V. flosculis singularibus cauliculis adhærentibus, *Raii Syn.* 279.

Alfine foliis *Veronicæ*. *Ger. em.* 613.

Alyssum. *Column. Phytob.* *t.* 28.

Angl. Wall Speedwell. Speedwell Chickweed.

In arvis ficcioribus, muris et rudcratis frequens.

Annua. Fl. Maio.

Caulis basi sæpiùs ramosus, erectus, spithamæus. Herba pallidior. Flores sessiles in axillis bractearum lanceolatarum, à foliis planè absimilium, quam ob rem inter spicatas recenseretur, modo non repugnaret affinitas cum præcedenti et sequentibus. Calycis lacinia lanceolata inæquales, acutæ. Corolla pallidè cærulea. Capiula sessilis, erecta, obcordata, compressa. Semina elliptica, complanata, hinc medio ferebiculata.

*hederi-
folia*.

15. VERONICA floribus solitariis, foliis cordatis planis quinquelobis, laciniiis calycinis cordatis, seminibus præcolatis.

Veronica hederifolia. *Sp. Pl.* 19. *Huds.* 7.

With. 17. *Relb.* 8. *Sibth.* 7. *Curt. Lond. fasc.* 2. *t.* 1. *Fl. Dan.* 428.

V. flosculis singularibus, *Hederulæ folio*, *Morsus gallinæ minor* dicta. *Raii Syn.* 280.

V. folio *hederæ*. *Riv. Monop. Irr.* *t.* 99. *f.* 3.

Alfine *hederacca*. *Ger. em.* 616.

Alfines quartum genus. *Fuchs. Ic.* 13.

Angl. Ivy-leaved Speedwell. Small Henbit.

In arvis, rudcratis, cultis ubique.

Annua. Fl. ab Aprili ad Septembrem usque.

Caulis prostrati, simplices. Folia alterna, longius petiolata, reniformi-cordata, quinqueloba, lobo centrali maximo. Pedunculi foliis longiores. Calycis laciniae latè cordatæ, acutæ, ciliatæ. Corolla pallidè cærulea. Capsula didyma, ventricosa. Semina magna, utrinque bina, umbilicato-concava, extus rugosa.

♂ Linn., species revera distincta, gaudet laciniis calycinis ellipticis, obtusis, undique hirtis, nec ciliatis.

16. VERONICA floribus solitariis, foliis super-triphyllis. rioribus digitato-partitis, pedunculis calyce longioribus, seminibus complanatis.

Veronica triphyllus. *Sp. Pl.* 19. *Syst. Veg.* Ed. 14. 60. *Huds.* 7. *With.* 17. *Engl. Bot.* t. 26. *Fl. Dan.* t. 627. *Curt. Lond. fasc.* 6, t. 2.

V. flosculis singularibus, foliis laciniatis, erecta. *Raii Syn.* 280.

Alfine recta. *Ger. em.* 612.

Angl. Finger'd Speedwell. Upright Chickweed.

In arvis arenosis, sed rarius.

In several parts of the sandy tract of country which connects Norfolk and Suffolk. Near Bury. *Sir John Cullum, Bart.* Near Rossington, Yorkshire; Mr. Tosfield. *Huds.*

Annua. Fl. Aprili.

Caulis plerumque ramosus, ramis divaricato-patulis, flexuosis. Folia infima integra, vix lobata; superiora subsessilia, ad basin usque trifida, lobo centrali maximo, obovato, lateralibus plerumque bipartitis. Flores longius pedunculati, erecti, ut et fructus. Calycis laciniae obovato-oblongæ, obtusæ, ciliatæ, inæquales. Corolla saturatè cærulea. Capsula orbiculata, emarginata, compressa. Semina utrinque plurima, obovata, complanata. Herba exsiccatione nigrescit.

verna.

17. VERONICA floribus solitariis, foliis digitato-partitis, pedunculis calyce brevioribus, caule stricto.

Veronica verna. *Sp. Pl.* 19. *Syst. Veg. Ed.* 14. 60. *Huds.* 8. *With.* 13. ~ *Rose Elem. App.* 444. t. 2. f. 1. *Engl. Bot.* t. 25. *Dicks.* *H. Sicc. fasc.* 10. 1. *Fl. Dan.* t. 252.
Angl. Vernal Speedwell.

In arvis arenosis aridis.

Near Bury. *Sir John Cullum, Bart.* At Foulden. Norfolk. *James Crowe, Esq.*

Annua. Fl. Aprili.

Habitu, colore pallido nunquam exsiccatione nigrescente, figurâ seminum, et situ florum, V. arvensi proxima; at foliis profundè divisis, lobo centrali obovato, obtuso ut in præcedente, floralibus etiam tripartitis, abundè discrepat. Caulis vix digiti longitudine, strictus, sæpè ramosus. Calycis laciniae lanceolatae, acutae, subæquales. Capsula obcordata, compressa.

10. PINGUICULA. *G. Pl.* 13. *Juss.* 98.
Gærtn. t. 112.

Cor. ringens, calcarata. *Cal.* bilabiatus, quinquefidus. *Caps.* supera, unilocularis.

lusitana.

1. PINGUICULA nectario obtuso petalo brevior, scapo villoso, capsulâ globosâ.

Pinguicula lusitanica. *Sp. Pl.* 25. *Huds.* *Ed.* 1. 7. *Engl. Bot.* t. 145. *With.* 18. *Cullum.* 6.

P. villosa. *Huds.* 8. *Lightf.* 77. t. 6.

P. flore minore carneo. *Raii Syn.* *281.

Viola palustris, Pinguicula dicta, Lusitanica. *Griseb. Virid.* 84.

Angl. Pale Butterwort.

Ad paludum margines.

In Hampshire, Dorsetshire, Devonshire and Cornwall plentifully. *Huds.* In the Isle of Skye. *Rev. Mr. Stuart.* Near Air, and in the island of Lamash; *Dr. Hope. Huds.* On Croagh Patrick, in the county of Mayo, Ireland. *A. B. Lambert, Esq.* Stoneham Park, Hampshire. *Sir Jos. Banks, Bart.* On many parts of the heath which, beginning at Yellow Ham hill near Dorehester, extends eastward, with little interruption, into Hampshire. In Purbeck, by Sandford bridge near Wareham, and about the midway between Wareham and Woodbury hill; also on Canford heath between Wimborne and Poole, and in several other places about Poole. It is usually found on the borders of bogs, and is abundantly more frequent in these counties than *P. vulgaris*. *Dr. Pulteney.*

Perennis. Fl. Junio, Julio.

Tota planta pubescit. Folia tenella, pallidè virentia, venis rubris picta, margine involuta. Scapi basin versùs imprimis pubescentes, pilis viscidis. Calyx ferè æqualis. Corollæ segmenta obcordata, æqualia; calcar basi inflato-conicum, dein paulò constrictum, apice obtusum, compressum. Germen pubescens. Capsula globosa, glabrata.

2. PINGUICULA nectario cylindraceo acuto longitudine petali, capsulâ ovatâ. *vulgaris.*

Pinguicula vulgaris. *Sp. Pl.* 25. *Huds.* 8. *With.* 18. *Relb.* 8. *Sibth.* 7. *Abbot.* 5. *Engl. Bot.* t. 70. *Dicks. H. Sicc. fasc.* 3. 1. *Fl. Dan.* t. 93.

*P. Giesneri. Raii Syn.** 281. *Bauh. Hist.* v. 3. 546.

P. five Sanicula Eboracensis. Ger. em. 788.

Butterwort. *Pet. H. Brit.* t. 35. f. 7.

Angl.

Angl. Common Butterwort. Yorkshire Sannicle.

In palustribus spongiosis, et ericetis humidis.
Plentiful throughout the north of England, and Scotland, as well as in Norfolk.

Perennis. Fl. Maio, Junio.

Planta glabra. Folia minùs involuta. Scapi glabri, apice tantum subpubescentes. Corolla violacea, limbo inæquali; calcar subulatum, acutum. Capsula ovata.

11. UTRICULARIA. *G. Pl.* 14. *Juss.* 98.

Cor. ringens, calcarata. *Cal.* diphyllus, æqualis, *Capsf.* supera, unilocularis.

vulgaris. 1. UTRICULARIA nectario conico, scapo paucifloro.

Utricularia vulgaris. *Sp. Pl.* 26. *Huds.* 8.

With. 19. *Relb.* 9. *Sibth.* 7. *Abbot.* 5.

Engl. Bot. t. 253. *Fl. Dan.* t. 138.

Lentibularia. *Raii Syn.* *286.

Millefolium palustre galericulatum. *Ger. em.* 828.

Great Bladder-snout. *Pet. H. Brit.* t. 36. f. 11.

Angl. Common Bladderwort, or Hooded Milfoil.

In aquis stagnantibus, paludibus, fossis et piscinis.

Perennis. Fl. Julio.

Herba ramosissima, sub aquâ natans, foliis minimis, fetaceis, alternis, sæpè utriculo compresso, aëre inflato, comitatis. Scapus supra aquam exsertus, erectus, teres, racemo 3 ad 8-floro terminatus. Bractææ solitariae, lanceolatae. Flores speciosi, lutei, lineis rubris. Nectarium incurvum, acutum, labio ferè æquale.

minor. 2. UTRICULARIA nectario carinato brevissimo obtuso.

Utricu-

Utricularia minor. *Sp. Pl.* 26. *Huds.* 9. *With.* 20.

Relb. 10. *Abbot.* 6. *Engl. Bot.* t. 254. *Fl.*

Dan. t. 128. *Schmidel.* *Ic.* t. 21. f. 1.

Lentibularia minor. *Raii Syn.* *286.

Small Bladder-snout. *Pet. H. Brit.* t. 36. f. 12.

Angl. Lesser Hooded Milfoil.

In fossis et paludibus rariùs.

In the river on Hounslow heath; Mr. Dandridge.

Dill. On turf bogs in Westmoreland and Cum-

berland, and near Glastenbury, Somersetshire,

plentifully. *Huds.* On Gamlingay bogs, Cam-

bridgeshire. *Relb.* At Cawston decoy, Norfolk.

Rev. Mr. Bryant. Near Yarmouth. *Mr. Dawson*

Turner. In little ditches on St. Faith's bogs,

near Norwich, along with the *U. vulgaris*.

Perennis. *Fl.* Julio.

A vulgari discrepat herbâ teneriori, at præcipuè floribus minoribus pallidioribusque, et nectario brevissimo, obtuso, vix calcarato.

12. LYCOPUS. *G. Pl.* 15. *Juss.* 111.

Cor. quadrifida: laciniâ unicâ emarginatâ. *Stam.*

distantia. *Sem.* quatuor, nuda, retusa.

1. LYCOPUS foliis finuato-ferratis.

europæus,

Lycopus europæus. *Sp. Pl.* 30. *Huds.* 9. *With.* 20.

Relb. 10. *Sibth.* 8. *Abbot.* 6. *Curt. Lond.*

fasc. 3. t. 2.

L. palustris glaber. *Raii Syn.* 236.

Marrubium aquaticum. *Ger. em.* 700.

Pseudo-marrubium palustre. *Riv. Monop. Irr.*

t. 22.

Angl. Water Horehound. Gypsywort.

Ad ripas rivulorum et lacuum, solo fabuloso.

Perennis. Fl. Julio, Augusto.

Herba inodora; flores quandoque suaveolentes. Affinis Menthæ, etiam foliis subtus resinoso-punctatis. Caulis tetragonus. Folia opposita. Verticilli multiflori, albi. Variat foliis profunde pinnatifidis, magis vel minus villosis, vix unquam omninò glabris.

13. SALVIA. *G. Pl.* 17. *Juss.* 111. *Gærtn. t.* 66.

Cor. inæqualis. *Filamenta* transversè pedicello affixa. *Sem.* quatuor, nuda.

pratensis. 1. SALVIA foliis cordato-oblongis crenatis: summis amplexicaulibus, verticillis subnudis, corollis galeâ glutinosis.

Salvia pratensis. *Sp. Pl.* 35. *Huds.* 10. *With.* 21.

Sibth. 8. *Abbot.* 6. *Engl. Bot. t.* 153.

Sclarea pratensis foliis ferratis flore cæruleo. *Dill. in Raii Syn.* 237.

Horminum sylvestre. *Fuchf. Hist.* 569.

H. sylvestre Fuchsi. *Ger. em.* 769.

H. pratense. *Riv. Monop. Irr. t.* 36.

Angl. Meadow Clary.

In pratis ficcis et ad sepes varius.

Near Cobham in Kent (not Essex), *Raii Syn.* and *Mr. Jacob Rayer.* Common in Surrey and Suffex. *Dr. Stokes.* At Wick-cliffs, Gloucestershire; *Mr. Swayne.* *With.* Between Middleton Stoney and Audley, Oxfordshire. *Sibth.* In a pasture near Ford-End Farm, Bedford; *Rev. T. O. Marsh.* *Abbot.*

Perennis. Fl. Julio.

Planta formosa, corollis magnis, cæruleo-violaceis, fornicatis, conspicua. Verticilli sexflori, bracteis vix calyce longioribus. Folia rugosa, parùm aromatica.

2. SALVIA foliis ferratis finuatis læviusculis, co-
rollis calyce angustioribus. *verbe-*
naca,

Salvia verbenaca. *Sp. Pl.* 35. *Huds.* 10. *With.* 21.

Relh. 10. *Sibth.* 8. *Abbot.* 7. *Engl. Bot. t.* 154.

Curt. Lond. fasc. 6. *t.* 1.

Horminum fylvestre Lavendulæ flore. *Raii Syn.*

237.

H. fylvestre. *Ger. em.* 771.

Angl. Wild English Clary.

In pratis, pascuis, et aggeribus, solo glareoso vel calcareo.

Perennis. Fl. à Junio usque ad Octobrem.

Præcedente minor, sed magis aromatica. Caulis basi decumbens. Flores exigui, violacei, tubo constricto.

II. DIGYNIA.

14. ANTHOXANTHUM. *G. Pl.* 18. *Juss.* 29.

Cal. gluma bivalvis, uniflora. *Cor.* gluma bivalvis, aristata. *Semen* unicum.

1. ANTHOXANTHUM spicâ ovato-oblongâ, *odoratum*,
flosculis subpedunculatis aristâ longioribus.

Anthoxanthum odoratum. *Sp. Pl.* 40. *Huds.* 11.

With. 58. *Relh.* 11. *Sibth.* 18. *Abbot.* 8. *Curt.*

Lond. fasc. 1. *t.* 4. *Schreb. Gram.* 49. *t.* 5.

Mill. Illustr. t. 4. *Mart. Fl. Rust. t.* 23. *Stil-*

lingf. t. 1. *Leers.* 6. *t.* 2. *f.* 1.

Gramen

Gramen vernum spicâ brevi laxâ. *Raii Syn.* 398.
Angl. Sweet-scented Vernal Grass.

In pratis et pascuis vulgaris.

Perennis. Fl. Maio.

Radix fibrosa, subpubescens. Culmi pedales, simplices, tenues, fulcati, glabri, subinde geniculati, supernè nudi. Folia brevissima, plana, acuminata, lætè viridia; caulina brevissima, patentia, vaginâ longissimâ, parùm ventricosâ, fulcatâ, ore fetosâ. Stipula lanceolata, erecta, alba, scariosa. Spica terminalis, solitaria, erecta, acuta, multiflora. Flores bracteolati. Glumæ calycinæ vaginantes, scariosæ, acutæ, interdum pilosæ, carinâ viridi, scabrâ; interior duplo longior, trinervis. Petala subæqualia, ferrugineâ, fetosa, calyce breviora, dorso basin versùs aristata, aristis inæqualibus, longiore geniculatâ.

Nectarium petalis duplo brevius, diphyllum, læve, ovatum. Antheræ lineares. Styli albi, demum exserti. Stigmata filiformia, villosa.

Spicæ, peractâ impregnatione, flavescent. Odor herbæ exsiccatæ ad Asperulam odoratam accedit, Holco odorato levior.

Classis III.

TRIANDRIA.

Stamina 3.

MONOGYNIA.

Stylus 1.

* *Flores superi.*

15. VALERIANA. *Cor.* quinquefida, basi gibba.
Sem. 1.
16. CROCUS. *Cor.* sexpartita, æqualis. *Stigmata* convoluta.
17. IRIS. *Cor.* sexpartita, petalis alternis reflexis.
Stigmata petaliformia.

** *Flores inferi, glumacei. Semen unicum.*

18. SCHOENUS. *Glumæ* paleaceæ, congestæ; exteriores steriles. *Sem.* subrotundum.
19. CYPERUS. *Glumæ* paleaceæ, distichè imbricatæ. *Sem.* imberbe.
20. SCIRPUS. *Glumæ* paleaceæ, undique imbricatæ.
21. ERIOPHORUM. *Glumæ* paleaceæ, undique imbricatæ. *Sem.* lanâ longissimâ cinctum.
22. NARDUS. *Cal.* nullus. *Cor.* Gluma bivalvis.

DIGYNIA.

Styli 2.

* *Calyces uniflori, vagi.*

24. PANICUM. *Cal.* trivalvis; valvulâ tertiâ minimâ.
26. ALOPECURUS. *Cal.* bivalvis. *Cor.* univalvis, apice simplici.
25. PHLEUM. *Cal.* bivalvis, truncatus, acuminatus, sessilis.
23. PHALARIS. *Cal.* bivalvis, valvis carinatis, æqualibus, corollam includentibus.
27. MILIUM. *Cal.* bivalvis, ventricosus, corollâ major. *Stigmata* villosa.
28. AGROSTIS. *Cal.* bivalvis; valvis acutis, corollâ brevioribus. *Stigmata* plumosa.
35. DACTYLIS. *Cal.* bivalvis, compressus; valvâ majori carinatâ.
39. STIPA. *Cal.* bivalvis. *Cor.* aristâ terminali, basi articulatâ.
41. LAGURUS. *Cal.* bivalvis, aristis villosis.
- Arundo epigejos, Calamagrostis et arenaria. Melica.*

* * *Calyces biflori, vagi.*

29. AIRA. *Cal.* bivalvis. *Flosculi* absque rudimento interjecto.
31. MELICA. *Cal.* bivalvis, sub-biflorus; rudimento tertii flosculi interjecto.
30. HOLCUS. *Cal.* bivalvis. *Cor.* aristata. *Flosculo* altero masculino, apetalò.

* * * *Calyces*

* * * *Calyces multiflori, vagi.*

34. BRIZA. *Cal.* bivalvis. *Cor.* ventricosa, valvis cordatis, obtusis. *Semen* corollæ adnatum.
33. POA. *Cal.* bivalvis. *Cor.* valvis ovatis, acutiusculis, muticis.
32. SESLERIA. *Cal.* bivalvis, subtriflorus, aristatus. *Cor.* quinquedentata.
37. FESTUCA. *Cal.* bivalvis. *Spicula* oblonga, teretiuscula; glumis acuminatis.
38. BROMUS. *Cal.* bivalvis. *Spicula* oblonga; glumis sub apice aristatis, inferiore ciliatâ.
40. AVENA. *Cal.* bivalvis. *Cor.* gluma teretiuscula, dorso aristata. *Arista* contorta.
42. ARUNDO. *Cal.* bivalvis. *Flosculi* lanâ persistente cincti.

Dactylis glomerata.

* * * * *Spicati, receptaculo communi scrobiculato.*

43. LOLIUM. *Cal.* monophyllus, fixus, multiflorus.
44. ROTTBOLLIA. *Cal.* monophyllus; subinde bipartitus, subuniflorus. *Rachis* articulata.
47. TRITICUM. *Cal.* bivalvis, solitarius, multiflorus. *Rachis* dentata.
36. CYNOSURUS. *Cal.* bivalvis, solitarius, multiflorus. *Receptaculum proprium* unilaterale, foliaceum.
45. ELYMUS. *Cal.* bivalvis, aggregatus, multiflorus.
46. HORDEUM. *Cal.* bivalvis, ternus, uniflorus.

TRIGYNIA.

Styli 3.

48. MONTIA. *Cal.* diphyllus. *Cor.* monopetala.
Caps. trivalvis, trisperma.
50. POLYCARPON. *Cal.* pentaphyllus. *Pet.* 5.
Caps. trivalvis, polysperma.
49. HOLOSTEUM. *Cal.* pentaphyllus. *Pet.* 5,
erofa. *Caps.* subcylindrica, apice dehiscens.
Tillæa muscosa. *Stellaria media.*

III. TRIANDRIA.

I. MONOGYNIA.

* *Flores superi.*

15. VALERIANA. *G. Pl.* 22. *Juss.* 195.
Gertn. t. 86.

Cal. nullus. *Cor.* monopetala, basi hinc gibba,
supera. *Semen* 1.

- *1. VALERIANA floribus monandris caudatis, *rubra*.
foliis lanceolatis integerrimis.

Valeriana rubra. Sp. Pl. 44. *Huds.* 12. *With.* 65.
Relb. 13. *Sibth.* 20.

V. rubra Dodonæi. Ger. em. 678.

V. marina. Riv. Monop. Irr. t. 3. *f.* 2,
Angl. Red Valerian.

In muris urbium antiquarum haud infrequens, sed
inter rejectamenta hortorum potiùs recensenda.

Frequent in Cornwall and Devonshire. *Huds.*
About Cambridge and Ely. *Relb.* Oxford. *Sibth.*
In Bishopgate-street, Norwich.

Perennis. Fl. Junio, usque ad Septembrem.

Herba lævis, glaberrima, foliis integerrimis, lanceolatis;
superioribus ovatis, longè acuminatis. Flores in co-
rymbo denso, ramosissimo, terminali, graciles, formosi,
saturatè rosei, rariùs albi. Semen papposum, villis
intertextis.

2. VALERIANA floribus triandris dioicis, foliis *dioica*.
radicalibus ovatis; caulinis pinnatis.

Valeriana dioica. Sp. Pl. 44. *Huds.* 12. *With.* 65,
Relb. 13. *Sibth.* 20. *Abbot.* 8. *Curt. Lond.*
fasc. 4. *t.* 3. *Fl. Dan. t.* 687.

V. sylvestris minor. Raii Syn. 200, mas.

V. sylvestris, seu *palustris*, minor altera. *Raii*
Syn. 200. *foemina*.

V. minor. *Ger. em* 1075.

Phu minimum *Matth. Valgr. v.* 1. 38.

Angl. Small, or Marth Valerian.

In pratis humidis et palustribus.

Perennis. Fl. Junio.

Altitudo spithamæa. Folia vix integerrima; in exemplariis Linnæanis etiam manifestè ferrata sunt: radicalia simplicia, ovata seu elliptica; caulina plerumque lyrato-pinnata. Planta mascula minor et tenerior est.

Flores utrique sexui, parco magnitudine, dilute carnei.

Variat floribus hermaphroditicis, quoad structuram saltèm, rariùs utroque organo perfectis.

officinalis. 3. VALEKIANA floribus triandris, foliis omnibus pinnatis: foliolis lanceolatis subuni-
 formibus.

Valeriana officinalis. *Sp. Pl.* 45. *Huds.* 12.

With. 66. *Relb.* 14. *Sibth.* 21. *Abbot.* 8.

Curt. Lond fasc. 6. t. 3. *Woodv. Med. Bot.*

t. 96. *Fl. Dan.* t. 50.

V. sylvestris major. *Raii Syn.* 200. *Ger. em.*
 1075

Phu. Column Phytob. 114.

P. parvum. *Matth Valgr. v.* 1. 37.

P. germanicum. *Fuchs Hist* 857.

β *Valeriana sylvestris major montana.* *Dill. in Raii*
Syn. 200.

V. sylvestris, radix. *Pharmac Lond.*

Angl. Great Wild Valerian.

In paludosis et propter fluvios frequens.

β in sylvis et pascuis montosis.

Perennis. Fl. Junio.

Antecedente major. Caulis quadripedalis. Folia sub-
 æqualiter pinnata, foliolo terminali vix reliquis majori,
 inæqualiter ferrata. Flores carnei. Radix nauseose
 aromatica,

aromatica, in officinis usitatissima. Var. β ob solum minus humidum præstantior, ut et minus virosa. Huic folia angustiora, herba tenerior.

4. VALERIANA floribus triandris, caule dichotomo, foliis lineari-lingulatis obtusis. *Locusta*.

Valeriana Locusta. *Sp. Pl.* 47. *Huds.* 13. *With.* 66.

Relb. 14. *Sibth.* 21. *Abbot.* 8. *Curt. Lond. fasc.* 5. *t.* 4. *Mart. Fl. Rust. t.* 24.

Valerianella arvensis præcox humilis femine compresso. *Raii Syn.* 201.

Lactuca agnina. *Ger. em.* 310.

Fedia olitoria. *Gærtn. v.* 2. 36. *t.* 86.

Angl. Corn Sallad. Lamb's Lettuce.

In arvis et agris ubique.

Annua. Fl. Aprili.

Caulis dichotomus, patulus. Folia cinereo-virentia, pallida, obovato-lanceolata, vel potius lineari-lingulata, sæpius integerrima, minutè ciliata. Flores exigui, cæruleo-albidi.

Variat foliis ferratis, subinde variè laciniatis. Fructus omnibus ferè varietatibus idem est.

16. CROCUS. *G. Pl.* 25. *Juss.* 59.

Cor. sexpartita, æqualis, supera; tubo longissimo.

Stigmata convoluta, erosa. *Spatha* univalvis, radicalis.

*1. CROCUS stigmatе exserto tripartito: *feg-fativus*.
mentis linearibus.

Crocus fativus α . *Sp. Pl.* 50. *With.* 68. *Relb.* 15.
Woodv. Med. Bot. t. 176.

C. officinalis. *Mart. Fl. Rust. t.* 58. *Huds.* 13 α .

C. autumnalis. *Engl. Bot. t.* 343.

Crocus. *Raii Syn.* 374. *Ger. em.* 151. *Trag. Hist.* 763. *Fuchs. Hist.* 441. *Id.* 249.

Crocum. *Matth. Valgr. v. 1. 62, 63. Camer. Epit. 33.*

Crocus, floris stigma. Pharmac. Lond. Angl. Saffron Crocus.

In pratis et pascuis, vix indigena.

Cultivated in Cambridgeshire and Essex, especially about Saffron Walden, and thence dispersed through the neighbouring fields.

Perennis. Fl. Septembri.

Radix bulbosa, solida, depressa, supra sobolifera. Caulis vel scapus planè nullus. Flos in radice sessilis, violaceus, tubo longissimo albo elevatus. Folia paulò post florem oriuntur, linearia, parùm revoluta, saturatè viridia, nervo albo, subtùs complanato. Stamina corollæ breviora, erecta. Stylus longitudine corollæ inter laciniis ejus lateraliter exsertus, dependens. Stigma profundè tripartitum, fulvum, odoratum; laciniis linearibus, margine involutis, apice crenatis.

vernus. 2. *CROCUS* stigmate incluso trifido: lobis cuneiformibus incis.

Crocus vernus. *Mart. Fl. Rust. t. 59. Engl. Bot. t. 344 Jacq. Austr. App. t. 36.*

C. lativus β. *Sp. Pl. 50. With. 63.*

C. officinalis β. *Huds. 13.*

C. vernalis cæruleus. *Deering. Nott. 60.*

C. vernus flore purpureo. *Ger. em. 154.*

Angl. Spring or Garden Crocus.

In pratis variis.

Near Nottingham plentifully. *Deering. Mrs. Sherbrooke. At Eatersca. Prof. Martyn.*

Perennis. Fl. Martio.

Habitus præcedentis, sed folia latiora, minùsque revoluta. Præcipuè verò differt stigmate incluso, erecto, minùs profundè trifido, pallido, inodoro; laciniis cuneatis, margine involutis, apice inæqualiter incis. Variat in hortis flore albo vel luteo.

3. CRO-

3. CROCUS stigma incluso trifido: lobis multi-*nudi-*
fido-laciniatis penicilliformibus, flore aphyllis.*florus.*

Crocus nudiflorus. *Engl. Bot. t. 491.*

C. sylvestris autumnalis. *Dodon. Pempt. 214.*
f. 1.

C. montanus autumnalis. *Ger. em. 154. f. 6.*

Colchicum commune. *Deering. Nott. 57.*

Angl. Naked-flowering Crocus.

In pratis fabulosis inundatis, rariùs.

Between Nottingham castle and the Trent. *Rev.*

Mr Becher. Near Halifax. Rev. Mr. Wood.

Perennis. Fl. Octobri.

Statura præcedentium. Flores omnino aphylli, purpuro-
violacei. Stigma inclusum, erectum, tripartitum, satu-
ratè fulvum, laciniis profundè multifidis, penicillifor-
mibus. Folia erectiora, pallidiora, vix revoluta, nec
apice emarcida. Capsula erecta, elliptica, trifurcata,
Maio perfecta.

17. IRIS. *G. Pl. 27. Juss. 57. Gærtn. t. 13.*

Cor. sexpartita, supera; petalis alternis reflexis.

Stigmata petaliformia.

1. IRIS corollis imberbibus: petalis interioribus *Pseud-*
stigma minoribus, foliis ensiformibus. *acorus.*

Iris Pseud-acorus. *Sp. Pl. 76. Hudsf 14. With. 69.*

Relb. 15. Sibth. 21. Abbot. 9. Engl. Bot. t. 578.

Fl. Dan. t. 494. Curt. Lond. fasc. 3. t. 4.

Woodv. Med. Bot. t. 40. Cord. Hist. 134.

I. palustris lutea. *Raii Syn. 374. Ger. em. 50.*

Acorus. Brunf. Herb. v. 2. 97.

Angl. Yellow Iris. Water Flower-de-luce.

In aquosis ubique.

Perennis. Fl. Julio.

Radix carnosa, horizontalis, depressa, acris, valdè astringens. Folia verticalia, ensiformia, tripedalia. Caulis compresso-teres, lævis, 3—6-florus. Flores lutei, nigro lineati. Petala interiora erecta, canaliculato-linearia, obtusa.

Variat flore pallidè flavo; etiam albo, ex auctoritate Raii.

fœtidissima,

2. IRIS corollis imberbibus: petalis interioribus patentissimis, caule uniangulato, foliis ensiformibus.

Iris fœtidissima. *Sp. Pl.* 57. *Huds.* 14. *With.* 70. *Relb.* 15. *Abbot.* 9. *Engl. Bot. t.* 596.

I. sylvestris quam Xyrim vocant. *Raii Syn.* 375. *Xyris Ger. em.* 60.

Sphatula fœtida. *Fuchs. Hist.* 794. *Trag. Hist.* 904.

Angl. Stinking Iris, or Gladwyn. Roast-beef plant.

In nemorosis et ad sepēs haud frequens.

At Black Notley, Essex. *Ray.* In Charlton wood, and near Hornsey. *Huds.* Several places about Cambridge. *Relban.* In the west of England not uncommon in lanes, *With.* Not rare in woods in Norfolk.

Perennis. Fl, Junio.

Folia obscurè viridia, subinde variegata, manibus trita fœtorem insolitum spargunt. Caulis subtriflorus, foliis vix altior. Flores plumbeo-cærulei, nigro striati. Semina fulva, nitida, globosa.

* * *Flores inferi, glumacei.*

18. SCHOENUS. *G. Pl.* 29. *Juss.* 27.

Glumæ paleaceæ, congestæ, exteriores steriles. *Cor.* nulla. *Semen* unicum, subrotundum, inter glumas,

1. SCHOE-

1. *SCHOENUS* culmo tereti, foliis margine dor- *Mariscus*,
foque aculeatis.

Schoenus Mariscus, *Sp. Pl.* 62. *Huds.* 14.

With. 79. *Reb.* 16.

Cyperus longus inodorus sylvestris. *Raii Syn.* 426.

Ger. em. 29.

C. longus major et elatior, foliis et carina ferratis.

Morif. Sec. 8. t. 11. f. 24.

Angl. Priekly or long-rooted Bog-rush.

In paludibus, subinde maritimis.

On Hinton and Teversham moors, and near Ely,
Relhan. In boggy places by the river Thame
at Dorsthill near Tamworth, Warwickshire, and
by the sea-side between Pensance and Mar-
ketjeu, Cornwall. *Ray*. On St. Faith's Newton
bogs, near Norwich. *Mr. Pitchford*. In a large
pool called Cars y Cromllyn, between Breton
Ferry and Swansea. *Sir Joseph Banks, Bart.*
Ellingham Fen, Norfolk. *Mr Woodward*. So
plentiful on the moors near Cambridge that it
is commonly used in that town for lighting fires,
Mr Crowe.

Perennis. Fl. Julio, Augusto.

Culmus 3 vel 4 pedalis, lævis, foliosus, supernè ramosus.

Folia lineari-lanceolata, longissima, acuminata; basi
vaginantia, integra; margine carin.que acutissimè fer-
rulata. Flores ferruginei, in corymbis terminalibus
lateralibusque; bracteis vaginantibus, alternis.

2. *SCHOENUS* culmo tereti nudo, capitulo ovato, *nigricans*,
involucri diphylli valvulâ alterâ subulatâ
longiore.

Schoenus nigricans. *Sp. Pl.* 64. *Huds.* 15.

With. 90 *Relb.* 16. *Sibth.* 22. *Dicks. Hort.*

Sicc. fasc. 1 1.

Cyperus nigricans. *With.* 78.

Juncus

Juncus lævis minor, panicula glomerata nigricante. *Raii Syn.* 430.

J. capitatus lithospermi femine. *Morif. f.* 8.
t. 10. *f.* 28.

Angl. Black Bog-rush.

In paludibus spongiosis et turfosis.

Perennis. Fl. Junio.

Culmus pedalis, simplicissimus, nudus, glaber. Folia linearia, semiteretia, scabriuscula; basi dilatata, fuscâ, nitidâ, margine tenuiori, vaginantia, culmo breviora. Capitulum subrotundum vel ovatum, e pluribus spiculis fusco-nigricantibus, nitidis, compositum. Involucrum duplex, membranaceum; interius brevissimum; exterius in acumen foliaceum, rectum, capitulo triplo altius, desinens.

compressus.

3. *SCHIOENUS* culmo teretiusculo nudo, spicâ distichâ involucro monophyllo breviora; spiculis multifloris, foliis planis.

Schoenus compressus. *Sp. Pl.* 65. *Huds.* 15.
With. 80. *Relb.* 17. *Leers* 9. *t.* 1. *f.* 1. *Rel.*
Rudb. 3. *f.* 2. *Dicks. Hort. Sicc. fasc.* 3. 2.

Carex uliginosa. *Sp. Pl.* 1381. *Fl. Suec.* 325.
Gramen cyperoides, spica simplici compressa disticha. *Raii Syn.* 425.

Angl. Compressed Bog-rush.

In paludibus turfosis, pratisque spongiosis.

Not uncommon in Norfolk.

Perennis. Fl. Julio.

Radix fibrosa. Culmus semipedalis, teres, hinc parùm complanatus, glaber, striatus, folio solitario ad basin vaginante, cætera nudus. Folia omnia ferè radicata, alterna, culmo breviora, linearia, plana, nervo dorsali carinata, margine scabriuscula. Spica terminalis, erectiuscula, nitidè fusca, e spiculis 5 ad 12 distichis, infimis subremotis, composita. Bractea solitaria, foliacea,
apicē

apicē triangulo, acutiuseulo, marginibus asperiuscula, spicā plerumque longior. Flores (quantū vidi) hermaphroditei. Semen basi fetis cinctum, *Leers*; quod etiam in Sch. glomerato Americæ observatur; hinc utrisque summa cum Eriophoro, quoad characterem, affinitas.

4. **SCHOENUS** culmo tereti nudo, spicā distichâ *rufus*. involucro monophyllo obtuso longiore; spiculis paucifloris, foliis canaliculatis.

Schoenus rufus. *Huds.* 15. *With.* 80. *Dicks.* *Hort. Sicc. fasc.* 10. 6.

S. brunneus. *Soland. Mss. in Herb. Lightf.*

S. ferrugineus. (*nec Linnæi*) *Huds.* 15. *With.* ed. 2. 42. *Lightf.* 86. (excluso syn. *Morisoni*, quod ad *Juncum triglumem* pertinet.)

S. compressi varietas. *Lightf.* 1138. t. 24. f. 2. rudis.

Angl. Brown Bog-rush.

In paludibus Scotiæ.

Isle of Mull. *Rev. Dr. Walker.* Skye. *Huds.* (doubtful if in Westmoreland.) Near Dunglas Castle. *Lightf.*

Perennis. Fl. Julio.

Radix repens, horizontalis. Culmus 4—6-uncialis, rarius elatior, teres omnino, striatus, glaber, et nudus, præterquam ad basin, ubi folia duo abbreviata gerit. Folia semiteretia, lævia, supernè canaliculata; dorso ecarinata, basi dilatatâ, membranaeâ, vaginantia; radicalia culmum æquantia, caulinis duplo longiora. Spica terminalis, nitidè fusca, e spiculis 5 vel 6 distichis, approximatis composita, (rachis subtriangulari. *Yalden.*) Bractea solitaria, foliacea, obtusa, canaliculata, lævis, spicam vix æquans, plerumque multo brevior, quandoque nulla. Spiculæ plerumque bifloræ, glumis tumidiusculis, lævibus, nec striatis. Antheræ lineares, exstantes. Stigmata revoluta. (Semen basi fetosum. *Yalden.*)

Duplicem glumarum ordinem, sub nomine calycis atque corollæ,

corollæ, descripserunt Hudsonus et Solander (in descr. mss. e Yaldeno mutuata), quæ forsitan *glumæ congestæ* Linnæi.

Recentem non vidi. Descriptionem ex exemplaribus Lightfootii et Walkeri ut potui elaboravi.—Stigmata duo invenit Yalden; Hudsonus tria adesse monuit.

S. ferrugineus verus Linnæi in Britannia nondum inventus est.

albus.

5. SCHOFNUS culmo triquetro folioso, floribus fasciculatis, foliis setaceis.

Schoenus albus. *Sp. Pl.* 65. *Hudf.* 16 *With.* 81. *Relb.* 17. *Fl. Dan.* t. 320. *Dicks. Hort. Sicc. fasc.* 1. 2.

Cyperus minor palustris hirsutus, ramiculis albis paleaceis. *Raii Syn.* 47. *Morif. sec.* 8. t. 9. f. 39.

Gramen junceum leucanthemum. *Ger. em.* 30.*

β. *Hudf.* 16.

Floribus fuscis; nec aliam vel minimam inveni differentiam, collatis exemplaribus originalibus Linnæi. Utrisque culmus triquetus est.
“*Quandoque bonus dormitat Homerus.*”

S. fuscus. *Sp. Pl.* 1664. *With.* 80.

Cyperus minor angustifolius palustris, capitulis fuscis paleaceis. *Dill. in Raii Syn.* 427.

Morif. sec. 8. t. 11. f. 40.

Angl. White-headed Bog-rush.

In palustribus spongiosis: frequens in Scotia.

On Felthorpe bogs, and near Heydon, Norfolk.
Rev. Mr. Bryant. Common in Westmoreland and Cumberland. *Mr. Crowe.*

Perennis. Fl. Augusto.

Radix repens. Culmi semipedales, adscendentes, foliosi, triquetri, striati, glabri; supernè ramis floriferis erectis diviti; basi squamis latis, concavis, vaginati. Folia omnia fere caulina, alterna, vaginantia, (vaginâ integrâ,)

tegrâ), linearia, obtusifuscula, parùm recurva, glabra, caulem nunquam superantia. Flores in fasciculis terminalibus erectis, subcorymbose, pedicellati, nivei, post anthesin fuscescentes; in β semper fusci. Bractæe foliis similes, sed breves.

19. CYPERUS. *G. Pl.* 29. *Gærtn. t.* 2. *Juss.* 27.

Glumæ paleaceæ, distichè imbricatæ. *Cor.* nulla.
Semen unicum, imberbe.

1. CYPERUS culmo triquetro folioso, umbellâ *longus*, foliosâ decompositâ; pedunculis nudis, spiculis alternis.

Cyperus longus. Sp. Pl. 67. *Huds.* 17. *With.* 79.
Raii Syn. 425. *Ger. em.* 30. † *Jacq. Ic. rar.* t. 297.

Morif. sec. 8. t. 11. f. 13.

Angl. Sweet Cyperus, or English Galingale.

In paludibus, rarissimè occurrit.

Isle of Purbeck, Dorsetshire; Mr. Newton. *Ray*,

By a little rivulet that runs into Whitford bay, between St. David's town and St. David's head, the only certain British station of this plant, July 25, 1775. *Sir John Cullum, Bart.* Never found near Norwich.

Perennis. Fl. Julio.

Radix tortuosa, longissima, gratè aromatica et astringens.

Culmus bipedalis, triquetus, striatus, angulis lævibus. *Folia* striata, margine scabra, basin (nec medium) culmi vaginantia; floralia tria vel quatuor, umbellam plerumque superantia. *Pedunculi* numerosi, longitudine varii, teretiusculi, striati, glabri; quandoque proliferi. *Spicæ* e spiculis 4 ad 7, alternis, parùm distantibus, bractæolatis, linearibus, multifloris. *Glumæ* xerampelinæ, carinâ virenti.

20. SCIRPUS. *G. Pl.* 30. *Gærtn. t.* 2. *Juss.* 27.

Glumæ paleaceæ, undique imbricatæ. *Cor.* nulla.
Semen unicum.

* *Spica unica.*

palustris. 1. SCIRPUS culmo tereti basi vaginato, spicâ sub-
ovali terminali, glumis acutis, radice repente.
Scirpus palustris. Sp. Pl. 70. *Huds.* 17. *With.* 73.
Relb. 18. *Sibth.* 22. *Engl Bot. t.* 131. *Fl.*
Dan. t. 273. *Rel. Rudb.* 27 f. 2. *Leers t.* 1.
f. 3.

S. Equiseti capitulo majori. *Raii Syn.* 49.

J. minor capitulis *Equiseti.* *Ger. em.* 35. ic. 1631.

Angl. Marsh creep ng Club-rush.

In fossis, paludibus et rivulis frequentissimus.

Perennis Fl. Junio, Julio.

Magnitudine variat. Radix repens, horizontalis. Culmi
numerosi, simplicissimi, erecti, nudi, teretes, glabri,
basi arctè seorsum vaginati, vaginis cylindricis, trun-
catis, integris. Folia nulla, nisi culmos steriles pro
foliis accipias. Spica solitaria, terminalis, crecta, ovata,
nuda, multiflora. Squamæ fuscae, acutæ, margine sca-
rioso albo. (*Semen* fetis quatuor ad basin. *Leers.*)

*multi-
caulis.*

2. SCIRPUS culmo tereti basi vaginato, spicâ
ovatâ terminali, glumis obtusis æqualibus, ra-
dice fibrosâ.

Scirpus. Fl. Dan. t. 167. conf. tab. 287 descript.

S. palustris varietas. *Linn Fl. Lapp. n.* 19 β.

S. multicaulis, *equiseti* capitulis minoribus. *Rel.*
Rudb. 8. f. 2.

S. equiseti capitulis crassioribus et habitioribus,
pumilus et multicaulis. *Rupp. Fl. Jen. ed.*
Hall. 319.

S. caule

S. caule aphylo, spicâ imbricatâ subrotundâ, glumis obtusis. *Hall. Hist. ed. 1. 249?*
Angl. Many-stalked Club-rush.

In paludibus turfosis rarissimè.

On a bog near the house of Mr. M'Kinnon, at Corryhattachan, Isle of Skye, discovered by Mr. John Mackay in 1794.

Perennis. Fl. Julio.

Radix fibrosa, fibris longis, crassioribus, non repens. Culmi numerosi, cæspitosi, laxè patentes, spithamæi, simplicissimi, teretes, glabri, basi præcipuè striati atque arctè seorsùm vaginati, vaginis cylindricis, strictis, truncatis, integris, subacuminatis, sæpiùs duplicatis, at squamis interstinctis nullis. Folia nulla. Spica ovata, vel elliptica, ferruginea, nuda, multiflora. Squamæ omnes æquales, imbricatæ, ovatæ, obtusæ, atro-purpureæ vel fuscae, apice subscariosæ, carinâ obsoletâ, viridi. Flosculorum unus vel alter, ad basin spicæ, sæpè viviparus et divaricatus evadit.

3. SCIRPUS culmo tereti striato basi vaginato *cæspitofus*.
 multiplici-squamoso, spicâ terminali, glumis
 exterioribus maximis.

Scirpus cæspitosus. Sp. Pl. 71. Hudf. 17. With. 73. Relb. 18. Rel. Rudb. 28. f. 1.

S. montanus capitulo breviori. Raii Syn. 429.

Juncus parvus montanus, cum parvis capitulis luteis. Baub. Hist. v. 2. 525. f. 2.

Angl. Scaly-stalked Club-rush.

In ericetis turfosis haud rarus.

Perennis. Fl. Julio.

Radix fibrosa, cæspitosa. Culmi numerosi, erecti, simplicissimi, altitudine valdè variantes, teretes, æquales, glabri, striati, basi vaginati, vaginâ unâ vel alterâ, laxâ, foliaceâ, striatâ, integrâ, apice obliquâ, in mucronem brevem obtusum desinente, squamis obvallatâ numerosis, membranaceis, albido-fuscis, striatis, concavis, vaginato-

vaginato-imbricatis, extûs minoribus, persistentibus. Spica parva, rufa. Glumæ duæ exteriores (parùm inæquales) longitudine totius spicæ, mucrone foliaceo, obtuso, (feminiferæ. Semen basi setosum. *Stokes.*)

pauciflorus.

4. SCIRPUS culmo tereti striato basi vaginato, spicâ terminali pauciflorâ glumis exterioribus longiore.

Scirpus pauciflorus. Lightf. 1078. Hudf. 648. With. 74. Sibth. 23.

S. Bæothryon. Linn. Suppl. 103. Ehrh. Phyt. n. 31. procul dubio.

S. minimus, spica brevior squamosa spadicea, Scheuchz. Agr. 564. t. 7. f. 19.

S. caule tereti, spicâ nudâ pauciflorâ, glumis calycinis longiore. Hall. Hist. n. 1335. v. 2. 176. Davall.

Angl. Chocolate-headed Club-rush.

In palustribus, præcipuè montosis.

On the highland mountains. *Lightf.* In Ellingham fen, Norfolk. *Mr. Woodward.* Plentiful on Poringland heath, near Norwich. *Mr. Crowe.*

Perennis. Fl. Augusto.

Radix cæspitosa, nigricans. Culmi suberecti, simplicissimi, 3—6 unciales, teretes, graciles, striati, glabri, basi vaginati, vaginâ strictâ, muticâ, æquali. Folia radicalia, culmo similia et sæpe altiora, basi pari modo vaginata. Spica parva, rufo-fusca, subquadriflora, glumis duabus inferioribus subæqualibus, ellipticis, spicâ brevioribus, muticis.

Distinctissima a præcedente, quâcum a Linnæo et Hudsono confunditur, squamarum radicalium defectu ferè totali, vaginis strictis, obtusis, muticis, glumis infimis spicâ brevioribus, foliorumque radicalium præsentia.

Annon folia quæ memoravimus potius culmi steriles dicenda sunt?

5. SCIRPUS culmo quadrangulo, basi vaginâ mu- *acicu-*
ticâ, spicâ ovatâ acutâ terminali, glumis exte- *laris,*
rioribus majoribus.

Scirpus acicularis. Sp. Pl. 71. excluso syn.

Scheuchzeri. Hudf. 18. Sibth. 23. Pollich.

v. 1. 40. Fl. Dan. t. 287.

S. minimus, capitulis equiseti. Raii Syn. 429.

Cyperus acicularis. With. 78.

Juncellus minimus capitulis equiseti. Moris.

sect. 8. t. 10. f. 37. Pluk. Phyt. t. 40. f. 7.

Angl. Least Club rush.

In ericetis humidis, inundatis et palustribus rarius.

On Epping Forest, near Wanstead. *Mr. Edward*

Forster, jun.

Perennis. Fl. Augusto.

Radix fibrosa, cæspitosa. Culmi tenuissimi, simplicissimi, erectiusculi, 2 vel 3-unciales, exactè quadranguli, vix striati, angulis lævibus, basi vaginâ retusâ, muticâ, tenuissimâ, arcuè vestiti. Folia pauca, culmo conformia sed breviora. Spica exigua, ovata, acuta, 5 vel 6-flora, ferruginea. Glumæ imbricatæ, nec distichæ, carinatæ, acutæ, margine membranaceo, pellucido, duæ inferiores vix majores, parùm obtusiores, subindè steriles.—
(Semina imberbia. *Haller, Linnæus.*)

Auctores omnes,* præter Hudsonum, culmum teretem incautè descripsere, quod erroneum.

6. SCIRPUS caule folioso flaccido natante, pedun *fluitans.*

culis alternis nudis, spicis solitariis termina-
libus

Scirpus fluitans. Sp. Pl. 71. Hudf. 18. With. 74.

Sibth. 23. Engl. Bot. t. 216. Dickf. Hort.

Sicc. fasc. 3. 4.

S. equiseti capitulo minori. Raii Syn. 431.

Scheuchz. Agrost. 365. t. 7. f. 20.

Gramen juncum clavatum repens, foliis et ca-

pitulis Psyllii. *Moris. sect. 8. t. 10. f. 31.*

L. 2

Angl.

Angl. Floating Club-rush.

In fossis et stagnis.

On Putney and Hounslow heaths. *Huds.* In several pits on the heath between Farnham and Godalmin. *Mr. Crowe* and *Mr. Dickson*. On St. Faith's Newton bogs, near Norwich.

Perennis. Fl. Julio, Augusto.

Radix exigua, fibrosa. Caules teretes, striati, longissimi, natantes, geniculis radicales, basi gracillimi, internodiis elongatis; supernè paulò incrassati, geniculis approximatis; foliosi. Folia linearia, basi vaginantia, flaccida, natantia, in fasciculis alternis, solitariis ad nodos. Pedunculi axillares, circa partem caulis superioris, solitarii, teretes, striati, nudi, recti, biunciales. Spica terminalis, parva, ovata, virens, glumis duabus exterioribus majoribus, mucronatis. Stigmata duo.

In exsiccatis aliquando magis luxuriat.

* * *Culmo tereti polystachyo.*

lacustris. 7. SCIRPUS culmo tereti nudo, paniculâ cymosâ decompositâ terminali, spiculis ovatis.

Scirpus lacustris. *Sp. Pl.* 72. *Huds.* 19. *With.* 75. *Relb.* 19. *Sibth.* 23.

S. palustris altissimus. *Raii Syn.* 428.

Juncus maximus seu Scirpus. *Baub. Theatr.* 178.

J. lævis maximus. *Morif. sec.* 8. t. 10. f. 1.

β. *Juncus* five *Scirpus* medius. *Baub. Theatr.* 181. *Raii Syn.* 428.

γ. *Juncus aquaticus medius*, caule carinato. *Raii Syn.* 428.

Angl. Bull-rush; β. Lesser Bull-rush.

In aquis purioribus, fluviorum sinibus, stagnis et lacubus frequens.

Perennis. Fl. Julio.

Radix repens, horizontalis, valida. Culmi recti 4 vel 5 pedales, nudi, teretes, glabri, intus spongiosi, basi vaginis

ginis pluribus, alternis. Panicula cymosè decomposita, ramis valdè inæqualibus, compressis, marginibus superne simbriatis. Bractææ exteriores duæ, lanceolatae, acutæ, spathacæ, paniculâ sæpiùs breviores; interiores plurimæ, minores. Spicæ congestæ, ovatae, ferrugineo-rutilantes. Glumæ concavæ, carinatæ, mucronatæ, margine simbriatæ, subinde fissæ. Semina hinc plana, basi setis 5 vel 6.

Variat longitudine bractearum, culmique altitudine; β paniculâ minori; (γ culmo hinc complanato. *Huds.*)

8. SCIRPUS culmo tereti nudo, capitulis glomeratis pedunculatis sessilibusvè, involucro di-*Holo-*
phylllo inæquali, foliis canaliculatis. *schoenus.*

Scirpus Holoschoenus. Sp. Pl. 72. Huds. 19.
With. 75. Dickf. Dr. Pl. 2.

S. maritimus, capitulis rotundioribus glomeratis.
Raii Syn. 429.

Scirpoides maritimum, capitulis sparsis glomeratis. Scheuchz. Agrost. 371. t. 8. f. 2.

Juncus maritimus capitulis rotundis. Moris. sec. 8.
t. 10. f. 17. Pluk. Phyt. t. 40. f. 4. Rel.
Rudb. 22. f. 1. et 25. f. 3.

J. acutus maritimus alter. Park. Theatr. 1194.

† β . *S. australis, culmo tereti nudo, capitulo laterali, bracteâ reflexâ, foliis canaliculatis. Syst.*
Veg. ed. 13. 85.

Juncus minor acutus maritimus prolifer. Pluk.
Phyt. t. 40. f. 5.

γ . *S. romanus, culmo tereti nudo, capitulo laterali*
conglobato, bracteâ reflexâ. Sp. Pl. 72.
Huds. 19. With. 76. Jacq. Austr. t. 448.

Scirpoides acutum maritimum, capitulo glomerato solitario. Scheuchz. Agr. 373. t. 8. f. 6.

S. maritimum, capitulis non omnibus in pediculis insidentibus, foliis et scapo supra paniculam

culam carinam ferentibus viridem. *Mich.*

Gen. 52. *Seguier. Veron.* v. 3. 73.

Angl. Round cluster headed Club-rush.

In palustribus, præcipuè maritimis, sed rariùs.
 β nondum inveni.

On Braunton Boroughs, Devonshire. *Ray. Rev.*
Dr. Goodenough. In Somersetshire and Hamp-
 shire. *Huds.* γ . In marshes near Throgmorton,
 Worcestershire. *Huds.*

Perennis. Fl. Augusto ad Novembrem.

Radix cæspitosa. Culmi erectiusculi, 2—3 pedales, te-
 retes, glabri, tenuissimè striati, basi membranâ lacerâ
 vaginati. Folia radicalia, linearia, acuta, pungentia,
 glabra, ecarinata, supra canaliculata fossilâ nivea. Ca-
 pitula sessilia vel pedunculata, pedunculis angulatis,
 basi vaginatis, subindè ramosis, fusca, c spiculis nume-
 rosis, obtusis, squamis interstinctis concavis, acutis,
 striatis, margine pubescentibus. Bractæ duæ, in-
 æquales, foliis omninò consimiles, quarum superior
 continuatio videtur culmi, longitudine valdè varians,
 biuncialis vel pedalis, recto-incurva; inferior multò
 brevior, magis vel minùs divaricata. Florum structura
 vix nota est. Semina ex auctoritate Scheuchzeri im-
 berbia videntur, stigma trifidum.

Involucrum superius, ut et inferius, lineâ albâ canalicu-
 latum, mucronato-pungens, more foliorum, demonstrat
 in h'c etiam fructificationem verè terminalem ut (pace
 Halleri) in præcedente descripsimus.

Tres species conjungo collatis exemplaribus originalibus
 Linnæi, Jacquinii, Seguieri, aliorumque, monente ipso
 Linnæo in manuscriptis recentioribus.

In β et γ culmus tenuior est, capitula pauciora, aliquan-
 tulò majora. Folia *S. romani* non viderat Linnæus
 cum ex Horto Upsaliensi *S. australem* in *Syst. Veg.*
 ed. 13. descripsit.

setaceus. 9. SCIRPUS culmo nudo setaceo, spicis latera-
 libus subgeminis sessilibus ebracteatis.

Scirpus

Scirpus setaceus. Sp. Pl. 73. (nec Mant. 2. 321.)

Huds. 20. With. 76. Relb. 19. Sibth. 24.

Fl. Dan. t. 311. Leers. 10. t. 1. f. 6.

S. foliaceus humilis. Raii Syn. 430.

Juncellus omnium minimus. Moris. sec. 8. t. 10.

f. 23.

Angl. Least Club-rush.

In glareosis inundatis, et arenosis maritimis.

Annua. Fl. Julio, Augusto.

Radix fibrosa, caespitosa. Culmi numerosi, bi- vel tri- unciales, setacei, striati, glabri. Folia culmis similia sed breviora, vaginâ cylindricâ striatâ. Spicæ plerumque geminæ, sæpè solitariae, quandòque ternæ vel qua- ternæ, paulò infra apicem culmi lateralitèr erumpunt, bractea infernè nullâ suffultæ, ovatæ, glabræ, e glumis subæqualibus, fuscis, nervo concinnè viridi carinatis et mucronatis. (Semina imberbia. *Leers.*)

Descriptio in Mantissa 321 ad aliam *Scirpi* speciem spectat e Lusitania, et Capite Bonæ Spei delatam, quæ forsitan *Cyperus minimus* Linnæi. Conf. *Rotboll. Ic. 47. t. 15.* *Scirpus supinus* Linnæi reverà ab utrisque distinctus est.

*** *Culmo triquetro, panicula nuda.*

10. *SCIRPUS* culino triquetro recto nudo acu- *triqueter.*
minato, spicis lateralibus sessilibus peduncu-
latisve.

Scirpus triqueter. Mant. 1. 29.

α. Spicis maximâ ex parte pedunculatis, numerosis,
culmi apicem quandòque attingentibus.

S. triqueter. With. 76.

S. mucronatus β, triqueter. Huds. 20.

Juncus acutus maritimus, caule triquetro
maximo molli, procerior nostras. *Raii Syn.*
428. *Phuk. Phyt. t. 40. f. 2.*

β. Spicis ferè omnibus sessilibus conglomeratis.

S. mucronatus. With. ed. 2. 50. Huds. 20. α.

Juncus acutus maritimus caule triquetro, rigido,

mucrone pungente. *Dill. in Raii Syn.* 429.

Pluk. Phyt. t. 40. *f.* 1.

J. acutus maritimus caule triangulo. *Bauh.*

Theatr. 175. *Moris. sect.* 8. *t.* 10. *f.* 20.

Rel. Rudb. 22. *f.* 2. 3.

Angl. Triangular Club-rush or Bull-rush.

In littoribus maritimis, et ad ripas fluviorum.

In ditches by the side of Acle dam, Norfolk,

Mr. Crowe. On the banks of the Thames above and below London. β In Jersey. *Dillenius.*

Perennis. Fl. Augusto.

Radix fibrosa, cæspitosa. Culmi erecti, tripedales, nudi, triquetri, lateribus planis angulisque lævibus, intus spongiosi, apice acuto, parum inflexo; (in var. β maritimâ firmiores. *Dill.* Spicæ laterales, bracteatae, squamis aliquot membranaceis interstinctæ, sessiles pedunculatae, pedunculis triangularibus, subinde ramosis, culmi apicem quandocumque æquantibus. Glumæ badiæ, ovatae, carinatae, mucronatae, margine dilatato, lacero.

Scirpus mucronatus Linnæi, *Dicks. Dr. Pl.* 26, nondum in Britannia obvius, discrepat lateribus culmi concaviusculis, acumine reflexo, glumis magis numerosis, striatis, viridibus, margine ferrugineo, integro, nec dilatato, lacero. Huc, ex auctoritate Herbarii Sherardiani, spectat synonymon Scheuchzeri, *tab.* 9. *f.* 14, mucrone quamvis in tabulâ non reflexo.

**** *Culmo triquetro, paniculâ foliaceâ.*

*mariti-
mus.*

II. *SCIRPUS* culmo triquetro, paniculâ conglobatâ foliaceâ terminali, glumis mucronatis lacero-trifidis.

Scirpus maritimus. *Sp. Pl.* 74. *Huds.* 21.

With 77. *Relb.* 20. *Curt. Lond. fasc.* 4. *t.* 4.

Engl. Bot. t. 542.

Gramen cyperoides palustris panicula sparsa.

Raii Syn. 425.

G. aquaticum cyperoides vulgatus. *Ger. em.* 22.

Cyperus longus inodorus latifolius, spicis tumidioribus. *Morif. sect. 8. t. 11. f. 25.*

β. *Cyperus rotundus littoreus*. *Ger. em. 31. inodorus. Raii Syn. 426.*

C. rotundus inodorus aquaticus septentrionalis.
Morif. sect. 8. t. 11. f. 9.

Angl. Salt-marsh Club-rush. β Round-rooted Bastard Cyperus.

In paludibus marinis, et ad ripas æstuariorum.

Perennis. Fl. Julio, Augusto.

Radix repens, in β nodosa. Culmi erecti, pedales vel tripedales, triquetri, striati, angulis asperulis, foliacei. Folia linearia, acuta, longissima, margine scabra; inferiora vaginantia, alterna; floralia sessilia, longitudine valdè inæqualia. Spicæ terminales inter folia floralia, badia, glomeratæ, sessiles pedunculatæve, pedunculis triquetris, simplicibus, ovatæ; subinde cylindricæ longissimæ. Glumæ nitidæ, fusco-ferruginæ, ovatæ, carinatæ, longiùs mucronatæ, apicc laceræ. Antheræ flavæ. Semina triangularia, hinc plana, basi setâ plerumque solitariâ.

12. SCIRPUS culmo triquetro folioso, cymâ fo-*sylva-*
liacæâ terminali; pedunculis nudis suprade-*ticus.*
compositis, spicis confertis.

Scirpus sylvaticus. Sp Pl. 75. Hudsf. 21. With. 77.

Relb. 24. Fl. Dan. t. 307. Leers. 10. t. 1. f. 4.

Cyperus gramincus. Raii Syn. 426. miliaceus.
Ger. em. 30.

C. longus miliaceus inodorus locustis plurimis brevioribus. Morif. sect. 8. t. 11. f. 15.

Angl. Wood Club-rush, or Millet Cyperus-grass.

In nemorosis humidis minùs frequens.

In Pembrokehire, Warwickshire, and Essex. *Ray.*
At Ditchingham (*Mr. Stone*), and other parts of
Norfolk. In Charlton wood, Kent.

Perennis.

Perennis. Fl. Julio.

Radix fibrosa. Culmus solitarius, erectus, triqueter, foliosus, glaber. Folia latiuscula, margine scabra; radicalia densè cæspitosa; caulina vaginantia, alterna; floralia sessilia. Cyma multiflora, supradecomposita, pedunculis triquetris, scabris, alternis, bracteis lanceolatis, membranaceis. Spicæ parvæ, solitariae vel congestæ, ovatæ, fusco-virides. Glumæ carinatae, obtusiusculæ, indivisæ. Semina fetis cincta, ipso semine longioribus.

21. ERIOPHORUM. *G. Pl.* 30. *Gærtn. t.* 2. *Juss.* 27.

Glumæ palcaceæ, undique imbricatæ. Cor. nulla. Semen unicum, lanâ longissimâ cinctum.

vagina- 1. ERIOPHORUM culmis vaginatis teretibus, *tum.* spicâ solitariâ, glumis scariosis.

Eriophorum vaginatum. *Sp. Pl.* 76. *Huds.* 22. *With.* 71. *Curt. Lond. fasc.* 4. *t.* 10. *Rel. Rudb.* 29. *Fl. Dan. t.* 236. *Dicks. Hort. Sicc. fasc.* 4. 3.

Juncus alpinus cum cauda leporina. *Raii Syn.* 436. *Bauh. Hist. v.* 2. 514. *f.* 2.

Gramen junceum, seu Juncus alpinus capitulo lanuginoso. *Morif. sect.* 8. *t.* 9. *f.* 6.

Angl. Single-headed Cotton-grass. Hare's-tail rush.

In ericetis paludosis turfosis.

Frequent in the alpine counties. Near Croydon. *Sherard.* On Blackstone edge, Lancashire, and on Bawsey bottom near Lynn, Norfolk.

Perennis. Fl. Martio.

Radix densè cæspitosa. Culmi obsolete triquetri, vel teretiusculi, læves, vaginâ alterâ aphyllâ ad medium, alterâ ad basin vaginati, vaginâ inflatâ, membranacâ, obtusâ, mucronulatâ. Folia omnia radicalia, basi membranacâ vaginantia, triquetro-linearia, striata, subrecurva. Spica terminalis, erecta, solitaria, ovata, subinde bracteata, primo

primo vere erumpens, e glumis numerosis, lanceolatis, scariosis, cærulescentibus, margine albo. Antheræ inclusæ, flavæ. Peraetâ inflorescentiâ, lana exferitur longissima, nivea, usque ad mediam æstatem persistens.

2. ERIOPHORUM culmis teretibus, foliis planis, *polystachion*,
spicis pedunculatis.

Eriophorum polystachion. *Sp. Pl.* 76. *Dickf.*

Tr. of Linn. Soc. v. 2. 289. *Hort. Sicc.*

fasc. 4. 1. *Engl. Bot. t.* 563. *Leers.* 11. *t.* 1.

f. 5. *With.* 72.

Linagrostis panicula minore. *Vaill. Par. t.* 16.

f. 2.

Gramen tomentosum, panicula sparsa. *Morif.*

sect. 8. *t.* 9. *f.* 1.

Angl. Broad leaved Cotton-grass.

In paludosis Angliæ et Scotiæ.

In Northamptonshire, Bedfordshire, Yorkshire, Cumberland, and very common in Scotland; first determined by Mr. Dickson.

Perennis. Fl. Aprili. *Herb. Lightf.*

Radix fibrosa, nec repens. (*Dickf.*) Culmus erectus, teres, striatus, foliosus, glaber. Folia culmo longè breviora, plana, lineari-lanceolata, lævia, apicem versùs acutè carinata, basi vaginantia, alterna; floralia duo vel tria, basi membranaceo-dilatata. Spicæ plurimæ, pedunculatæ, ovatæ, glumis lineari-lanceolatis, scariosis, post florescentiam, elongatis pedunculis, cernuæ, lanâ exfertâ albâ.

Variat pedunculis ramosis,

3. ERIOPHORUM culmis teretibus, foliis cana- *angustifolium*,
liculatis apice triquetris, spicis pedunculatis.

Eriophorum angustifolium. *Dickf. Tr. of Linn.*

Soc. v. 2. 289. *Hort. Sicc. fasc.* 4. 2. *Engl.*

Bot. t. 564. *Hoffm. Pl. Germ.* 19. *With.* 72.

E. poly-

- E. polystachion.* *Huds.* 21. *Relb.* 20. *Sibth.* 24.
Curt. Lond. fasc. 4. *t.* 9.
Linagrostis. *Raii Syn.* 435.
L. paniculâ majore. *Vaill. Par. t.* 16. *f.* 1.
Angl. Common Cotton-grass.

In paludosis turfosis et cœnosis frequens.

Perennis. Fl. Aprili.

Radix repens. Culmus teres, tenuior, striatus, glaber, subunifolius. Folia culmo parùm breviora, semiteretia; canaliculata, lævia, apice triquetra, basi vaginantia; floralia duo vel tria, linearia, plana, longitudine varia. Spicæ subternæ, pedunculatæ, ovatæ, glumis fuscis, margine scariosis, post florescentiam erectiusculæ, lanâ longissimâ, albâ.

alpinum. 4. **ERIOPHORUM** culmis nudis angulatis, spicâ solitariâ erectâ pappo brevior.

Eriophorum alpinum. *Sp. Pl.* 77. *Transf. of Linn. Soc. v.* 2. 290 & 356. *Fl. Dan. t.* 620. *Dicks. Hort. Sicc. fasc.* 8. 3. *Engl. Bot. t.* 311. *Witb.* 72.

Juncus alpinus bombycinus. *Scheuchz. Prod.* 27. *t.* 8. *f.* 1.

Angl. Alpine Cotton-grass.

In palustribus turfosis alpinis Scotiæ.

On a moss three miles east of Forfar, in the shire of Angus; found by Mr. Brown and Mr. Don in 1791. *Tr. of Linn. Soc.*

Perennis. Fl. Aprili, Maio.

Radix repens. Culmi numerosi, erecti, 4—9-unciales, triquetri, subindè tetragoni, angulis scabris, superne nudi. Folia vaginantia, basin culmi vestientia, linearia, carinata, obtusa, margine scabra, culmo longè breviora, vaginâ tubulosâ, truncatâ. Spica solitaria, erecta, parva, ovata. Glumæ ovatæ, carinatæ, flavo-ferrugineæ; exterior in mucronem foliaceum desinens,
 Semen

Semen acutè triquetrum, lanâ candidâ, spicâ duplò longiore, erectâ, vestitum.

22. NARDUS. *G. Pl.* 30. *Juss.* 33.

Calyx nullus. *Cor.* gluma bivalvis.

1. NARDUS spica setacea recta secunda. *stricta.*

Nardus stricta. Sp. Pl. 77. *Huds.* 22. *With.* 71.

Relb. 21. *Sibth.* 33. *Leers.* 11. *t.* 1. *f.* 7.

Engl. Bot. 290. *Dicks. Hort. Sicc. fasc.* 8. 4.

Mart. Fl. Rust. t. 27.

Gramen sparteum juncifolium. *Raii Syn.* 393.

G. sparteum capillaceo folio, minus, erectum.

Morif. sect. 8. *t.* 7. *f.* 8.

Spartum nostras parvum Lobclio. *Ger. em.* 41.

ic. 1631.

Angl. Mat-grafs.

In ericetis fabulosis humidiusculis.

Perennis. Fl. Julio.

Radix e fibris crassis, pubescentibus. Culmi parùm incurvi, semipedales, rigidi, subangulosi, læves. Folia radicalia, densè cæspitosa, vaginis inflatis, striatis, albidis, nitidis, squamis muticis brevioribus obvallatis; supernè divaricata, angustissima, acuta, tetragona, minutissimè pubescentia, subglaucæ. Spica terminalis, solitaria, gracilis, floribus secundis, cinereis, glumis aristatis, scabris.

II. DIGYNIA.

23. PHALARIS. *G. Pl.* 32. *Gærtn. t.* 80. *Juss.* 29.

Cal. bivalvis, carinatus, longitudine æqualis, uniflorus, corollam includens.

*canari-
ensis.*

* 1. PHALARIS paniculâ ovatâ spiciformi, glumis calycinis navicularibus integris, corollâ quadrivalvi.

Phalaris canariensis. Sp. Pl. 79. *Ait. Hort. Kew. v.* 1. 85. *Huds.* 23. *With* 113. *Sibth.* 33. *Leers.* 18. *t.* 7. *f.* 3 +. *Schreb. Gram.* 83. *t.* 10. *f.* 2. *Mart. Fl. Rust. t.* 17.

Phalaris. Ger. em. 86. major semine albo. *Dill. in Raii Syn.* 394. *Morif. sect.* 8. *t.* 3. *f.* 1. *Angl.* Manured Canary-grass.

In ruderatis et ad vias, vix indigena.

Annua. Fl. Junio—Augusto.

Radix fibrosa, albida. Culmi plures, bipedales, striati, foliacei, geniculis fuscis. Folia lanceolata, molliuscula, parùm scabra; caulina vaginâ striatâ, inflatâ, apice constrictâ. Spica terminalis, erecta, densa, ramosa, speciosa. Glumæ calycinæ albidae, lineis viridibus pictæ, latius carinatæ, naviculares, integræ, plerùmque pilosæ, carinâ tamen imberbi, hiantes, corollâ duplò longiores. Corolla exterior e glumis duabus lanceolatis, carinatis, glabris, minoribus, calyci oppositis: interior duplò longior, valvula exterior sericeo-villosa, compressa, alteram interiorem, glabram, parùm brevior, arctè amplectens. Semen ovatum, compressum, nitidum.

arenaria. 2. PHALARIS spicâ ovato-lanceolatâ obtusâ, glumis exactè lanceolatis ciliatis, culino basi ramoso.

Phalaris

Phalaris arenaria. *Huds.* 23. *With.* 113. *Engl.*
Bot. t. 222.

P. phleoides β . *Ait. Hort. Kew.* v. 1. 86.

Phleum arenarium. *Sp. Pl.* 88. *With.* 62.

Gramen typhinum maritimum minus. *Raii*
Syn. 398. *Pluk. Phyt. t.* 33. *f.* 8.

Angl. Sea Canary-grass.

In arenosis maritimis, nec non ericetis editioribus.
On Swaffham and Newmarket heaths. *Engl. Bot.*

Annua. Fl. Junio.

Radix fibrosa, fibris simplicibus pubescentibus. Culmi plures, nisi solo sterilissimo, divaricati, foliosi, supernè nudi, teretes, purpurei, politi, semipedales. Folia vaginantia, lanceolata, glaucescentia, margine scabriuscula, vaginis inflatis, striatis, glabris. Spica terminalis, solitaria, ovata vel ovato-lanceolata, sub-simplex, vix paniculata vel in lobos divisibilis, multiflora, obtusa. Calyx exactè ovatus, e glumis duabus æqualibus, lanceolatis, compressis, trinerviis, mucronulatis, dorso ciliatis, margine membranaceis. Corolla bivalvis, retusa.

3. PHALARIS paniculâ sub-cylindricâ spiciformi, *phleoides*.
glumis lineari lanceolatis glabriusculis margine interiore retuso, culmo simplici.

Phalaris phleoides. *Sp. Pl.* 80. *With.* 114. *Fl.*

Dan. t. 531. *Engl. Bot. t.* 459. *Ait. Hort.*

Kew. v. 1. 86. α . excluso synonymo Hudsoni.

Gramen typhinum, spicâ conoide striatâ, culmo violaceo. *Barrel. Ic. t.* 21.

Angl. Cat's-tail Canary-grass.

In campis elatis arenosis.

In a field at Narborough, Norfolk. *Mr. Crowe* and
Mr. Woodward. In Cambridgeshire, *Lyons.*
Herb. Banks.

Perennis. Fl. Julio.

Radix fibrosa, fibris glabris. Culmus simplex, sæpè solitarius,

tarius, basi præcipuè foliosus, teres, glaber, purpureus, nitidus, pedalis vel sesquipedalis. Folia vaginantia, lineari-lanceolata, margine scabra, vaginis striatis, parùm inflatis, scabriusculis. Panicula terminalis, solitaria, densa, ovato-cylindrica, acutiuscula, omninò spiciformis, sed in lobos ramosos divisibilis, multiflora. Calyx ovato-linearis, sub apice aliquantulùm contractus, e glumis duabus æqualibus, lineari-lanceolatis, compressis, trinerviis, mucronulatis, summitate subdivergentibus, margine membranaceo, apice, Phleorum more, retuso, dorso nervisque aliquando pubescentibus. Corolla bivalvis, lanceolata.

Gramen Hudsono, ut videtur, ignotum, a Phleo suo paniculato, ut et a præcedenti specie, distinctissimum, quod e descriptionibus nostris facillè intelligi potest.

24. PANICUM. *G. Pl.* 32. *Gærtn. t. 1. Juss.* 29.
Digitaria. *Juss.* 29.

Cal. trivalvis, uniflorus; valvulâ tertiâ minimâ.
Semen corollâ persistente, cartilagineâ, corticatum.

verticillatum. 1. PANICUM spicâ verticillatâ; spiculis quaternis, involucellis unifloris bifetis scabris, culinis diffusis.

Panicum verticillatum. Sp. Pl. 82. *Huds.* 24.
With. 114. *Curt. Lond. fasc. 4. t. 6.*

Gramen paniceum spicâ asperâ. *Raii Syn.* 394.
simplici. *Morif. sect. 8. t. 4. f. 11.*

G. geniculatum. Ger. em. 15. *f. 1.*

Angl. Rough Panick-grass.

In arvis humidis rariùs.

Between Putney and Roehampton, and near Chelsea. *Ray.* Battersea fields. *Curtis.*

Annua. Fl. Junio, Julio.

Radix fibrosa. Culmi plures, diffusè patentes, foliosi, glabri, bipedales. Folia margine scabra, stipulâ intrafoliaceâ, brevissimâ, ciliari. Spica luteo-virens, solitaria,

taria, erecta, composita; spiculis verticillatis, subquaternis, brevibus, patentibus, verticillis inferioribus remotioribus. Flores ovati, pedicellati. Glumæ calycinæ carinata, concavæ. Bractææ binæ plerumque ad singulum florem, setaceæ, scabræ, flore duplo longiores.

2. PANICUM spicâ cylindricâ; spiculis confertis, *viride*, involucellis bifloris multifetosis scabriusculis, corollâ leviusculâ.

Panicum viride. *Sp. Pl.* 83. *Huds.* 24. *With.* 115.

Curt. Lond. fasc. 4. *t.* 5. *Leers.* 13. *t.* 2. *f.* 2.

P. Crus-galli. *Fl. Dan.* *t.* 852!

Gramen paniceum spicâ simplici lævi. *Raii Syn.* 393.

G. panici effigie spicâ simplici. *Ger. em.* 17.

Angl. Green Panick-grass.

In arvis, præcipue arenosis, sed minùs frequens.

Near Guildford, and in Battersea fields. *Hudson.*

About Norwich.

Annua. Fl. Julio.

Radix fibrosa. Culmi plures, patentes, geniculati, scabri.

Folia lato-lanceolata, acuminata, aspera, stipulis e pilis densissimis, brevibus. Spica viridis, hinc subinde purpureascens, composita; spiculis approximatis, subalternis, brevibus, erecto-patentibus. Flores ovati, pedicellati. Glumæ calycinæ ovatæ, concavæ, vix carinatæ, nervosæ. Corolla oculo armato scabriuscula, non vero transversè rugosa ut in *P. glauco*, necque, ut e differentia specifica Linnæana intelligendum videtur, nervosa. Bractæa e fetis numerosis, scabriusculis, plerumque biflora, longitudine varia. (Calyx quadrivalvis. *Stokes.*)

3. PANICUM spicâ decompositâ; spiculis alternis *Crus-* conjugatisque subdivisis, calycibus aristatis hispidis, rachi subquingulari. *galli.*

Panicum Crus galli. *Sp. Pl.* 83. *Huds.* 24.
With. 115. *Curt. Lond. fasc.* 4. t. 8. *Leers.* 13.
 t. 2. f. 3.

P. vulgare. *Ger. em.* 85.

P. sylvestre Herbariorum. *Park.* 1154.

Gramen paniceum spicâ divisâ. *Raii Syn.* 394.
Morif. sec. 8. t. 4. ord. 1. f. 15 & 16.

Angl. Loose Panick-grafs.

In arvis humidis rariùs.

About Martha's Chapel, near Guildford. *Huds.*

About Battersea and Putney, in moist places.
Lightf. Herb.

Annua. Fl. Julio.

Radix fibrosa, cæspitosa. Culmi plurimi, validi, erecti, bipedales, læves. Folia scabra, exstipulata. Spica purpurascens, spiculis subsecundis, subdivisis & ramosis, rachi communi angulatâ, angulis 3—5, scabris. Glumæ calycinae angulatæ, tuberculis setiferis scabræ, aristatæ, aristis longitudine mirè variantibus, scabris. Corolla lævissima, nitida, apice mucronata & subpubescens.

Inter hoc et *Panicum Crus Corvi* vix solidam inveni differentiam.

sanguinale.

4. *PANICUM* spicis digitatis basi interiore nodosis, floribus geminis muticis, vaginis foliorum punctatis.

Panicum sanguinale. *Sp. Pl.* 84. *Huds.* 25.
With. 115. *Curt. Lond. fasc.* 4. t. 7. *Leers.* 14.
 t. 2. f. 6. *Schreb. Gram.* 119. t. 16. *Mart.*
Fl. Rust. t. 78.

Gramen dactylon latiore folio. *Raii Syn.* 399.
Herb. Buddle. *Morif. sect.* 8. t. 3. ord. 1. f. 2.

Ischæmon vulgare. *Ger. em.* 27.

Angl. Cock's-foot Panick-grafs.

In arvis, sed rariùs.

Near Guildford, with the preceding. *Huds.* Plentiful

tiful about Elden in Suffolk, *Ray*; and at Witchingham near Norwich, (but not found there at present.) In Battersea fields, and in sandy fields between Brandon and Moundeford, Norfolk; also near Henham, between Beccles and Saxmundham, Suffolk. *Lightf. Herb.*

Annua. Fl. Julio.

Radix fibrosa. Culmi numerosi, decumbentes, geniculati, ramosi, pedales vel sesquipedales, striati, glabri. Folia læviuscula, quandoque pilosa, vaginis propriis breviora. Vaginæ striatæ, subinde pilosæ, tuberculis adpersæ raris, minutis, apice subexaratis, subinde piliferis. Spicæ terminales, umbellatæ, erecto-patentes, in plantâ luxurianti numerosæ et dein parum remotæ vel alternæ. Rachis partialis indivisa, alata, flexuosa, multiflora, basi interiore quandoque, nec semper, nodosa. Flores secundi, per paria alterni, altero pedicellato. “Glumæ calycinæ sæpe quaternæ” (*Curtis*), “hinc verum Panicum, nec distincti generis, censeri oportet; conjungit nempe cum Panicis Digitariam Halleri et Jusseui.” *Stokes*. Ipse in speciminibus indigenis duas tantum inveni, in exoticis aliquando tres; hæ valde inæquales, nervosæ, corollinis oppositæ.

5. PANICUM spicis digitatis basi interiore villosis, *dactylon*. floribus solitariis, calycibus æqualibus corollæ alternis, sarmentis repentibus.

Panicum Dactylon. *Sp. Pl.* 85. *Huds.* 25. *With.* 116. *t.* 21. *Dicks. Dr. Pl.* 53. *H. Sicc. fasc.* 11. 1.

Gramen repens cum paniculâ graminis Mannæ. *Raii Syn.* 399. *Herb. Buddle.*

G. dactylon folio arundinaceo minus. *Baub. Theatr.* 114. *Morif. sect.* 8. *t.* 3. *f.* 4.

G. dactiloides radice repente. *Ger. em.* 28.

Angl. Creeping Panick-grass.

In arenosis maritimis Cornubiæ.

About Pensance and Marketjeu. *F. Borone.*

Perennis. Fl. Julio, Augusto.

Radix sublignosa, repens, geniculata, longissima, glabra, radiculis ramosissimis, lævibus, sarmenta exferens longissima, prostrata vaginis brevifoliis vestita. Culmi 3 vel 4 unciales, basi geniculato decumbentes, foliosi, striati, glabri. Folia linearia, acutissima, hirta, vaginisque striatis albidis. Stipulæ fetosæ. Spicæ 3 vel 4, umbellatæ, purpurascens, erecto-patentes. Rachis partialis flexuosa, angulata, nec alata. Flores solitarii, numerosi, secundi. Calyx e glumis binis, lanceolatis, carinatis, æqualibus, lateralibus. Corolla calyci alterna, ovata, lævis.

25. PHLEUM. *G. Pl.* 33. *Gærtn. t.* 1. *Juss.* 29.

Cal. bivalvis, truncatus, acuminatus, sessilis, uniflorus, corollam includens.

pratense. 1. PHLEUM spicâ cylindricâ longissimâ, glumis dorso ciliatis aristâ longioribus.

Phleum pratense. *Sp. Pl.* 87. *Huds.* 25. *With.* 117.

Relb. 23. *Sibth.* 34. *Leers.* 17. *t.* 3. *f.* 1.

Schreb. Gram. 102. *t.* 14. *Mart. Fl. Rust.* *t.* 5.

Gramen typhinum majus, seu primum. *Raii Syn.* 398. *Ger. em.* 11.

G. typhoides maximum, spicâ longissimâ. Moris. sect. 8. *t.* 4. *ord.* 3. *f.* 1. *Bauh. Theatr.* 49.

β. Huds. 26.

Gramen typhinum minus. *Raii Syn.* 398. *Ger. em.* 12.

G. typhoides medium five vulgatissimum. Moris. sect. 8. *t.* 4. *ord.* 3. *f.* 2.

γ. Huds. 26.

Phleum nodosum. *Sp. Pl.* 88. *Relb.* 23. *Sibth.* 34. *Leers.* 17. *t.* 3. *f.* 2.

Gramen nodosum spicâ parvâ. *Raii Syn.* 398. *Bauh. Theatr.* 20.

G. typhoides minus, nodosum. Moris. sect. 8. *t.* 4. *ord.* 3. *f.* 3.

Angl.

Angl. Common Cat's-tail-grass. Timothy-grass.

In pratis et pascuis humidis. β in locis sterilibus et ad vias. γ in apricis, soloque sterili rariùs inundato.

Perennis. Fl. a Junio ad Octobrem usque.

Radix fibrosa, in β parum tuberosa, in γ bulbosa, bulbis ovatis. Culmi erecti, 3 vel 4-pedales, striati, foliosi, superne nudi; in β et γ basi decumbentes, et breviores. Folia scabriuscula, vaginis cylindricis, longis, striatis; stipula brevissima, obtusa, membranacea. Spica solitaria, erecta, cylindrica, obtusa, e floribus numerosissimis, confertis; in α 4 vel 5-uncialis, in β et γ vix biuncialis. Glumæ calycinae in omnibus virides, nervis, margineque membranaceo interiore, albis, carinà ciliatis, aristis brevibus, subdivariatis. Antheræ purpureæ.

“Variat in insulâ Monâ floribus viviparis.” *D. Davies.*

2. PHLEUM spicâ ovato-cylindraccâ, aristis longitudine glumarum.

Phleum alpinum. *Sp. Pl.* 88. *Fl. Lapp. Ed.* 2. 20. *Dicks. Tr. of Linn. Soc.* v. 2. 288. *Engl. Bot. t.* 519. *Fl. Dan. t.* 213. *With.* 119.

Gramen typhoides alpinum, spicâ brevi, densâ, et veluti villosâ. *Scheuchz. Agrost.* 64. *Prod.*

t. 3.

Angl. Alpine Cat's-tail-grass.

In pratis alpinis Scotiæ.

On mountains near Garway moor. *Dickson.*

Perennis. Fl. Julio.

Radix tuberosa, subrepens. Culmus solitarius, adscendens, pedalis, subinde sesquipedalis, foliosus, superne nudus, striatus, glaber. Folia margine scabra, stipulis brevissimis subdeltoidibus, nec retusis; inferiora vaginis brevibus, cylindraceis; superiora vaginis inflatis,

folio longioribus. Spica vix uncialis, ovato-cylindrica, obtusa, atro-purpurea. Glumæ calycinæ longius ciliatæ, subdivaricatæ, nervis viridibus, aristis rectis, glumarum fere longitudine, purpurcis.

paniculatum.

3. PHLEUM paniculâ cylindricâ spiciformi, glumis nudis sursum incrassatis, culmo ramoso.

Phleum paniculatum. *Huds.* 26, exclusis synonymis forsân omnibus. *With.* 117. *Relb. Suppl.* 1. 8.

Phalaris paniculata. *Ait. Hort. Kew.* v. 1. 87. *Sibth. Ox.* 34.

Ph. aspera. *Retz. Obs.* 4. 14.

Gramen typhinum paniceum utriculatum amplo utriculo. *Barrel. Ic. t.* 28. *f.* 2.

Angl. Panicked Cat's-tail-grass.

In pratis, campis elatis et muris.

Below King's Weston near Bristol, and on Newmarket heath. *Huds.* Near Bourn bridge. *Mr. Crowe.* Corn-fields near Newmarket. *Relban.*

Annua. Fl. Julio.

Radix fibrosa, parva, annua proculdubio. Culini plures, ramosi, subpedales, adscendentes, geniculati, foliosi, glaberrimi, basi attenuati. Folia plurima, erecta, asperiuscula, nuda, vaginis longis, subinflatis, striatis, nudis; stipulæ membranacæ, laceræ. Panicula solitaria, recta, 2—4 uncialis, virens, cylindrica, densa, acutiuscula, omnino spiciformis, sed in lobos compositos divisibilis, tactu aspera, e floribus numerosissimis, imbricatis, rigidis, constans. Calyx cuneiformis, oculo armato scaber, superne gibbosus, e glumis duabus æqualibus, cuneatis, gibbis, angulosis, obtusissimè truncatis, mucronulatis, subdivergentibus; margine interiori rectangulo, vix membranaceo; carinâ imberbi. Corolla bivalvis, inæqualis, calyce brevior. Anthæræ parvæ, subrotundæ, albidæ.

Gramen adhuc valde obscurum, certè Phleum, nec Phalaris, quod bene descripsit Hudsonus, quamvis synonyma
et

et ipse et omnes male intellexere. Halleri et Bauhini citationes conscribere, prout dubias, nolui. Barrelicri t. 54. est Phalaridis (vel forsitan Crypsæos *Hort. Kew.* species) Gallo-provinciæ indigena, quam e Cel. Gerardo habui.—Phleum paniculatum in Horto Parisiense colitur sub nomine Phalaridis asperæ Jusseui. *Herb. Thouin. N. 211.*

4. PHLEUM paniculâ spicatâ, glumis linearibus *crinitum*.
basi subventricosus hispidis aristatis: aristis capillaceis longissimis.

Phleum crinitum. *Schreb. Gram. 151. t. 20. f. 3. With. 64.*

Alopecurus aristatus. *Huds. 28.*

A. monspeliensis. *Sp. Pl. 89. With. 121.*

A. maxima Anglica. *Raii Syn. 396. paludosa. Moris. sect. 8. t. 4. ord. 2. f. 3.*

Gramen alopecurum majus, spicâ virescente divulsâ, pilis longioribus. *Barrel. Ic. t. 115. f. 2.*

β. *Huds.*

Alopecurus paniceus. *Sp. Pl. 90. With. 121.*

Cynosurus paniceus. *Sp. Pl. ed. 1. 73.*

Gramen alopecurum minus, spicâ virescente divulsâ. *Barrel. Ic. t. 115. f. 1.*

Angl. Bearded Cat's-tail-grass.

In pratis maritimis udis rariùs.

Near Portsmouth, and in ditches on the Essex coast. *Ray.* Near Cley, Norfolk, *Mr. Humphrey.*
Herb. D. Rose.

Annua. Fl. Julio, Augusto.

Radix fibrosa, parva. Culmi plures, erectiusculi, geniculati, teretes, lævissimi, glabri, foliosi. Folia striata, margine nervisque scabra, vaginis longis striatis, glabris, stipulâ membranaceâ, laccrâ, nervosâ, extus scabrâ. Panicula erecta, densa, spiciformis, e spiculis composita ternis vel quaternis, secundis, lobatis et ramosis.

Flores numerosi, congesti, pedicellati, secundi, parvi. Glumæ calycinæ æquales, lineari-lanceolatae, compressæ, pubescenti-hispidae, superne virides, margine membranaceo, retuso, albo; arista terminalis alba, scabra, glumâ duplo longior. Corolla bivalvis; valvulâ exteriori majore, brevius aristatâ, alteram imberbem amplectente.

Varietas β minor et quodammodo tenerior est, nullam tamen differentiam in floribus inveni, examinatis exemplariis ipsius Linnæi.

Optimè ut videtur ad Phleum retulit hanc speciem Celebrerrimus Schreberus.

26. ALOPECURUS. *G. Pl.* 33. *Gærtn. t.* 1. *Juss.* 29:

Cal. bivalvis, uniflorus. *Cor.* univalvis.

pratensis. 1. ALOPECURUS culmo erecto lævi, spicâ sublobatâ, glumis calycinis villosis basi connatis.

Alopecurus pratensis. *Sp. Pl.* 88. *Huds.* 27.

With. 119. *Curt. Lond. fasc.* 5. *t.* 5. *Relb.* 24.

Sibth. 34. *Schreb. Gram.* 133. *t.* 19. *f.* 1.

Leers. 15. *t.* 2. *f.* 4. *Mart. Fl. Rust.* *t.* 6.

Gramen Alopecuro simile glabrum cum pilis longiusculis in spicâ. *Raii Syn.* 396.

G. alopecuroides majus. *Ger. em.* 10.

G. alop. spicâ longiore medium vulgare, folio lanuginoso. *Morif. sect.* 8. *t.* 4. *f.* 8. ex amanuensis errore in *Mant.* 324 excluditur.

Angl. Meadow Fox-tail-grass.

In pratis et pascuis ubique, pecoribus gratum.

Perennis. Fl. Maio.

Radix fibrosa. Culmus bipedalis, erectus, foliosus, lævis.

Folia glabriuscula, glaucescentia, stipulâ brevi, subpubescente. Spica simpliciuscula, vix paniculata, densa, erecta, biuncialis, mollis, multiflora. Glumæ omnes fere æquales, lanceolatae, compressæ, albæ, lineis utrinque duabus, viridibus, longitudinalibus, pictæ; calycinæ ad carinam præcipue sericeo-villosæ, muticæ: gluma

gluma corollina solitaria, glabra nisi carinæ apicem versus, basi aristata; arista geniculata, flore duplo longior, nuda.

2. **ALOPECURUS** culmo erecto scabriusculo, *agrestis*.
 spicâ simplicissimâ attenuatâ, glumis nudiusculis basi connatis carinâ dilatatis.

Alopecurus agrestis. *Sp. Pl.* 89. *Huds.* 27.
With. 119. *Relb.* 24. *Sibth.* 35. *Schreb.*
Gram. 140. t. 19. f. 2. *Leers.* 16. t. 2. f. 5.
Mart. Fl. Ruß. t. 22.

A. myosuroides. *Huds. ed.* 1. 23. *Curt. Lond. fasc.* 2. t. 7.

Gramen myosuroides majus, spicâ longiore, aristis rectis. *Raii Syn.* 397.

G. alopecuroides spicâ longâ tenuiore. *Morif. sect.* 8. t. 4. f. 12.

G. alopecurinum minus. *Ger. em.* 11.

- β. *G. myosuroides minus*, spicâ brevior, aristis recurvis. *Raii Syn.* 397. *Herb. Buddle.*

Varietas parva, fusca, c. situ aprico sterili.

Angl. Slender Fox-tail-grass.

In arvis et ad vias.

Annua. Fl. Julio.

Radix fibrosa, parva, annua ni fallor. Culmus sesquipedalis, erectus, foliosus, apice nudo, scabriusculo. Folia supra scabra, stipulâ lanceolatâ, pubescente. Spica fere triuncialis, erecta, gracilis, acuta, multiflora, subinde purpurascens. Glumæ subæquales, præcedente majores, minusque pubescentes, albo viridique variæ, nervis prominentibus; calycinæ basi villosæ, connatæ, carinâ dilatâ subpubescente, muticæ; gluma corollina glabra, basi aristata; arista geniculata, flore duplo longior, oculo armato scabra, exsiccatione recurva.

3. **ALOPECURUS** culmo erecto, spicâ simpli- *bulbosus*.
 cissimâ attenuatâ, glumis calycinis distinctis villosis, radice bulbosâ.

Alope-

Alopecurus bulbosus. Sp. Pl. 1665. With. 120.

A. geniculatus β . *Huds. 27.*

Gramen myosuroides nodosum. Dill. in Raii Syn. 397. t. 20. f. 2.

G. typhinum phalaroides, pilosâ spicâ, aquaticum bulbosum. Barrel. Ic. t. 699. f. 1.

Angl. Bulbous Fox-tail-grass.

In aquosis maritimis, rariùs.

Salt marshes near Yarmouth. *Mr. Woodward.*

Cardiff marshes, Glamorganshire; and near the Aust passage. *Lightf. Herb.*

Perennis. Fl. Julio.

Radix bulbosa, cæspitosa, fibras e parte inferiori emittens, membranâ fuscâ, striatâ, tunicata. Culmus solitarius, vix pedalis, simplicissimus, erectus, basi parum decumbens, foliosus, binodis, striatus, glaber. Folia glabra; radicalia pauca, brevia; caulina fere linearia, patentia, longitudine vaginarum. Stipula brevis, striata. Spica sesquiuncialis, erectiuscula, simplicissima, gracilis, acuta, multiflora. Glumæ calycinæ parum inæquales, acutæ, muticæ, basi omnino discretæ, nervis carinâque (non dilatatâ) villosæ; corollina emarginata, basi aristata ut in prioribus.

Species procul dubio distincta, ab *A. geniculato* diversissima, spicâ et structurâ florum *A. agrestis* magis quidem affinis.

*genicu-
latus.*

4. **ALOPECURUS** culmo adscendente geniculato, spicâ sublobatâ cylindricâ, glumis retusis pilosis.

Alopecurus geniculatus. Sp. Pl. 89. Huds. 27.

With. 120. Relb. 25. Sibth. 35. Curt. Lond. fasc. 5. t. 6. Leers. 16. t. 2. f. 7. Mart. Fl. Rust. 1. 97.

Gramen aquaticum geniculatum spicatum. Raii Syn. 396.

G. fluviatile spicatum. Ger. em. 14.

Gramen aquaticum spicatum. *Park. Theatr.*
1275.

G. alopec. fluviatile geniculatum procumbens.
Morif. sect. 8. t. 4. f. 15.

β. G. fluviatile album. *Dill. in Raii Syn. 396.*

γ. Radice bulbosâ. *Huds. 27. With. 60.*

Angl. Floating Fox-tail-grass.

In aquosis et stagnis frequens. γ in muris, soloque
sterili, exsiccato.

Perennis. Fl. Julio.

Radix e fibris longissimis, simplicissimis. Culmi natantes,
longissimi, geniculati, geniculis inferioribus radican-
tes, apice adfurgentes, foliosi, ramosi, glabri. Folia glabri-
uscula. Stipula albida, tenuissima. Spica cylindrica,
brevis, obtusiuscula, in lobos divisibilis, multiflora, pur-
purascens, in β albida. Glumæ præcedentibus duplo
minores, subæquales, obtusissimæ; calycinæ longius
pilosæ; corollina crenata, glabra, basin versus aristata;
arista geniculata, longitudine varia. Radix in ficcis
bulbosa evadit, culmo erectiusculo: hanc varietatem
cum A. *bulboso* vero confudit Hudsonus.

27. MILIUM. *G. Pl. 33. Juss. 29.*

Cal. bivalvis, uniflorus, tumidus. Cor. bivalvis,
brevissima. Stigmata villosa.

1. MILIUM floribus paniculatis diffusis muticis. *effusum.*

Milium effusum. *Sp. Pl. 90. Huds. 29. With. 122.*

Relb. 25. Sibth. 35. Curt. Lond. fasc. 4. t. 12.

Leers. 18. t. 8. f. 7.

Gramen miliaceum. *Raii Syn. 402. Ger. em. 6.*

G. sylvaticum, paniculâ miliaceâ sparsâ. *Morif.*
sect. 8. t. 5. f. 10.

Angl. Millet-grass.

In nemorosis humidis frequens.

Annua.

Annua. Fl. Junio, Julio.

Radix fibrosa. Culmi plures, erecti, tripedales, foliosi, glabri, striati. Folia latiuscula, plana, margine tantum scabra. Stipula ovata, lacera. Panicula terminalis, erecta, laxa, multiflora; ramuli patentes, capillares, ramosi, flexuosi, supernè scabri, fasciculati, fasciculis alternatim per rachin communem positis, unde primo intuitu verticillata videtur panicula. Glumæ ovatæ, concavæ; calycinæ tenuiores, trinerviæ, scabriusculæ; corollinæ calyci contrariæ, subcorneæ, nitidæ, alterâ latiore, alteram amplectente. Stamina glumis duplo longiora, sub anthesin exstantia. Stigmata lateraliter exserta, plumosa.

lendigerum. 2. MILIUM paniculâ subspicatâ, floribus aristatis. *Milium lendigerum.* *Sp. Pl.* 91. *Witb.* 122.

Schreb. Gram. vol. 2. 14. t. 23. f. 3.

Agrostis australis. *Linn. Mant.* 1. 30.

A. rubra. *Huds. ed.* 1. 26.

A. ventricosa. *Gouan. Hort.* 39. t. 1. f. 2.

Alopecurus ventricosus. *Huds. ed.* 2. 28.

Panicum terotinum arvense spica pyramidata.

Raii Syn. 394.

Gramen Alopecuro accedens, ex culmi geniculis spicas cum petiolis longiusculis promens.

Pluk. Phyt. t. 33. f. 6.

Angl. Panick Millet-grass.

In arvis, præcipuè ubi aquæ stagnarunt, sed rariùs.

In the Isle of Shepey plentiful. *Huds.* Corn-fields at Gillingham, Norfolk. *Mr. Woodward.* Corn-fields at Weymouth plentifully. *Lightf. Herb.*

Annua. Fl. Augusto.

Radix exigua, fibrosa. Culmus subsolitarius, basi subinde ramosus, pedalis, erectus, foliosus, lævissimus, plerumque trinodis. Folia utrinque scabra, vaginâ subventricosâ, stipulâ striatâ lacerâ. Panicula erecta, uncialis, bi-vel tri-uncialis, densa, spiciformis, ramulis erectis, adpressis, ramosis. Flores albidî, nitidî, adscendentes. Calyx basi tumidus, lævissimus; supernè compressus.

compressus, exasperatus; apice membranaceus, acutus, valvulis parum inæqualibus. Corolla bivalvis, calyce triplo brevior; valvula exterior ovata, concava, retusa, infra apicem aristata, aristâ geniculatâ; interior minutissima, tenuissima, mutica. Semen compressum, nitidum, fuscum, corollâ vestitum.

28. AGROSTIS. *G. Pl.* 33. *Juss.* 29.

Cal. bivalvis, uniflorus: valvulis acutis. *Cor.* bivalvis, inæqualis, calyce major. *Stigmata* plumosa.

* *Aristatæ.*

1. AGROSTIS petalo exteriore aristâ rectâ strictâ *Spica*
longissimâ infra apicem, paniculâ patulâ. *venti.*

Agrostis Spica venti. Sp. Pl. 91. *Huds.* 30.

With. 126. *Relb.* 26. *Leers.* 19. *t.* 4. *f.* 1.

Gramen miliaceum majus, glumis aristatis, spadicis et pallidis. *Raii Syn.* 405.

G. harundinaccum. Ger. em. 5.

G. agrorum venti spica. Park. Theatr. 1158.

Angl. Silky Bent-grass.

Inter segetes, solo arenoso et sub humido minùs frequens.

In Gayton field and Witton field, Norfolk. *Mr. Crove.* Near Kingston upon Thames, plentifully. *Rev. Dr. Goodenough.* St. Faith's near Norwich; and at Warrington, Lancashire.

Annua. Fl. Junio, Julio.

Radix e fibris numerosis, tortuosis, basi incrassatis, statim attenuatè capillaceis. Culmi plures, tripedales, basi præcipu geniculati, foliosi, læves. Folia latiuscula, supra-pubescentula, subtus scabra. Stipula laciniata. Panicula amplissima, subsecunda; ramulis semiverticillatis, patentibus, inæqualibus, ramosissimis multifloris, angulatis, scabris. Glumæ albidæ vel spadiceæ, carinâ scabræ; calycinæ acuminatæ, muticæ, parum inæquales;

inæquales; corollina exterior paulò infra apicem aristata, interiorem muticam, longèque tenuiorem, amplectens; arista scabra, flore quintuplo longior.

canina. 2. AGROSTIS calycibus ovatis coloratis, corollâ nudâ: aristâ dorsali incurvâ, culmis prostratis subramosis.

Agrostis canina. *Sp. Pl.* 92. *Huds.* 30 (α).
Relb. 26. *Sibth.* 36.

A. vinealis. *With.* 127.

Angl. Brown Bent-grass.

In pratis et pascuis humidiusculis.

Perennis. Fl. Julio.

Culmi decumbentes, pedales vel tripedales, foliosi. Folia utrinque scabra. Panicula erecta, semiverticillata. Flores ovati, nitidi, basi purpurascens. Glumæ calycinæ ovatæ, minùs patulæ, parum inæquales, exterior dorso scabra, interior lævis. Corolla calyce paulò brevior, albida. Arista e basi ferè valvulæ majoris, câque duplo longior, fuscescens, medio geniculata, inflexa.

littoralis. 3. AGROSTIS calycibus lineari-lanceolatis aristatis, corollâ nudâ: aristâ sub-terminali rectâ, culmis decumbentibus.

Agrostis littoralis. *With.* 129. *t.* 23. corolla ex errore mutica delineatur.

Angl. Sea-side Bent-grass.

In littoribus marinis lutosi.

In salt marshes near Cley, Norfolk. *Rev. Mr. Bryant.* *Herb. D. Rose.*

Perennis. Fl. Augusto.

Radix repens ramosa. Culmi basi decumbentes, geniculati, subpedales, simplices, foliosi, glabri. Folia latiuscula, plana, nervosa, utrinque margineque scabra. Stipula striata, subpubescens, minùs lacera. Panicula erecta, bi-vel tri-uncialis, densa, lobata, semiverticillata, pedunculis capillaribus, scabris. Flores albidii, apice virentes

virescentes vel purpurascens. Glumæ calycinæ linearilanceolatae, patentes, subæquales, carinâ scabrâ, aristis terminalibus, rectis, longitudine valvularum. Corolla calyce paulo brevior, alba. Arista ex apice ferè valvulæ majoris, recta, brevis, subinde calycem vix excedente.

4. AGROSTIS calycibus lanceolatis corollâ basi *setacea*. aristatâ; aristâ geniculatâ, foliis radicalibus setaceis.

Agrostis setacea. Curt. Lond. fasc. 6. t. 12.

A. alpina. With. 128.

A. mutabilis. Sibth. 36?

A. canina β & γ . Hudf. 30, 31.

Angl. Briffly Bent-grass.

In ericetis aridis.

In the west of England frequent. *Earl of Gainsborough*, and *the Rev. Dr. Goodenough*. Seacoast near Weymouth. *Duchess Dowager of Portland.* Herb. Lightf.

Perennis. Fl. Julio, Augusto.

Radix fibrosa, tomentosa. Culmus spithamæus, foliosus, declinatus. Folia pallidè viridia, glaucescentia, scabriuscula; radicalia setacea, cæspitosa, erecta, vaginis albidis; caulina solo fertiliore dilatata, striata. Stipula alba, tenuis, erecta, apice erosa. Panicula albida, erecta, semiverticillata; pedunculis cretiusculis, ramosis, scabris. Flores lanceolati, attenuati. Glumæ calycinæ angustè lanceolatae, acutæ, clausæ, parum inæquales, albidæ, carinâ virescente, scabrâ, apice subinde purpurascens. Corolla calyce duplo brevior, tenuissima; valvula exterior basi aristata, aristâ calycem excedente, albâ, ultra medium geniculatâ.

* * *Muticæ.*

5. AGROSTIS paniculâ patente: ramulis divari-*vulgaris*.
catis capillaribus, calyceibus æqualibus, petalo
interiore duplo brevior retuso.

Agrostis

Agrostis vulgaris. With. 132.

A. tenuis. Sibth. 36.

A. capillaris. Relb. 27. Leers. 20. 1. 4. f. 3.

A. polymorpha α. Hudf. 31.

Gramen miliaceum locustis minimis, paniculâ fere arundinaccâ. *Raii Syn. 402.*

β. *A. canina. With. 127.*

Nulla modo ab α differt nisi quod aristata est.

γ. *A. pumila. Linn. Mant. 1. 31. Relb. 28.*

Lightf. 1081. Ic. in titulo vol. 2. malè.

Variat corollâ aristatâ. *D. H. Bryant; nec rarò D. Goodenough.*

δ. Corollâ post anthesin elongatâ. *With. 133. var. 3. Angl. Fine Bent-grass.*

In pratis, pascuis, et versuris agrorum vulgaris.

Perennis. Fl. Julio.

Radix e fibris numerosis, capillaribus, subinde repens. Culmi erecti, glabri, striati, foliosi, altitudine varii. Folia angusta, acuta, scabriuscula, longissimè vaginata; stipula brevissima, crosa, mutica, decurrens. Panicula erecta, purpurascens, alternatim semiverticillata, ramulis patentibus, capillaribus, flexuosis, trichotomis, dichotomis et subdivisis, læviusculis. Flores ovato-lanceolati. Calycis valvulæ subæquales, lanceolatae, acutæ, muticæ, patentes; basi purpuræ; margine scariosæ. Petala tenuissima, albida; exterius calyci fere æquale; interius duplo brevius, retusum. Antheræ fusco-flavæ, longitudine corollæ.

Ab *A. capillari* verâ Linnæi omnino diversa. *Conf. Plant. Ic. ex Herb. Linn. tab. 54.*

A. pumila levissima quidem varietas est *Agrostidis vulgaris* nostræ, culmis e solo steriliori humilioribus.

Var. δ gaudet corollâ calyceque post anthesin elongatâ, ut in *A. albâ* β, quæ vera *A. sylvatica* est.

Stolonifera.

6. **AGROSTIS** paniculâ coarctatâ, culmo ramoso repente, floribus congestis, calycibus æqualibus lanceolatis pubescentibus.

Agrostis

Agrostis stolonifera. *Sp. Pl.* 93. *With.* 131.

Relb. 27. *Sibth.* 26. *Mart. Fl. Rus.* t. 120.

A. polymorpha δ . *Huds.* 31.

Gramen montanum miliaceum minus, radice repente. *Raii Syn.* 402?

G. caninum supinum. *Ger. em.* 26.

Angl. Creeping Bent-grass.

In arvis et pratis humidis.

Perennis. Fl. Julio, Augusto.

Radix perennis, fibrosa, ramosissima, pubescens. Culmi decumbentes, ramosi, foliosi, glabri, geniculis contractis, atro-purpureis, infimis radicanibus. Folia latiuscula, nervosa, utrinque scabra. Stipula multifida. Panicula erecta, ramulis alternatim semiverticillatis, subsecundis, coarctatis; inferioribus patentibus; ab insertione ad apicem densè floriferis, ramosissimis, flexuosis. Flores lineari-lanceolati. Calycis valvulae aequales, obtusiusculae, carinatae, purpurascens, extus pubescentes. Petala calyce breviora, inaequalia, obtusa, mutica, quandoque basi aristata.

7. *AGROSTIS*. paniculâ laxâ, culmo repente, *alba*. calycibus aequalibus lanceolatis nitidis carinâ scabris.

Agrostis alba. *Sp. Pl.* 93. *With.* 129. *Relb.* 28.

Sibth. 37? *Leers.* 21. t. 4. f. 5. *Cullum.* 23.

A. polymorpha ζ (palustris) *Huds.* 32.

Gramen miliaceum majus, paniculâ spadiceâ, et paniculâ viridi. *Dill. in Raii Syn.* 404.

G. palustre, paniculâ speciosâ. *Morif. sect.* 8. t. 6. f. 27?

β . *Agrostis sylvatica.* *Huds. ed.* 1. 28. *Sp. Pl.* 1665. *With. ed.* 2. 74?

A. polymorpha η . *Huds.* 32.

Gramen miliaceum sylvestre, glumis oblongis. *Dill. in Raii Syn.* 404.

Angl. Marsh (β Wood) Bent-grass.

In fossis et palustribus; β in nemorosis.

Perennis. Fl. Julio.

Radix perennis, e fibris longis fibrillosis. Culmus basi decumbens, geniculis inferioribus radicans, foliosus, supernè nudus, erectus, glaber. Folia scabra, vaginis lævibus. Stipula obtusa, bifida, sæpè lacera. Panícula ut in præcedente, etiam a basi ad apicem florifera, sed longè minùs densa. Flores lanceolati, nitidi. Calycis valvulæ æquales, acutæ, carinâ tantum scabræ. Petala calyce breviora, inæqualia, acuta, rariùs infra apicem aristata.

Varietas β maximè notabilis est corollâ calyceque post anthesin elongatâ, extra calycem protrusâ, et incrassatâ. Dissert etiam paniculæ ramis infimis basi nudis, nec undique floriferis. Hujus synonymon Raianum ad Agr. arundinaceam in Sp. Pl. perperàm relatum est.

minima. 8. AGROSTIS paniculâ filiformi, floribus ellipticis retusis muticis, corollâ villosâ.

Agrostis minima. *Sp. Pl.* 93. *Huds.* 32. *With.* 134.

Stillingfl. Misc. præf. 28.

Gramen sparteum, capillaceo folio, minimum.

Dill. Giff. Append. 172. t. 16. f. 1.

G. minimum unciale aut biunciale. *Morif. sect.* 8.

t. 2. f. 10.

G. minimum. *Baub. Hist.* v. 2. 465.

G. min. Anglo-Britannicum. *Lobel. Illust.* 20.

Raii Syn. Indic. Pl. Dub.

Angl. Smallest Bent-grass.

In arenosis maritimis rariùs.

In Wales. *Stillingfleet.* Frequent in sandy pastures on the south-west coast of Anglesea. *Rev. Mr. H. Davies.*

Annua. Fl. Martio, Aprili.

Radix e fibris tenuissimis, albidis, annua. Culmi erecti-
nsculi, unciales vel triunciales, simplicissimi, enodes,
capillares, lævissimi, folio solitario prope basin vaginati,
cæterum

cæterùm nudi, apice purpurascens. Folia omnia ferè radicalia, linearia, obtusa, culmo duplo breviora, basi dilatata, membranacea, vaginantia. Stipula obtusa, plicata, utrinque in marginem membranaceum decurrens. Racemus (potiùs quàm panicula) terminalis, erecta, simplex, rachi flexuosâ, floribus secundis, erectis, adpressis, lævibus, ellipticis, obtusis, muticis, compressis, viridi-purpureis. Calycis valvulæ æquales, earinatæ, retusæ, nitidæ, clausæ. Corolla calyce brevior, villosa, nervosa, apice crosa. Antheræ parùm exsertæ, apice bicornes. Styli filiformes, pubescentes. Flores quandoque monoiei mihi visi sunt.

Ad Poam refert clar. Stokes, fortè novi generis gramen.

29. AIRA. *G. Pl.* 34. *Gærtn. t. 1. Juss.* 31.

Cal. bivalvis, biflorus. *Cor.* bivalvis. *Flosculi* absque interjecto rudimento.

* *Muticæ.*

1. AIRA paniculâ spicatâ, calycibus pedunculo *cristata*, longioribus, petalis acuminatis inæqualibus.

Aira cristata. *Sp. Pl.* 94. *Huds.* 33.

Poa cristata. *With.* 145. *Relb.* 37. *Sibth.* 42.

Gramen pumilum hirsutum, spicâ purpureo-argenteâ molli. *Raii Syn.* 396.

Gr. spicâ cristatâ subhirsutum. *Morif. sect.* 8. *t. 4. f. 7.*

Angl. Crested Hair-grass.

In pascuis elatis sterilibus, murisve, minùs frequens.

Perennis. Fl. Julio, Augusto.

Radix cæspitosa, e fibris simplicibus, tomentosis. Culmi solitarii, simplices, erecti, spithamæi, læves, ad basin præcipuè foliosi. Folia linearia, glabra, rigidiuscula, longiùs vaginata, stipulâ brevi. Panicula erecta, densa, spiciformis, viridi-argentea, pedunculis pubescentibus, ramosis, pedicellis brevissimis, erectis. Flores ovato-lanceolati. Calyx biflorus, rariùs triflorus, nec

quod seiam quadriflorus, obsoletè pubescens, e glumis duabus inæqualibus, æutis, carinatis, utrinque nervosis, margine scariosis, dorso viridibus. Petala exteriora calyci confimilia; interiora longè tenuiora et angustiora, parùm breviora.

Ex habitu *Aira* potiùs quam *Poa* censenda, nec multùm obstat flosculorum numerus, qui plerumque bini sunt. Glumarum muero quandoque in brevem aristam protruditur.

Vereor ne gramen quoddam alienum in Systemate Naturæ cum hocce confuderit Linnæus, et inde *Pois* adfociaverit.

aquatica. 2. *AIRA* paniculâ patente, floribus muticis lævibus obtusis calyce longioribus, foliis planis.

Aira aquatica. *Sp. Pl.* 95. *Huds.* 33. *With.* 135.

Relb. 29. *Sibth.* 38. *Curt. Lond. fasc.* 1. t. 5.

Poa dulcis. *Salisb. Prod.* 20.

Gramen miliaccum aquaticum. *Raii Syn.* 402.

Angl. Water Hair-grass.

In aquosis et ad ripas fluviorum.

Perennis. Fl. Maiò, Junio.

Radix repens, fibris albis, longissimis. Culmus natans, longissimè extensus, ramosus, geniculis inferioribus radicantibus, foliosus, apice erectus. Folia plana, lævia, natantia. Stipula alba, tenuis, brevis, integra. Panicula erecta, semiverticillata, ramulis inæqualibus, patentibus, multifloris, glabris. Glumæ calycinæ inæquales, angulato-truncatæ, purpurascens; exterior basi trinervis: corollinæ calyce ferè triplo longiores, æquales, nervosæ, plicatæ, apice membranacæ, retusæ.

* * *Aristatæ*.

cæspitosa. 3. *AIRA* foliis planis, paniculâ patente, petalis aristatis basi villosis: aristâ rectâ brevi.

Aira cæspitosa. *Sp. Pl.* 96. *Huds.* 34. *With.* 136.

Relb. 29. *Sibth.* 38. *Leers.* 23. t. 4. f. 8.

Gramen miliaceum segetale majus. *Raii Syn.*

403.

G. segetale. *Ger. em.* 5.

Angl. Turfy Hair-grass.

In pratis et nemorosis humidis.

Perennis. Fl. Junio, Julio.

Radix densè caespitosa, fibrosa. Culmi tripedales, erecti, laeves, foliosi, binodes. Folia plana, angusta, mucronata, rigidiuscula, nervosa; subtùs laevia; suprà nervis margineque scabra; radicalia, basi dilatata, vaginantia. Stipula elongata, acuta, bifida. Panícula saepius pedalis, patens, nitida, fuscescens, in umbrosis alba; pedunculi semiverticillati, ramosissimi, capillares, scabri. Flores parvi. Glumæ calycinae subæquales, dorso scabrae. Flosculus alter pedicellatus, pedicello villosus, ultra basin flosculi protensus. Glumæ corollinae glabrae; exterior latior, apice erosa, dorso aristata, aristâ inflexâ, brevi, rariùs glumam excedente.

“ Variat floribus viviparis in locis alpinis.” *D. Davies.*

4. AIRA foliis setaceis, culmo subnudo, paniculâ *flexuosa*, patente trichotomâ, pedunculis flexuosis, aristis geniculatis.

Aira flexuosa. *Sp. Pl.* 96. *Huds.* 34. *With.* 136.

Relb. 30. *Sibth.* 38. *Leers.* 23. t. 5. f. 1.

Fl. Dan. t. 157.

Gramen paniculatum locustis parvis, purpureo-argenteis, majus et perenne. *Raii Syn.* 407.

G. avenaceum paniculatum Alpinum, foliis capillaceis brevibus, locustis purpureo-argenteis splendentibus, et aristatis. *Scheuchz. Prod.* 23.

t. 4. f. 4.

β. Paniculâ minùs patente, pedunculis vix flexuosis.

Aira montana. *Huds.* 35. *With.* 136. *Relb.* 31.

Leers. 24. t. 5. f. 2. nec *A. montana* Linnæi.

G 3

γ. Culmo

- γ. Culmo magis folioso, paniculâ albâ, vix flexuosâ.
 Gramen nemorosum, paniculis albis, capillacco
 folio. *Dill. in Raii Syn.* 407. *Scheuchz. Prod.* 24.
t. 6. *f.* 1. *Morif. sect.* 8. *t.* 7. *f.* ult.
Angl. Waved Mountain Hair-grass.

In cricetis et montosis arenosis. γ in umbrosis.

Perennis. Fl. Julio.

Radix fibrosa, fibris in aridis tomentosis. Culmus pedalis vel sesquipedalis, erectus, nudiusculus, lævis, purpurascens, basin versus uninodis; in γ subtrinodis, atque foliosus. Folia setacea, margine scabriuscula, basi dilatata vaginantia; in α & β brevia, rigidula, omniaque ferè radicalia; in γ longitudine dimidii culmi, sublaevica, caulina longius vaginata. Stipula omnibus brevis, obtusa, bifida. Panicula pauciflora, triehotoma, patens; pedunculis scabris, valdè flexuosis, (in β & γ rectiusculis, minùsque patentibus); pedicellis alternis rectis. Flores erecti, scariosi, argenteo-nitidi, spadicei; in γ virescente-albidi. Glumæ omnes ferè longitudine æquales, apice erosæ; corollinæ interiores angustiores; flosculi basi pilosi, (altero brevius pedicellato), dorso basin versus aristati, aristâ geniculatâ, tortili, glumis sesquilongiori.

Aira setacea *Huds. ed.* 1. nullo modo a varietate nostrâ β differt; sed *A. montana* Linnæi longè diversum est gramen, nondum in Britannia observatum.

canescens. 5. *AIRA* foliis setaceis, culmo folioso, paniculâ coarctatâ, aristis apice clavatis calyce brevioribus.

Aira canescens. *Sp. Pl.* 97. *Huds.* 36. *With.* 137. *t.* 24. *Dicks. Dr. Pl.* 4.

Gramen miliacum maritimum molle. *Dill. in Raii Syn.* 405.

G. paniculatum capillacco folio minimum. *Tourn. Inst.* 523.

G. exile durius Norvegicum aut Danicum, Scopario

Scopario Gramini cognatum. *Lobel. Advers. nova* 466. f. 1.

Angl. Grey Hair-grafs.

In arenosis maritimis.

On the coast of Norfolk and Suffolk.

Perennis. Fl. Julio.

Radix e fibris longissimis, capillaceis, cæspitosa. Culmi plures, 3 ad 6 unciales, binodes, adscendentes, læves, foliosi, basi geniculati. Folia fetacea, erecta, rigida, scabra, glauca; caulina longius vaginata. Stipula lanceolata, acuta. Panicula erecta, uncialis, lanceolata, coarctata, alternatim ramosa, subsecunda. Glumæ calycinæ cinereo-argenteæ, acutæ, dorso subpubescentes, flosculis duplo longiores. Flosculi basi pilosi, altero pedicellato. Arista e basi ferè petali exterioris; infernè tortilis, ferruginea; medio geniculata, dentibus cincta; apice clavata, albida, calycem non excedens. Antheræ violacæ, inclusæ.

6. AIRA foliis fetaceis: vaginis angulatis, floribus *præcox.* paniculato-spicatis, flosculis sessilibus basi nudis dorso aristatis.

Aira præcox. *Sp. Pl.* 97. *Huds.* 36. *With.* 137. *Relb.* 31. *Sibth.* 39. *Abbot.* 15. *Curt. Lond. fasc.* 3. t. 7. *Dicks. H. Sicc. fasc.* 4. 4. *Fl. Dan. t.* 383.

Gramen parvum præcox, paniculâ laxâ canescente. *Raii Syn.* 407. t. 22. f. 2. *Pluk. Phyt. t.* 33. f. 9.

Angl. Early Hair-grafs.

In glareosis et sterilibus frequens.

Annua. Fl. Maio, Junio.

Radix parva, fibrosa. Culmi erecti, bi- vel tri-unciales, solo fertili quandoque spithamæi, lævissimi, foliosi, pallidi. Folia fetacea, pallidè virentia; radicalia citò marcescentia; caulina longius vaginata, vaginis inflatis, angulatis,

latis, nervosis, lævibus. Stipula obtusa. Panicula simpliciuscula, pauciflora, erecta, coarctata, vix uncialis. Flores albido-virescentes, nitidi, erecti, glumis calycinis æqualibus, dorso scabris. Flosculus uterque sessilis, glumis calycem subæquantibus, basi nudis, exterioribus basin versus aristatis. Arista calyce duplo ferè longior, setacea, geniculata.

*caryo-
phylla.*

7. AIRA foliis setaceis, paniculâ trichotomâ divaricatâ, flosculis sessilibus, aristâ geniculatâ dorsali.

Aira caryophylla. Sp. Pl. 97. Hudf. 36. With. 138. Relb. 32. Sibth. 39. Curt. Lond. fasc. 6. t. 6. Dickf. H. Sicc. fasc. 4. 5. Fl. Dan. t. 382. Stillingfl. Misc. t. 5.

Gramen paniculatum, locustis parvis purpureo-argenteis, annuum. *Raii Syn. 407. Moris. sect. 8. t. 5. f. 11.*

Angl. Silver Hair-grass.

In glareosis et arenosis frequens.

Annua. Fl. Julio.

Radix exigua, fibrosa. Culmus basi ramosus, foliosus, glaber. Folia radicalia marcescentia; caulina longius vaginata, vaginis striatis, parùm inflatis, margine scabris. Stipula lanceolata, elongata. Paniculæ terminales, sæpiùs quadrifidæ, ramis trichotomis, lævibus, divaricatis. Flores parvi, divergentes, ovati, purpureo-argentei, nitidi. Glumæ calycinæ scariosæ, dorso scabræ. Flosculus uterque sessilis, basi obsolete pilosus. Corolla e glumis durioribus, apice scariosis; exteriori basin versus aristatâ, aristâ calycem excedente, setacâ, geniculatâ.

30. HOLCUS. *G. Pl. 541. Gærtn. t. 80. Juss. 30.*

Cal. bivalvis, biflorus; flosculo altero masculino. Cor. bivalvis; exteriori aristatâ.

1. HOLCUS

1. *HOLCUS* glumis villosis, flosculo hermaphro-*lanatus*.
ditico mutico, masculo aristâ arcuato-re-
curvâ.

Holcus lanatus. *Sp. Pl.* 1485. *Hudsf.* 440. *With.*

134. *Relb.* 370. *Sibth.* 40. *Curt. Lond. fasc.* 4.

t. 11. *Dicksf. H. Sicc. fasc.* 9. 16. *Leers.* 219.

t. 7. *f.* 6. *Schreb. Gram.* 145. *t.* 20. *f.* 1.

Gramen miliaceum pratense molle. Raii Syn.

404.

Angl. Meadow Soft-grass.

In pratis et pascuis frequens.

Perennis. Fl. Junio, Julio.

Radix fibrosa, cæspitosa. Culmī plures, erecti, striati, foliosi, retrorsum villosi, apice glabriusculi. Folia plana, pallida, mollia, subtus magis villosa. Stipula truncata, brevis, denticulata. Panicula erecta, effusa, mollis, purpureo-albida, demum pallidior; ramulis semiverticillatis, ramosissimis, capillaribus, villosis. Calycis glumæ longitudine æquales, mucronatæ, villosæ, subinde tuberculatæ, coloratæ; interior latior, trinervis. Flosculi pedicello communi flexuoso inserti, masculus autem supra hermaphroditicum, calyce breviores, basi nudi, glumis firmioribus, nitidis, glabris; exterioribus latioribus. Arista e glumâ exteriori floris masculini, mox arcuato-recurva, calyce inclusa. Antheræ longitudine corollæ.

Pistillum abortivum, à Curtisio detectum, in aliquot floribus masculinis inveni, nec tamen in omnibus.

2. *HOLCUS* glumis subæqualibus nudiusculis, *mollis*.
flosculo hermaphroditico inferiore mutico,
masculo aristâ geniculatâ, radice repente.

Holcus mollis. *Sp. Pl.* 1485. *Hudsf.* 440.

With. 135. *Relb.* 376. *Sibth.* 40. *Curt. Lond.*

fasc. 5. *t.* 8. *Dicksf. H. Sicc. fasc.* 9. 17.

Leers. 218. *t.* 7. *f.* 7. *Schreb. Gram.* 148.

t. 20. *f.* 2.

Gramen

Gramen miliaceum aristatum molle. *Raii Syn.*

404.

Angl. Creeping Soft-grass.

In arvis, nemorosis, et ad sepes.

Perennis. Fl. Julio, Augusto.

Radix repens, longissimè extensa. Culmus solitarius, ascendens, foliosus, glaber, geniculis lanatis. Folia virentia, glabriuscula. Stipula rotundata, dentata. Panicula erecta, effusa, purpureo-albida; ramulis plerumque binis, ramosis, capillaribus, villosis. Calycis glumæ ut in præcedente, sed carinâ tantum ciliatâ. Flosculi ut in præcedente, basi vero hispidi. Arista tortilis, recta, calyceem excedens, demùm geniculato-inflexa, neque areolata. Flos superior pistillo imperfecto quoque gaudet, monente Curtio.

avena-
ceus.

3. **HOLCUS** glumis inæqualibus glabris, flosculo hermaphroditico superiore subinutico, masculo aristâ geniculatâ refractâ, radice nodosâ.

Holcus avenaceus. *Wiggers. Fl. Holst.* 72.
Sibth. 40.

Avena elatior. *Sp. Pl.* 117. *Huds.* 53. *With.* 163.
Relh. 48. *Abbot.* 25. *Curt. Lond. fasc.* 3.
t. 6. *Leers.* 40. *t.* 10. *f.* 4. *Schreb. Gram.* 25.
t. 1.

A. nodosa. *Cullum.* 41.

Gramen nodosum avenaceâ paniculâ. *Raii Syn.*
406.

G. avenaceum elatius, radice tuberculis prædita.
Morif. sect. 8. *t.* 7. *f.* 38.

G. caninum nodosum. *Ger. em.* 23.

β. *G. avenaceum* paniculâ accrosâ, semine paposo. *Dill. in Raii Syn.* 406.

Angl. Oat-like Soft-grass.

In pascuis, apricis, et ad vias.

Perennis.

Perennis. Fl. Junio, Julio.

Radix magis vel minus nodosa, e tuberibus plerumque binis, altero super alterum, radiculis villosis. Culmus erectus, tri- vel quadripedalis, simplex, foliosus, glaber. Folia saturatè viridia, suprà margineque scabra. Stipula brevis, retusa, denticulata. Panicula erecta, ramulis semiverticillatis, ramosis, scabris, sub anthesin præcipuè patentibus. Calyx petalis brevior, e glumis scariosis, acutis, carinâ scabris; altero interiore, longiore, trinervi. Flosculus masculinus sessilis, basi setosus, dorso basin versus aristatus, aristâ tortili, geniculatâ, flosculo duplò longiore; hermaphroditicus parùm elevatus, aristâ subterminali, rectâ, brevi. Glumæ corollinæ exteriores firmiores, nervosæ, pubescentiusculæ; interiores tenuissimæ, angustiores, parùmque breviores, nudæ.

Flores masculini sæpius pistillo obsoleto, abortivo, gaudent,

31. MELICA. *G. Pl.* 34. *Gærtn. t.* 80. *Juss.* 31.

Cal. bivalvis, sub-bisflorus; rudimentum tertii floris inter flosculos. *Cor.* bivalvis.

1. MELICA petalis imberbibus, paniculâ secundâ *uniflora*, ramosâ, floribus erectis, calyce unifloro.

Melica uniflora. *Retz. Obs. Bot. fasc.* 1. 10.

With. 139. *Relb. Suppl.* 1. 8. *Sibth.* 40.

Abbot. 16. *Curt. Lond. fasc.* 5. t. 10. *Dicks.*

H. Sicc. fasc. 8. 6. *Mart. Fl. Rust. t.* 64.

M. nutans. *Huds.* 37. *Rel. Rudb.* 18. f. 2.

Gramen avenaceum nemorense, glumis rarioribus ex fusco xerampelinis. *Raii Syn.* 403.

G. avenaceum spicâ muticâ rariore glumâ.

Moris. sect. 8. t. 7. f. 49.

Angl. Wood Melic-grafs.

In nemoribus et dumetis.

Perennis. Fl. Maio, Junio.

Radix

Radix fibrosa. Culmi plures, erecti, sesquipedales, graciles, foliosi, glabri. Folia plana, tenuia, subtus margineque scabra; vaginâ folio duplo circiter brevior, pubescente; stipulâ brevi, polymorphâ. Panícula erecta pauciflora; ramulis secundis, capillaribus, racemosis, scabris, infimis geminis. Flores erecti. Calyx purpureus, parum inæqualis, glaber, obtusus, muticus, nervosus. Flosculus fertilis sessilis, extus gibbosus, intus concavus, glumis viridibus, nervosis; sterilis pedunculo crasso, inflexo, insertus, e glumis aliquot parvis vacuis constans.

nutans. 2. *MELICA* petalis imberbibus, paniculâ coarctatâ secundâ subsimplici, floribus pendulis, calyce bifloro.

Melica nutans. *Sp. Pl.* 98. *With.* 138. *Leers.* 24.

t. 3. f. 4. Dickf. H. Sicc. fasc. 8. 5. Rel.

Rudb. 18. f. 1. Schreb. Gram. 62. t. 6. f. A.

Mart. Fl. Rust. t. 65. Curt. Lond. fasc. 6. t. 4.

M. montana. *Huds.* 37.

Grainen avenaceum, locustis rubris, montanum.

Raii Syn. 403.

G. avenaceum locustis rubris. *Park. Theatr.* 1151.

Angl. Mountain Melic-grass.

In nemoribus montosis Angliæ borealis.

Near Kendal. *Ray.* Helks-wood by Ingleton, Yorkshire. *Mr. Crowe and Mr. Woodward.*

Perennis. Fl. Junio, Julio.

Radix repens. Culmus erectus, sesquipedalis, foliosus, angulatus, angulis scaber. Folia plana, margine scabra; vagina longa, aspera; stipula brevissima, erosa. Panícula erecta, vix ramosa; pedunculis plerumque simplicibus, secundis, capillaribus, flexuosis, scabris, unifloris. Flores nutantes. Calyx purpureus, margine scariofo, albo, parum inæqualis, glaber, obtusiusculus, muticus, obsolete nervosus. Flosculi virecentes; glumâ exteriori glabrâ nervosâ, gibbosâ; interiori pubescenti, planiusculâ; subquaterni, pedicello communi affixi, alterni,

terni, distichi; primus ad basin, sessilis; secundus parum elevatus, uterque fertilis; tertius minor, sterilis; quartus minimus, e glumâ parvâ, retusâ, inani constans.

3. MELICA petalis acutis imberbibus, paniculâ *cærulea*, coarctatâ, floribus erectis cylindricis.

Melica cærulea. *Linn. Mant.* 2. 325. *With.* 139.

Relb. 32. *Sibth.* 39. *Abbot.* 16. *Curt. Lond. fasc.* 5. *t.* 11.

Aira cærulea. *Sp. Pl.* 95. *Huds.* 33. *Leers.* 22. *t.* 4. *f.* 7. *Fl. Dan.* *t.* 239.

Gramen pratense serotinum, paniculâ longâ purpurascente. *Raii Syn.* 404. *Morif. sect.* 8. *t.* 5. *f.* 22.

Angl. Purple Melic-grafs.

In inundatis sterilioribus frequens.

Perennis. Fl. Augusto.

Radix bulbosa, fibris crassis, tortuosis. Culmus pedalis vel sesquipedalis, erectus, rigidus, teres, glaber, striatus, basin versus uninodis et foliosus, superne nudus. Folia rigidula, acuminata, vix scabra; stipula e pilis aliquot brevissimis. Panicula erecta, violacea, ramosa, coarctata. Pedunculi plerumque bini, simplices aut divisi, erecti, angulati, flexuosi, angulis scabri. Flores erecti, glabri. Calyx e glumis subæqualibus, ovatis, acutis, nervosis, carinatis, purpureis. Flosculi quaterni, pedicello communi alternatim affixi, glumis lanceolatis, acutis, dorso striatis, basi nudis nec pilosis, quorum duo inferiores tantum fertiles sunt. Anthræ atropurpureæ.

Variat paniculâ pallidâ in nemorosis. *Lightf. Herb.*

32. SESLERIA. *Scop. Carn. ed.* 1. 189. *Juss.* 31.

Cal. bivalvis, subtriflorus, aristatus. *Cor.* bivalvis; valvulâ exteriori tridentatâ; interiori bidentatâ.

1. SES-

cærulea. 1. *SESLERIA* spicâ ovato-oblongâ imbricatâ,
bracteis alternis.

Sesleria cærulea. *Scop. Carn. ed. 2. vol. 1. 63.*

Arduin. Spec. 2. 18. t. 6. f. 3, 4, 5. With. 140.

S. n. 1446. Hall. Hist. v. 2. 217.

Cynosurus cæruleus. Sp. Pl. 106. Hudsf. 59.

Jacq. Misc. v. 2. 66. Ic. rar. v. 1. 21. Dickf.

H. Sicc. fasc. 6. 3. Mart. Fl. Rust. t. 20. Curt.

Cat. n. 18.

Gramen glumis variis. Bauh. Prod. 21. f. 1.

Scheuchz. Agrost. 83. t. 2. f. 9. A. B.

*G. parvum montanum spicâ crassiore purpureo-
cæruleâ brevi. Raii Syn. 399.*

Angl. Blue Moor-grass.

In humidis alpinis calcareis.

Crevices of the limestone rocks at the foot of
Ingleborough. *Dr. Stokes. About Settle. Curtis.*

Conzick Scar, near Kendal; *Mr. Gough. With.*

At the top of Barrow-field wood in the same
neighbourhood. *Mr. Woodward and Mr. Crowe.*

Perennis. Fl. Aprili.

*Radix fibrosa, cæspitosa, longissima. Culmus vix pedalis,
erectus, simplex, enodis, teres, striatus, glaber, basi sub-
foliosus. Folia linearia, obtusa, recurva, striata, mar-
gine, quandoque etiam dorso, scabra. Vaginæ tubu-
losæ, breves, striatæ, glabræ, pallidæ. Stipula obso-
leta. Spica erecta, brevis, obtusa, cæruleo vel fusco
varia, nitida, subsimplex. Bractææ solitariae, ad basin
spicularum duarum vel trium inferiorum, amplexi-
caules, membranaceæ, apice dilatatæ, crosæ. Spiculæ
fæpius binatæ, pedicellatæ, erectæ, ovato-lanceolatæ,
glabræ, plerumque bi- vel tri-floræ. Glumæ calycinae
subæquales, ovatæ, brevius aristatæ, membranaceæ,
carinatae. Petala nervosa, membranacea; exterius tri-
dentatum, dente intermediâ in aristam brevem pro-
ductâ; interius bidentatum. Antheræ lineares, sub
anthesin exsertæ. Stigmata subulata, pubescentia.*

Cum

Cum Cynofuro nullam habet affinitatem, neque pertinent ad fructificationem bractæe istæ pedicellis tantum inferioribus subjectæ. Cum Phleo in multis convenit, monente Scopio, at distinctum genus est.

33. POA. *G. Pl.* 34. *Juss.* 32.

Cal. bivalvis, multiflorus. *Spicula* basi rotundata.

Cor. bivalvis, valvulis ovatis, acutiusculis, muticis.

1. POA paniculâ erectâ ramosâ laxâ, spiculis sex-*aquatica.*
floris linearibus, flosculis obtusis septem-
nerviis.

Poa aquatica. Sp. Pl. 98. *Huds.* 38. *With.* 140.

Relb. 33. *Sibth.* 41. *Abbot.* 17. *Curt. Lond.*
fasc. 5. *t.* 12. *Leers.* 26. *t.* 5. *f.* 5.

Gramen aquaticum majus. Raii Syn. 411.

Ger. em. 6.

G. paniculatum aquaticum latifolium. Moris.

sect. 8. *t.* 6. *f.* 25.

Angl. Reed Meadow-grass.

In fossis et ad ripas fluviorum.

Perennis. Fl. Julio.

Radix fibrosa, albida. Culmi plures, subsexpedales, erecti, ancipites, striati, glabri, foliosi, geniculis inferioribus verticillatim radicanibus. Folia ensiformia, acuta, recta, margine carinâque scabra, vaginâ læviusculâ. Stipula brevis, obtusissima, integra. Panicula amplissima, ramosissima, erecta, laxa; ramis semiverticillatis, scabris; ramulis alternis, flexuosis. Calyx subæqualis, albidus, nitidus. Flosculi 5 ad 8, vix plures, subcylindrici, remotiusculi, obtusi, extus purpurascens, scabri. Glumæ corollinæ exteriores septemnerviæ; interiores emarginatæ. "Variat floribus viviparis." *D. Davies.*

Gramen pecoribus gratum. Withering.

2. POA paniculâ ramosâ divaricatâ, spiculis ad-*fluitans.*
pressis

pressis teretibus multifloris, flosculis obtusis septemnerviis basi duplicato-nervosis.

Poa fluitans. *Salisb. Prod.* 21.

P. n. 1453. *Hall. Hist. v.* 2. 219.

Festuca fluitans. *Sp. Pl.* 111. *Huds.* 46. *With.* 156. *Relb.* 42. *Sibth.* 45. *Abbot.* 21. *Curt. Lond. fasc.* 1. *t.* 7. *Mart. Fl. Rust. t.* 113. *Schreb. Gram.* 37. *t.* 3. *Fl. Dan. l.* 237. *Leers.* 35. *t.* 8. *f.* 5.

Gramen aquaticum cum longissimâ paniculâ.
Raii Syn. 412.

G. loliaceum fluviatile, longissimâ paniculâ.
Morif. sect. 8. *t.* 3. *f.* 16.

Angl. Flote Meadow-grass.

In fossis et rivulis lentè fluentibus frequens.

Perennis. Fl. totâ æstate.

Radix repens, longè extensa. Culmus basi radicans atque ramosus, obliquè adscendens, teres, striatus, foliaceus, glaber. Folia plana, obtusiuscula, latiuscula, lævia, striata, erecta; inferiora flaccida, natantia; omnia longius vaginata, vaginis compressis, striatis, lævibus. Stipulæ subdeltoideæ, breves, crosæ, tenuissimæ. Panicula erectiuscula, glabra, striata, ramis alternis, patentibus, spiculis alternis, subseffilibus, ramulo adpressis, lincariteretibus, 8 ad 12-floris. Calyx inæqualis, membranaceus, obtusus, estriatus, carinâ viridi. Flosculi remoti, subcylindrici, obtusi, oculo armato scabriusculi, septemnervii, nervis minoribus insuper basin versus interstinctis, apice scariosi, albi, sæpiùs denticulati; glumis interioribus emarginatis.

Variat paniculâ minùs ramosâ. Glumæ muticæ; structura florum affinis præcedenti et tribus vel quatuor sequentibus speciebus; omnia verò Poam nec Festucam planè declarant.

distans. 3. POA paniculâ ramosâ effusâ; ramis demùm reflexis, spiculis quinquefloris, flosculis obtusissimis obsoletè quinquenerviis nitidis.

Poa

Poa distans. *Linn. Mant.* 32. *With.* 141. *t.* 25.

P. retroflexa. *Curt. Lond. fasc.* 6. *t.* 1.

P. salina. *Pollich. v.* 1. 89.

Aira aquatica β . *Hudsf.* 34. *Linn. Syst. Nat. ed.* 12. *v.* 2. 91.

Angl. Reflexed Meadow-grass.

In pratis et ruderatis maritimis.

On a sandy soil in Devonshire, Kent, Yorkshire, and Lancashire. *Hudson.* Frequent near the sea-coast; also between London and Hampstead. *Curtis.*

Perennis. Fl. Julio.

Radix fibrosa. Culmi plures, basi decumbentes, subpedales, teretes, foliosi, striati, glabri. Folia longius vaginata, acutiuscula, lævia, glauca, plana; radicalia parum involuta. Stipula obtusa, emarginata. Panicula erecta, ramis semiverticillatis, angulatis, scabris, subflexuosis, spicatis vel ramosis, longitudine variis, demum maximè retrofractis. Spiculæ lineares, glabræ, 4 ad 7-floræ, albido vel purpureo variegatæ. Calyx inæqualis, carinatus, parvus, obtusus, margine membranaceus; glumâ exteriori obsoletè trinervi. Flosculi remoti, ferè ut in *Poa aquatica*, subcylindrici, obtusissimi, retusi, quinquenerves, apice scariosi; glumis interioribus emarginatis, (in margine scabris *Curtis.*)

4. POA paniculâ ramosâ subcoarctatâ, spiculis *maritima.* quinquefloris, flosculis obtusiusculis teretiusculis obsoletè quinquenervibus, radice repente.

Poa maritima. *Hudsf.* 42. *With.* 147. *Lightf.* 98.

Relb. 38. *Dicks. H. Sicc. fasc.* 11. 2. *Flo.*

Dan. t. 251. *Pourret. Act. Tolos. v.* 3. 325.

Græmen paniculatum maritimum vulgatissimum.

Raii Syn. 409.

β . G. caninum maritimum paniculatum. *Raii Syn.* 410. *Hudsf.* 42. non vidi.

Angl. Creeping Sea Meadow-grass.

In palustribus maritimis.

Perennis. Fl. Julio.

Radix repens. Culmi basi decumbentes, pedales, striati, glaberrimi, teretes, foliosi. Folia longius vaginata, involuta, glauca, acuta. Stipula obtusa, subinde acutiuscula. Panicula erecta, stricta, ramis plerumque binatis, angulatis, scabris, parum flexuosis, racemosis, sub florescentiâ patentibus, dein erectis, aliquando divaricatis ut in præcedente. Spiculæ lineares, glabræ, subquinquefloræ, purpurascentes. Glumæ calycinæ inæquales, ovatæ, acutiusculæ, carinatæ; exterior trinervis. Flosculi remotiusculi, oblongi, subcylindrici, obtusiusculi nec retusi, vix manifestè quincunervos, apice minus scariosi; glumæ interiores margine scabræ, apice emarginatæ.

Præcedenti valdè affinis, at, ni fallor, distincta, radice repente, flosculis longioribus, acutioribus, minimè retusis.

procumbens.

5. POA paniculâ lanceolatâ secundâ coarctatâ scabrâ; rachi tereti, spiculis subquinquefloris, flosculis obtusiusculis nervosis.

Poa procumbens. *Curt. Lond. fasc. 6. t. 11. Engl. Bot. t. 532.*

P. rupestris. *With. 146. t. 26.*

Angl. Procumbent Sea Meadow-grass.

In ruderatis maritimis inundatis.

At the foot of St. Vincent's rocks, found by Mr. Curtis; on waste ground near the floating dock, Bristol; also on Scarborough new pier, within the spray of the sea. *Sir T. Frankland.*

Annua. Fl. Julio, Augusto.

Radix fibrosa, cæspitosa. Herba glauca. Culmi plures, prostrati, spithamæi, geniculati, foliosi, glabri. Folia longissimè vaginata, latiuscula, acuta, plana, nervosa, scabriuscula. Stipula acutiuscula, crosa. Panicula rigida, secunda, semicylindrica, rachi ramulisque teretiusculis, scabris. Spiculæ distichæ, sessiles, lanceolatae, 4 vel 5-floræ, glaucæ, glabræ. Glumæ calycinæ obtusæ,

tusæ, valdè inæquales, carinatae; exterior quinque-nervis. Flosculi obtusifusculi, quinquenerves, apice membranacei erosi; glumæ interiores ciliatæ. Cynofurus durus Linnæi, gramen ad Poas proculdubio amandandum, huic proximum est; vix enim discrepat nisi habitu rigidiori, paniculâ simpliciore, rachi flexuosâ complanatâ glabrâ estriatâ, glumis calycinis exterioribus novemnervibus.

6. POA paniculâ lanceolatâ disticho-secundâ *co-rigida*. arctatâ glabrâ; rachi marginatâ, spiculis septemfloris, flosculis teretibus enervibus.

Poa rigida. Sp. Pl. 101. Hudf. 42. With. 146.

Relb. 35. Sibth. 42. Curt. Lond. fasc. 2. t. 4.

Gramen exile duriusculum in muris et aridis proveniens. *Raii Syn. 410.*

G. loliaceum murorum duriusculum, spicâ erectâ rigidâ. Moris. sect. 8. t. 2. f. 9.

Engl. Hard Meadow-grass.

In muris et arenosis frequens.

Annua. Fl. Junio.

Radix fibrosa. Culmi plures, erectiusculi, in maritimis sæpè prostrati, spithamæi, rigidi, teretes, purpurascens, glaberrimi, nitidi, basi foliosi. Folia acuta, sæpiùs involuto-setacea, nervosa, subtùs lævissima, suprà scabra. Stipula obtusa, lacera. Panicula rigida, lanceolata, duplici ordine secunda, rachi flexuosâ, alatâ, angulis scabris. Spiculæ lineares, pedicellatæ, 4 ad 8-floræ, glabræ. Glumæ calycinæ subæquales, carinatae, acutæ. Flosculi obtusifusculi, cylindrici, vix carinati, omninò enerves, apicem versus scabri; glumæ interiores ciliatæ.

7. POA paniculâ secundâ coarctatâ, culmo adscen- *compress-* dente compresso, flosculis angulosis basi villo *sa.* complicato connexis.

Poa compressa. Sp. Pl. 101. Hudf. 41. With. 147.

Relb. 36. Sibth. 42. Engl. Bot. t. 365. Dickf.

H. Sic. fasc. 6. 1. Leers. 30. t. 5. f. 4.

Gramen pratense paniculatum medium. *Raii*

Syn. 409.

Angl. Flat-stalked Meadow-grass.

In muris, pratense ficioribus.

Perennis. Fl. Julio, Augusto.

Radix repens. Culmus basi decumbens, foliosus, geniculatus, supernè erectus, nudus, striatus, valdè compressus, quâ notâ facilè dignoscitur. Vaginæ longitudine foliorum, ancipites, striatæ, parùm ventricosæ. Folia plana, acuta, glaucescentia, lævia. Stipula brevis, obtusiuscula. Panicula erecta, secunda, obtusiuscula, coarctata, rachi compresso-tereti, ramulis angulatis, scabris, subflexuosis, sub florescentiâ patulis, dein erecto-adpressis. Spiculæ subsessiles, ovatæ, vel ovato-lanceolatæ. Glumæ calycinæ subæquales, ovatæ, nervosæ. Flosculi a tribus ad novem, arctè imbricati, ovati, angulosi, trinerves, obtusiusculi, sub apicem purpurei, apice scariosi albi, basi villis tenuissimis, complicatis, brevioribus, connexi.

alpina. 8. POA paniculâ diffusâ, spiculis quadrifloris cordatis, glumis ovatis subfalcatis liberis, stipulis inferioribus brevissimis.

Poa alpina. *Sp. Pl.* 99. *Lightf.* 96.

P. pratensis *β.* *Huds.* 39. ex *Herb. D. Pitchford.*

Gramen paniculatum alpinum majus, paniculâ speciosâ variegatâ. *Scheuchz. Prod.* 20. t. 3.

β. vivipara. *Sp. Pl.* 99. *Huds.* 39.

Gramen alpinum latifolium, paniculâ laxâ, foliaceâ, foliolis in paniculâ paucioribus et magè crispis. *Scheuchz. Agrost.* 212. t. 4. f. 14.

Angl. Alpine Meadow-grass.

In alpibus Scoticis.

Inch Earn, north of Forfar. *Mr. Muckay.*

Perennis. Fl. Julio.

Radix fibrosa, fibris simplicibus, elongatis, glabris. Folia radicalia plurima, patentia, plana, latiuscula, nervosa, subtus

subtùs præcipuè glaucescentia, apicc obtusiuscula, mucronulata, margine scabra, vaginis brevibus, erectis, compressis, stipulâ brevi, obtusâ; caulina duo, brevissima, vaginis longissimis, stipulis lanceolatis, acutis. Culmus pedalis, adscendens, binodis, supernè nudus, teres, purpurascens, nitidus. Panicula brevis, subrotunda vel ovata, diffusa, ramis binatis alternatim ramosis, angulatis, læviusculis, subindè scabris. Spiculæ cordatæ vel latè ovatæ, compressæ, purpureo et spadiceo nitentes, tri- vel quadrifloræ, in hortis sexfloræ. Glumæ calycinæ subæquales, ovatæ, acutæ, subfalcatæ, obsoletè nervosæ, carinâ scabræ. Flosculi ovati, acutiusculi, margine scariosi et sæpiùs sericei, basi setosi, sed liberi, neque pilis complicatis connexi; valvulæ interiores emarginatæ, ciliatæ.

Sub cœlo pluvioso alpium, mutatis in folia glumis, fructificatione denique in gemmam, vivipara sæpè evadit.

9. POA paniculâ flexuosâ, spiculis trifloris, glumis *flexuosa*.
ovatis basi villo connexis, stipulis omnibus lanceolatis.

Angl. Zigzag Meadow-grass.

In alpibus Scoticis.

On Ben Nevis. *Mr. Muckay*.

Perennis. Fl. Julio.

Radix fibrosa. Culmi plures, spithamæi, adscendentes, geniculati, subtrinodes, foliacei, supernè subangulati, striati, glaucescentes. Folia erecto-patentia, plana, angusta, acuta, suprâ scabra, vaginis omnibus elongatis, laxis, parùm compressis, striatis, glabris, stipulis omnibus lanceolatis, acutis, foliorum caulinorum sæpiùs laceris. Panicula ovata, abbreviata, subcondensata, ramis binatis, alternatim ramosis, magis minùsve basi flexuosis, supernè angulatis, scabris. Spiculæ latè ovatæ, compressæ, glaucæ, albido et purpureo variegatæ, plerumque trifloræ. Glumæ calycinæ parùm inæquales, lanccolato-ovatæ, acutæ, vix falcatæ, carinâ scabræ. Flosculi lanccolati, acutiusculi, margine denudati, carinâ scabri, basi villis complicatis tenuissimis connexi; valvulæ interiores vix ciliatæ.

Distincta a præcedente colore glauco, culmo magis folioso, foliis angustioribus, acutioribus, suprâ scabris, vaginis longioribus, stipulis omnibus conformibus, pedunculis flexuosis, glumis angustioribus, basi villo connexis, neque carinâ neque margine setoso-sericeis.

bulbosa. † 10. POA paniculâ subflexuosâ, spiculis quadrifloris, glumis villo connexis, foliis ferrulatis, culmo basi bulboso.

Poa bulbosa. *Sp. Pl.* 102. α et γ . *Huds.* 41.

Gramen xerampelinum, miliaceâ, pertenui ramosâque sparsâ paniculâ. *Scheuchz. Agr.* 185, ex descr. opt. *Vaill. Par.* 91, ex descr. excluso synonymo Lobelii.

G. vernum, radice Ascalonitidis. *Vaill. Par.* t. 17. f. 8. opt.

Angl. Bulbous Meadow-grass.

In pratis et pascuis. *Huds.*

Near Clapham in Surrey. *Huds. ed.* 1.

Perennis. Fl. Maio, Junio.

Radix fibrosa, albida, “ex quâ (ut verbis Scheuchzerianis aptissimi utar) confertim plures nascuntur quidam veluti bulbuli, tunicis albicantibus, aut ex albedo fuscis, membranaceis, pluribus, quæ foliorum vaginæ sunt, ad unciam ferè unam exporrectis, vestiti, qui vel foliorum fasciculos, vel etiam eulmos, producunt.” Culmus pedalis, tenuis, glaber, erectus, foliosus. Folia brevia, lincaria, eanaliculata, glaucescentia, margine ferrulato-scabra; caulina brevissima, vaginis longis, striatis. Stipula brevis, obtusiuscula. Panicula ovata, ramulis subflexuosis. Spiculæ latè ovatæ, compressæ, glauæ, xerampelino variegatæ, sæpius quadrifloræ. Glumæ calycinæ ovatæ, trinerves, carinâ apice scabrâ, margine sericeæ, albidæ. Flosculi lanccolati, marginem versus sericei, basi villis complicatis connexi.

Hanc speciem nondum in Angliâ vidi; fide Hudsoni hîc admisi, et ex Linnæano exemplario descripsi. Varietas

β Linnæi diversa species est, at utraque a *P. alpinâ* distinctissima. *Conf. With.* 143.

11. POA paniculâ diffusâ, spiculis ovatis quinque-*cæsia.*
floris, glumis lanceolatis subsericeis liberis,
stipulis brevissimis obtusis.

Poa glauca. With. 148?

Ang. Sea-green Meadow-grass.

In Scotiâ. *D. Fairbairn.*

Mountains in Bredalbane. *Mr. Mackay.*

Perennis. Fl. Junio, Julio.

Radix fibrosa, subcæspitosa. Herba glauca. Culmus pedalis, erectus, teres, striatus, læviusculus, basin versus binodis et foliosus, supernè simplex, rectus, nudus. Folia lanceolata, obtusiuscula, plana, dorso basin versus lævigata, cæteroquin scabra. Vaginæ longitudine foliorum, striatæ, subscabræ. Stipula valdè variabilis, aliquando brevissima, vaginâ inclusâ, quandoque elongata, lanceolata, lacera, præcipuè ad folia caulina. Panicula erecta, diffusa, ramosissima, parùm elongata, ramis semiverticillatis, angulatis, scabris, ramulis binatis. Spiculæ ovatæ, erectæ, sæpiùs quinquefloræ, glauco albido purpureoque variæ. Glumæ calycinæ subæquales, lanceolatæ, acutæ, carinatæ, obsoletè trinerves, apicem versus scabræ. Flosculi remotiusculi, lanceolati, obtusi, apice scariosi, erosi, dorso lanati, margine scariosi, sericei, basi liberi; valvulis interioribus margine scabris; rachi flexuosâ, villosâ.

Diversissima videtur a Poâ glaucâ Vahlîi in *Flo. Dan.* t. 964, quæ proprior est *P. nemorali.*

12. POA paniculâ diffusâ, spiculis trifloris, glumis *trivialis.*
lanceolatis quinquenervibus basi villo con-
nexis, stipulâ elongatâ.

Poa trivialis. Sp. Pl. 99. *Huds.* 39. *With.* 143.

Relb. 33. *Sibth.* 41. *Curt. Lond. fasc.* 2. t. 6.

P. dubia. *Leers.* 28. t. 6. f. 5.

Gramen pratense paniculatum medium. *Raii*
Syn. 409.

β. *Poa setacea.* *Huds. ed.* 1. 34.

Angl. Roughish Meadow-grass.

In pratis et pascuis, præcipuè humidis, ubique.

Perennis. Fl. a Junio ad Septembrem usque.

Radix fibrosa. Culmi basi decumbentes & radicales, dein erecti, sesquipediales, simplices, teretes, striati, scabriusculi, foliosi, sæpè purpurascens. Folia cretiuscula, flaccida, acutiuscula, carinata, subtus præcipuè margineque scabra. Vaginæ longitudine ferè foliorum, subcompressæ, striatæ, parùm scabræ. Stipula elongata, lanceolata, acuta, quâ notâ, monente Curtio, hæc species a *Poa pratensi* optimè dignoscitur. Panicula elongata, erecta, ramosissima, diffusa, scabra, ramis alternatim decompositis, sub florescentiâ horizontaliter patentibus. Spiculæ ovatæ, sæpius trifloræ, purpurco-virides. Glumæ calycinæ dorso scabræ, longitudine æquales; exterior lineari-lanceolata, acuminata; interior margine dilatata, trinervis. Flosculi obtusiusculi, argutè quinquenerves, apice scariosi, margine dorsoque subsericei, basi villis complicatis, numerosis, connexi; valvulæ interiores margine incrassato, viridi, vix pubescente.

Varietas β gaudet foliis inferioribus involuto-setaceis, vaginis tamen justæ latitudinis; nec ad *P. pratensem* referri potest, ob stipulam lanceolatam acutam. Cum *P. angustifoliâ* Hudsoni a Cel. Withering confundi videtur.

pratensis. 13. *POA* paniculâ diffusâ, spiculis quadrifloris, glumis lanceolatis quinquenervibus villo connexis, stipulâ abbreviatâ obtusâ.

Poa pratensis. *Sp. Pl.* 99. *Huds.* 39. *Willd.* 141.

Relb. 34. *Sibth.* 41. *Curt. Lond. fasc.* 2. t. 5.

Leers. 28. t. 6. f. 4?

Gramen pratense paniculatum majus latiore folio.

Raii Syn. 409.

β. *Poa*

β. *Poa angustifolia*. *Sp. Pl.* 99. *Relb.* 34? *Leers.* 27.
t. 6. f. 3.

Angl. Smooth-stalked Meadow-grass.

In pratis, pascuis et muris ubique.

Perennis. Fl. Maio, Junio.

Radix fibrosa, stolonibus repentibus. Culmi plures, erecti, pedales vel sesquipedales, simplices, teretes, minutè striati, læves, foliosi. Folia patentia, obtusiuscula, carinata, glabra, subinde glauca. Vaginæ longitudine foliorum, striatæ, glabræ. Stipula brevis, obtusa. Panicula ut in præcedente, sed minùs scabra. Spiculæ ovatæ, quadrifloræ, sæpè quinquefloræ, virides, subindè purpurascèntes. Glumæ calycinæ acutæ, dorso scabræ, parùm inæquales, trinerves. Flosculi obtusiusculi, argutè quinquenervi, apice scariosi, dorso scabri, basi villis numerosis, complicatis, longissimis, connexi; valvulæ interiores margine subpubescentes.

Varietas β dignoscitur foliis inferioribus angustissimis, rigidulis, paniculâ minori, vaginis inferioribus scabriusculis.

14. POA paniculâ divaricatâ, spiculis ovatis, flos- *annua.*
culis remotiusculis quinquenervibus liberis,
culmo obliquo compresso.

Poa annua. *Sp. Pl.* 99. *Huds.* 42. *With.* 143.

Relb. 35. *Sibth.* 42. *Curt. Lond. fasc.* 1. t. 6.

Mart. Fl. Rust. t. 98. *Leers.* 29. t. 6. f. 1.

Stillingf. Misc. t. 7.

Gramen pratense minus, seu vulgatissimum.

Raii Syn. 408.

Angl. Annual Meadow-grass.

In pascuis, rudcratis, cultis, arcis ubique.

Annua. Fl. totâ æstate.

Radix fibrosa, multiplex. Culmi numerosi, erectiusculi, graciles, compressi, foliosi, læves, basi radicanter atque ramosi. Folia lincaria, obtusiuscula, plana (subinde transversim crispa vel corrugata), flaccida, lævia, lætè virentia,

virentia, margine scabra. Vaginæ elongatæ, compressæ, glabræ, striatæ, pallescentes. Stipula brevis, subdeltoidea, erosa. Panícula erecta, glabra, ramis binatis, divaricatis, subsecundis, parùm ramosis. Spiculæ ovato-oblongæ, nitidæ, viridi alboque variegatæ, glabræ. Glumæ calycinæ lanceolatæ, acutæ, carinatæ, inæquales. Flosculi remotiusculi, ovato-lanceolati, obtusi, quinquenerves, margine membranacei, dorso subsericei, basi omninò liberi, nec pilis connexi.

“In alpinis vivipara evadit.” *D. Davies.*

nemoralis. 15. **POA** paniculâ foliisque attenuatis, spiculis lanceolatis subtrifloris, glumis acutis obsoletè quinquenervibus, stipulâ brevissimâ crenatâ.

Poa nemoralis. *Sp. Pl.* 102. *With.* 146. *Relb.* 36. *Sibth.* 43. *Leers.* 30. *t.* 5. *f.* 3. *Fl. Dan.* *t.* 749.

P. angustifolia β . *Huds.* 41.

Gramen paniculatum angustifolium alpinum, locustis rarioribus & angustioribus, non aristatis. *Scheuchz. Prod.* 18. *t.* 2.

β . *Poa angustifolia* α . *Huds.* 40.

P. pratensis, var. 2. *With.* 142. ex *Herb.* *D. Woodward.*

P. trivialis. *Leers.* 26. *t.* 6. *f.* 2?

Gramen paniculatum pratense majus, angustiore folio. *Morif. sect.* 8. *t.* 5. *f.* 19.

Angl. Wood Meadow-grass.

In nemorosis.

Most plentiful in the north of England.

Perennis. *Fl.* Junio.

Radix parùm repens. Culmi plures, erecti, sesquipedales, graciles, subcompressi, striati, glabri, foliosi. Folia omnia ferè caulina, angusta, attenuato-setacea, acuta, striata; superiora utrinque margineque aspera; inferiora subtus glabrata. Vaginæ vix longitudine foliorum, compressæ, glabriusculæ. Stipula brevissima, crenata.
Panícula

Panicula erectiuscula, diffusa, ramis gracilibus, semi-verticillatis, angulatis, scabris, sæpè flexuosis, magis vel minùs ramosis. Spiculæ lanceolatæ, acutæ, nitidiusculæ, glabræ. Glumæ calycinæ subæquales, spiculâ parùm breviores, lanceolatæ, acuminatæ, trinerves, carinâ apicem versus scabræ. Flosculi 2 ad 4, approximati, lanceolati, acuti, vix manifestè quinquenerves, margine apiceque membranacei, dorso magis minùsve sericei, basi subpilosi, sed liberi, neque villis congestis protrahendis connexi; valvulæ interiores acutæ, intùs carinatæ, margine subpubescentes.

Varietas β gaudet habitu firmiori, caule foliisque minùs attenuatis, paniculâ densiori, flosculis magis plerumque numerosis; faciem verò habet Poæ pratensis vel trivialis. Dignoscitur tamen flosculis liberis, basi licet pilosis, acutioribus et minùs nervosis, stipulâ brevissimâ, subcrenatâ vel ciliatâ, adeo ut eum illis associari nequeat.

16. POA paniculâ subsimplici coarctatâ erectâ, *decumbens*.
spiculis ovatis quadrifloris longitudine calycis, *bens*.
stipulis ciliaribus.

Poa decumbens. *With.* 147.

Festuca decumbens. *Sp. Pl.* 1110. *Huds.* 47.

Relb. 43. *Sibth.* 46. *Fl. Dan.* t. 162.

Leers. 34. t. 7. f. 5. *Dicks.* H. *Sicc. fasc.*

11. 3.

Melica decumbens. *Web. Gott.* 3.

Gramen avenaceum parvum procumbens, paniculis non aristatis. *Raii Syn.* 408.

G. triticeum palustre humilius, spicâ breviori.

Morif. sect. 8. t. 1. f. 6.

Angl. Decumbent Meadow-grass.

In pascuis sterilibus humidiusculis, præcipuè montosis spongiosis.

Perennis. Fl. Julio.

Radix fibrosa, ramosa, subrepens. Culmus decumbens, sub florescentiâ erectus, pedalis, geniculatus, rigidus, glaberrimus,

glaberrimus, foliaceus. Folia linearia, obtusiuscula, involuta, rigidula, striata, dorso apicem versus asperrima. Vaginæ striatæ, apice præcipuè pilosæ. Stipula brevissima, e ciliis densissimis, æqualibus, erectis. Panícula erecta, subsimplex, pauciflora, rachi et pedunculis flexuosis, angulatis, scabriusculis. Spiculæ ovatæ, tumidæ, acutæ, glabræ, sæpiùs purpurascentes. Glumæ calycinæ longitudine circiter ipsius spiculæ, vel parùm longiores, subæquales, ovatæ, acutæ, obsoletè nervosæ, carinâ scabræ. Flosculi 3 ad 5, arctè imbricati, elliptici, glaberrimi, nitidi, obsoletè multinerves, basi pilosi, liberi, margine sericei, apice trifidi, nec aristati; glumis interioribus ellipticis, obtusis, margine ciliatis.

34. BRIZA. *G. Pl.* 35. *Juss.* 32. *Gærtn. t.* 1.

Cal. bivalvis, multiflorus. *Spicula* disticha. *Cor.* bivalvis, ventricosa, valvulis cordatis obtusis. *Semen* corollæ adnatum, depresso.

minor.

1. BRIZA spiculis triangularibus septemfloris, calyce flosculis longiore, stipulâ lanceolatâ longissimâ.

Briza minor. Sp. Pl. 102. *Huds.* 38. *With.* 148. *Dicks. H. Sicc. fasc.* 5. 3.

Gramen tremulum minus, paniculâ amplâ, locustis parvis triangulis. *Raii Syn.* 412.

G. tremulum minus, locustâ deltoide. Moris. sect. 8. *t.* 6. *f.* 47.

Angl. Small Quaking-grass.

In arvis Angliæ australioris rarissimè.

Near Bath. *Mr. Alchorne. Huds.* In cornfields between Pensance and Marketjew, Cornwall, 1774. *Lightf. in Herb.* Jersey. *Sherard.* Guernsey. *Yulden.*

Annua. Fl. Julio.

Radix fibrosa, parva. Culmus erectus, spithamæus, teres, glaberrimus, foliosus, basi sæpiùs ramosus. Folia vaginantia, erecta, lanceolata, acuta, plana, lætè viridia, striata,

friata, margine scabra. Vagina friata, glabra. Stipula lanceolata, longissima, amplexicaulis, decurrens, supernè folio adnata, tenuissima. Panicula diffusa, patentissima, multiflora, ramis binatis, ramulis dichotomis, divaricatis, capillaribus, scabris, viridibus. Spiculæ pendulæ, tremulæ, deltoideæ, glabræ, albo viridique variæ, subseptemfloræ. Glumæ calycinæ subæquales, concavæ, obtusissimæ, friatæ, margine membranaceæ, carinâ læves, flosculis longiores. Flosculi calyci conformes, alterni, sensim minores; glumis interioribus longè minoribus, emarginatis.

2. BRIZA spiculis ovatis septemfloris, calyce flos- *media.*
culis brevior, stipulâ brevissimâ obtusâ.

Briza media. *Sp. Pl.* 103. *Huds.* 38. *Relb.* 38.
Sibth. 43. *Engl. Bot. t.* 340. *Mart. Fl. Rust.*
t. 39. *Dicks. H. Sicc. fasc.* 5. 2. *Leers.* 25.
t. 7. *f.* 2. *Fl. Dan. t.* 258.

Gramen tremulum. *Ruii Syn.* 412.

G. tremulum vulgare minus, Phalaris minor Lobelii. *Morif. sect.* 8. *t.* 6. *f.* 45.

Phalaris pratensis. *Ger. em.* 86.

Angl. Common Quaking-grass.

In pascuis frequens.

Perennis. Fl. Maio, Junio.

Radix fibrosa, elongata. Culmus pedalis et ultra, erectus, teres, foliosus, glaberrimus. Folia vaginantia, erectiuscula, lanceolata, acuta, plana, striata, scabriuscula. Vaginæ caulinae longissimæ, striatæ, glabræ. Stipula brevissima, obtusâ, vix decurrens, neque folio supernè adnata, quibus notis a B. minore et a virenti abundè differt. Panicula diffusa, patens, multiflora, ramis binatis, ramulis ut in priore, sed fuscis nec viridibus. Spiculæ pendulæ, tremulæ, ovatæ vel subcordatæ, glabræ, nitidæ, albo fuscoque variæ, septem- (subinde novem-) floræ. Glumæ calycinæ subæquales, concavæ, obtusæ, vix striatæ, margine scariosæ, carinâ læves, flosculis parùm breviores. Flosculi calyci conformes, alternati,

alternatim, ut in præcedente, digesti ; glumis interioribus etiam emarginatis.

35. DACTYLIS. *G. Pl.* 35. *Juss.* 31.

Cal. bivalvis, compressus ; alterâ valvulâ majore, carinatâ.

stricta. 1. DACTYLIS spicis geminis erectis lævibus, spiculis secundis pubescentibus.

Dactylis stricta. *Soland. in Ait. Hort. Kew.* v. 1. 104. *Engl. Bot. t.* 380. *With.* 149. t. 27.

D. cynosuroides. *Huds.* 43. *With. ed.* 2. 93. *Loefling. It. Hisp.* 115.

Spartum Effexianum, spicâ geminâ clausâ. *Dill. in Raii Syn.* 393.

Angl. Smooth Cock's-foot-grass.

In pratis limosis maritimis humilioribus.

About the mouths of rivers in Kent, Essex and Suffolk principally. In the Isle of Shepey plentifully. *Rev. Dr. Goodenough.*

Perennis. Fl. Augusto.

Radix repens, fibris longissimis. *Culmus erectus*, subpedalis, rigidus, simplex, teres, glaberrimus, foliosus. *Folia alterna*, vaginantia, rigida, involuta, apice subulato-mucronata, glabriuscula, supernè profundius striata, dorso læviora. *Vaginae culmum* arctè amplectentes, cylindricæ, striatæ, intùs nitidæ, glabræ, foliis delapsis persistentes, dein parùm divaricatæ et subcarinatæ. *Stipula brevissima*, erosa. *Spicæ geminæ*, rariùs ternæ, erectæ, subæquales, sibi invicem dorso applicatæ, simplices, rachi angulatâ, striatâ, lævi. *Spiculæ secundæ*, duplici serie alternatim imbricatæ, sessiles, erecto-adpressæ, unifloræ. *Glumæ calycinæ lineares*, acuminatæ, complicato-carinatæ, pubescentes ; interior brevior et multò angustior, glabra. *Petala subæqualia*, calyce majora, carinata, acutiuscula. *Antheræ lineares*, exsertæ.

exfertæ. Styli basi connati, stigmatibus subulatis, villosis.

A D. cynosuroide abunde differt spicis paucissimis, neque numerosis vel paniculato-spicatis, rachi spiculisque inermibus, nec spinoso-exasperatis.

2. DACTYLIS paniculâ secundâ supradecompositâ *glomeratâ* glomeratâ.

Dactylis glomerata. Sp. Pl. 105. Hudf. 43.

With. 150. Relb. 39. Sibth. 43. Schreb.

Gram. 72. t. 8. f. 2. Engl. Bot. t. 335.

Mart. Fl. Rust. t. 14. Leers. 21. t. 3. f. 3.

Gramen asperum. Raii Syn. 400.

Angl. Rough Cock's-foot-grass.

In pratis et umbrosis vulgaris.

Perennis. Fl. Junio—Augusto.

Radix fibrosa, subcæspitosa. Culmus erectus, bipedalis, teres, striatus, supernè nudus, scabriusculus. Folia linearia, acuta, patentia, longa, obscurè viridia, striata, margine præcipuè scabra. Vaginæ striatæ, compressæ, scabræ. Stipula elongata, lacera. Panicula alternitè ramosa, ramis remotis, angulatis, scaberrimis, rigidis, decompositis. Spiculæ secundæ, conglomeratæ, scabræ. Glumæ calycinæ maximè inæquales, carinatæ, acuminatæ; exterior trinervis, carinâ ciliatâ; interior membranacea, glabra. Flosculi tres vel quatuor, rariùs solitarii, calyce majores, quinquenerves, mucronati, carinâ pectinato-ciliati; gluma interior bifida, margine ciliata. Antheræ purpureæ, exfertæ, tremulæ.

36. CYNOSURUS. *G. Pl. 36. Juss. 31. Gærtn. t. 1.*

Cal. bivalvis, multiflorus. Receptaculum proprium unilaterale, foliaceum.

1. CYNOSURUS bracteis pinnato-distichis *multicristatus*, tics, spicâ simplici lineari.

Cynosurus

Cynosurus cristatus. *Sp. Pl.* 105. *Huds.* 59.
With. 150. *Relb.* 39. *Sibth.* 52. *Schreb.*
Gram. 69. *t.* 8. *f.* 1. *Engl. Bot.* *t.* 316.
Mart. Fl. Rus. *t.* 106. *Leers.* 49. *t.* 7. *f.* 4.
Gramen cristatum. *Ger. em.* 29.* *Raii Syn.* 398.
Angl. Crested Dog's-tail-grass.

In pascuis ficcioribus vulgaris.

Perennis. *Fl.* Julio.

Radix fibrosa, caespitosa. *Culmi* plures, erecti, stricti, pedales, foliosi; supernè nudi, teretes, striati, glaberrimi. *Folia* linearia, acuta, lævia. *Vaginæ* teretes, striatæ, glaberrimæ. *Stipula* brevis, obtusa, erosa. *Spica* simplex, linearis, obtusa, stricta, secunda, rachi subflexuosa, angulatâ, lævi. *Spiculæ* alternæ, ovatæ, multifloræ, subsessiles. *Glumæ* calycinæ lineares, acutæ, carinatæ, carinâ scabræ. *Flosculi* alterni, tumiduli, vix carinati, apicem versus scabri, mucronati. *Bractææ* pectinatæ, e glumis (monente Stokesio) abortivis, alternis, distichis, compresso-carinatis, dorso scabris, acutis, vix aristatis.

Varietatem tetrastricham, tetragonam, descripsit Raius in *Syn.* p. 399.

Quandoque vivipara evadit.

echina-
tus.

2. *CYNOSURUS* bracteis pinnatis scariosis longissimè aristatis, spicâ compositâ ovatâ.

Cynosurus echinatus. *Sp. Pl.* 105. *Huds.* 59.
With. 150.

Gramen alopecuroides spicâ asperâ brevi. *Raii Syn.* 397. *Barell. Ic.* *t.* 123. *f.* 2.

G. panicum, spicâ asperâ latiore. *Morif. sect.* 8. *t.* 4. *f.* 13.

Angl. Rough Dog's-tail-grass.

In arenosis, sub cælo mitiore.

On all the sandy grounds of Jersey. *Sherard.*
 Near Sandwich. *Hudson.* Near Hastings, but very sparingly. *Rev. Dr. Goodenough.*

Annua.

Annua. Fl. Julio.

Radix fibrosa, parva. Culmus erectus, foliosus, teres, striatus, lævis, basi ramosus. Folia lineari-lanceolata, acuta, letè virentia, striata, utrinque scabra. Vaginæ parum inflatæ, subancipites, scabriusculæ, fulcatæ. Stipula lanceolata, membranacea. Panicula ovata, conglomerata, ramosa. Spiculæ alternæ, ovatæ, pallidæ, scabræ. Glumæ calycinæ æquales, tenuissimæ, membranaceæ, acutæ, scabræ. Flosculi plures, apicem versus scabri, longissimè aristati, aristâ scabrâ, purpurascente; valvulis interioribus muticis. Bractææ glumis alternis, lanceolatis, nervosis, scabris, flosculorum more aristatis.

37. FESTUCA. *G. Pl.* 36. *Juss.* 32.

Cal. bivalvis. *Spicula* oblonga, teretiuscula, disticha; glumis acuminatis.

1. FESTUCA paniculâ secundâ coarctatâ, floscu-*ovina* lis teretiusculis basi glabris, culmo tetragono, foliis setaceis scabris.

Festuca ovina. *Sp. Pl.* 108. *Huds.* 44. *With.* 152. *Relb.* 40. *Sibth.* 43. *Engl. Bot.* t. 585. *Mart. Fl. Rust.* t. 102. *Leers.* 32. t. 8. f. 3.

Gramen foliolis junceis brevibus, majus, radice nigrâ. *Scheuchz. Agr.* 279. t. 6. f. 8.

β. *Festuca rubra.* *With.* 153. (nec Linnæi), ex ipso auctore.

γ. *F. tenuifolia.* *Sibth.* 44. *With.* 155.

F. ovina β. *Leers.* 33. t. 8. f. 4.

Gramen capillaceum, locustellis pennatis non aristatis. *Raii Syn.* 410. *Pluk. Phyt.* t. 34. f. 2.

G. loliaceum, foliolis junceis brevibus minus. *Morif. sect.* 8. t. 3.

Angl. Sheep's Fescue-grass.

In pascuis ficcioribus, collibusque apricis frequens.

Var. γ . vulgarissima est.

β . In the New Forest, Hampshire. *Dr. Withering*.

Perennis. Fl. Junio.

Radix fibrosa, cæspitosa, fibris capillaribus, nigris, glabris. Culmi spithamæi, erecti, simplices, tenues, rigiduli, glabri, basi foliosi, apicem versus præcipuè tetragoni. Folia longitudine et directione varia, tenuissima, angulato-setacea, acuta, scabra, sæpe glaucescentia. Vaginæ striatæ, glabræ. Stipula brevissima. Panícula lanceolata, secunda, ramosa, rachi pedunculisque angulatis, subflexuosis, scabris. Spiculæ erectæ, ovato-lanceolatæ, glabræ. Glumæ calycinæ inæquales, lincarilanceolatæ, acuminatæ, carinatæ, angulatæ, glabræ. Flosculi 4 vel 5, remotiusculi, teretiusculi, parum compressi vix carinati, enerves, glabri, basi lævissimi, apicem versus tuberculis minutis scabri, sæpius mutici, quandoque (ut in herbario Linnæano) aristati, aristis scabris, rectis, longitudine variis. Valvula interior mutica, acutiuscula, concava, margine omnino glabro, subrevoluto.

β . differt colore tantum paniculæ purpurascente. Flosculi in γ mutici; folia verò erecta subflaccida, vel rigida incurva, nullæ varietati peculiaria sunt.

vivipara. 2. FĚSTUCA paniculâ secundâ coarctatâ, flosculis compresso-carinatis muticis calyceque subpubescentibus, culmo tetragono, foliis setaccis lævibus.

Festuca ovina β . *Sp. Pl.* 108. *Huds.* 44. *Lightf.* 101. *With.* (var. 4.) 152.

Gramen sparteum montanum, spicâ foliaceâ gramineâ, majus et minus. *Raii Syn.* 410. t. 22. f. 1.

G. paniculatum sparteum alpinum, paniculâ angustâ, spadiceo-viridi, proliferum. *Scheuchz. Prod.* 21. t. 1. f. 2.

Angl. Viviparous Fescue-grass.

In

In cacuminibus montium altissimorum.

On Ingleborough, Skiddaw, Snowdon, and on most of the Scottish mountains.

Perennis. Fl. Julio.

Radix et culmus ut in præcedente. Folia recurva, angulato-setacea, minus rigida, et omnino lævia, lætè viridia. Panicula abbreviata, subsimplex, erecta, secunda, rachi pedunculisque angulatis, scabris. Spiculæ erectæ, ovato-lanceolatæ, plerumque mollissimè pubescentes. Glumæ calycinæ valdè inæquales, lanceolatæ, acuminatæ, carinatæ, angulatæ, pubescentes. Flosculi 4 vel 5, arctè imbricati, lanceolati, acuminati, mutici, acutè carinati, magis vel minus compressi, angulati, nervosi, toti pubescentes, rariùs glabri; superiores 2 vel 3, metamorphosi grumarum in folia, vivipari. Glumæ interiores margine nunquam non pubescunt.

In hortis, monente Linnæo, ut et Witheringio ex observationibus D. Gough, perpetuò vivipara manet. Videtur species a priore distincta.

3. *FESTUCA* paniculâ secundâ diffusâ, flosculis *durius-*
aristatis, culmo tereti, foliis caulinis planis, *cula,*
radice fibrosa.

Festuca duriuscula. *Sp. Pl.* 108. *Huds.* 44.

With. 153. *Relb.* 40. *Sibth.* 44. *Leers.* 33.
t. 8. *f.* 2. *Engl. Bot. t.* 470.

Gramen pratense, paniculâ duriore laxâ, unam
partem spectante. *Raii Syn.* 413. *t.* 19. *f.* 1.

β . *Huds.* 45.

Festuca dumetorum. *Sp. Pl.* 109. *With.* 154.

Fl. Ddn. t. 700.

Angl. Hard Fescue-grass.

In pratis et pascuis ficcioribus ubique. β in dume-
tis et ad sepes.

Perennis. Fl. Junio.

Radix fibrosa, fusca, villosa. Culmi pedales vel bipedales,
erecti, simplices, foliosi, glabri, teretes, striati, quan-
doque

doque subcompressi. Folia longitudine varia, angusta, acuta; radicalia setacea, scabra; caulina plana, lævia, subinde supra pubescentia. Vaginæ striatæ, glabræ, longissimæ. Stipula minuta. Panicula ferè Festucæ ovinæ, at omni parte major, floribusque magis carinatis, semper aristatis, vix apice scabris, glumis interioribus margine pubescentibus.

Flosculi in β mollissimè pubescunt, calyce semper glabro.

rubra. 4. **FESTUCA** paniculâ secundâ, flosculis teretiusculis aristatis, foliis supra pubescentibus, radice repente.

Festuca rubra. *Sp. Pl.* 109. *Huds.* 45. *Lightf.* 102. *Stillingfl. Misc. t.* 9.

Gramen alpinum pratense, paniculâ duriorè laxâ spadiceâ, locustis majoribus. *Scheuchz. Agr.* 287. *t.* 6. *f.* 9. ex descr.

β . *Festuca cambrica.* *Huds.* 45. *With.* 155.

γ . *F. glabra.* *Lightf.* 1085. *Huds.* 648. *With.* 154. *Angl.* Creeping Fescue-grass.

In pascuis montosis et cacuminibus alpinis, etiam in arenosis maritimis.

α . At Yarmouth. *Mr. Woodward.* Scarborough. *Mr. Teesdale.* Exmouth sands, Devonshire. *Mr. Newberry,* and *Rev. Dr. Goodenough.* Dumfries-shire. *Mr. J. Burges's Mac-Garrock.* Near Kirkby Lonsdale, Westmoreland, 1783.

β . On the ledges of Clogwin y Garnedth, the highest rock of Snowdon. *Herb. Banks.*

γ . At Ardbiglen in Galloway. *Herb. Lightf.*

Perennis. Fl. Julio.

Radix repens, flagellis longissimè (in maritimis) per arenam mobilem extensis, filiformibus, glabris, hinc infimis fibris nigris radicantibus. Culmi subpedales, decumbentes, simplices, basi præcipuè foliosi, teretiusculi, glaberrimi, striati. Folia linearia, demum involuta, acuta, subpungentia; subtus lævia, vix carinata; supra fulcata, pubescentia. Vaginæ fulcatæ; caulinae longissimæ,

gissimæ, glabræ; radicales abbreviatæ, dilatatæ, magis vel minus retrorsum pubescentes. Stipula brevissima, erosa. Panicula secunda, ramosa, rachis pedunculisque sulcatis, angulis scabris. Spicæ erectiusculæ, ovatæ, purpurascens; in maritimis glaucescentes. Glumæ calycinæ lineari-lanceolatæ, acutæ, nervosæ, inæquales, glabræ, rarius carinâ vel margine ciliatæ. Flosculi plerumque quinque vel sex, parum distantes, lanceolati, subcompressi, vix carinati, brevius aristati, sæpius apice margineque tantum pubescentes, dorso glaberrimo, at interdum in maritimis undique villosi, ut in *F. duriuscula* β , dumetorum. Glumæ interiores parum breviores, obtusiusculæ, margine pubescentes vel ciliatæ ut in *Bromis*.

Dignoscitur radicibus latè repentibus. Var. β gaudet paniculâ simpliciore, flosculis majoribus, latioribus, ferè ovatis, vel omnino glabris, vel margine pubescentibus, foliis suprà minus pubescentibus, non verò glabris.

Spiculæ in γ trifloræ, flosculis plerumque glaberrimis.

5. *FESTUCA* paniculâ secundâ erectâ, flosculis *bromoides*.
subulatis aristatis apice scabris, foliis setaceis *oides*.
vaginâ brevioribus.

Festuca bromoides. *Sp. Pl.* 110. *Huds.* 46.

With. 151. *Sibth.* 44.

Bromus dertonensis. *Allion. Ped. v.* 2. 249.

Gramen paniculatum bromoides minus, paniculis aristatis, unam partem spectantibus. *Raii*

Syn. 415. *Pluk. Phyt. t.* 33. *f.* 10. *Rel.*

Rudb. 17. *f.* 3.

G. bromoides, paniculâ heteromallâ, longioribus aristis donatâ. *Scheuchz. Agr.* 290. *t.* 6. *f.* 10.

Herb. Cher. ex ipso Scheuchzero.

Angl. Barren Fescue-grass.

In muris, et locis arenosis sterilioribus.

Annua. Fl. Junio.

Radix fibrosa, cæspitosa, ramosissima, fusca. Culmi plurimi, spithamæi vel pedales, erecti, teretes, glaberrimi,

rimi, basin versus foliosi, geniculis 3 vel 4, fuscis. Folia pallidè viridia, erecta, setacea, angulata, acuta, lævia, brevia. Vaginæ foliis longiores, subventricosæ, fuscatae, glabræ. Stipula exigua. Panicula erecta, uncialis, vix unquam biuncialis, secunda, spicata, basi subramosa; rachi pedunculisque angulatis scabriusculis. Spiculæ erectæ, glabræ, ovatae. Glumæ calycinæ acutæ, glabræ; exterior angustissima, ferè setacea; interior duplò longior et triplò ferè latior, trinervis. Flosculi circiter sex, remotiusculi, subulati, parùm compressi, ecarinati, apicem versus tuberculis exasperati, aristati, aristis rectis, scabris, valvulâ longioribus. Valvula interior acutiuscula, margine pubescens. Semen lineare, sulco exaratum.

Magnitudine variat infinitè.

myurus. 6. *FESTUCA* paniculâ secundâ nutante elongatâ, flosculis subulatis aristatis apice scabris, foliis setaceo-subcarinatis brevissimis.

Festuca myurus, *Sp. Pl.* 109. *Huds.* 46. *With.* 151. *Relb.* 41. *Sibth.* 44. *Dicks.* *H. Sicc. fasc.* 10. 2. *Leers.* 34. *t.* 3. *f.* 5.

Gramen murorum, spicâ longissimâ. *Raii Syn.* 415. *Ger. em.* 29.*

G. avenaceum murorum, spicâ longissimâ nutante aristatâ, *Morif. sect.* 8. *t.* 7. *f.* 43.

Angl. Wall *Fescue-grass.*

In muris, et locis sterilibus.

Annua. Fl. Junio.

Præcedenti proxima, at distincta. Differt foliis minùs setaceis, culmò plerumque elatiori, paniculâ quadruplò longiore, ramosâ atque nutante. Spiculis flosculisque cum priore omninò convenit. Variat glumis pilosis, sed rariùs.

uniglumis. 7. *FESTUCA* paniculâ secundâ erectâ subsimplici, flosculis subulatis compressis aristatis, calycis valvulâ alterâ brevissimâ.

Festuca

Festuca uniglumis. *Soland. in Ait. Hort. Kew.*
v. 1. 108.

Stipa membranacea. *Sp. Pl.* 116? *Mant.* 2.
326?

Lolium bromoides. *Hudsf.* 55. *With.* 169.

Festuca avenacea sterilis humillima, spicâ unam
partem spectante. *Petiv. Conc. Gram.* 101.
Dill. in Raii Syn. 413. *sub n.* 3. *t.* 17. *f.* 2,
exclusis omninò citationibus *Syn. ed.* 2, quæ
ad *Bromum diandrum* pertinent.

Angl. Single-husked Fescue grass.

In arenosis maritimis.

In Mersey island, near Colchester. *Dale.* On the
coast of Suffex. *Dillenius.* At Little Hamp-
ton, Suffex; also near Landguard Fort. *Rev.*
Dr. Goodenough.

Annua, vel Biennis. Fl. Junio.

Radix fibrosa. Culmi plures, spithamæi, erectiusculi, an-
gulati, læves, foliosi, basi ramosi. Folia lanceolata,
acuta, subinvoluta, striata, lævia. Vaginæ foliis lon-
giores, ventricosæ, striatæ, glabræ. Stipula brevis, ob-
tusa. Panícula erecta, sesqu-uncialis, secunda, basiparum
ramosa, supernè spicata; rachi pedunculisque compressis,
angulis scabris. Spiculæ erectæ, lanceolatæ, compressæ.
Gluma calycina exterior longitudine ferè spiculæ, subu-
lata, compressa, acuminata, carinata, trinervis, apice ca-
rinâque scabra; interior minutissima, vix nisi oculo
armato cernenda, albida. Flosculi 5 ad 8, subulato-
triquetri, carinati, aristati, scabri, remoti, superiores
abortivi. Aristæ flosculis duplò longiores, rectæ, ca-
pillares, scabræ. Valvulæ interiores lineares, angustif-
simæ, acutæ, margine scabræ.

Præcedentibus speciëbus valde affinis est, nec ullam cum
Lolio habet affinitatem vel similitudinem.

Stipa membranacea Linnæi, ad *Festucas* proculdubio re-
vocanda, huic quam maximè affinis est, et ferè eadem
species; vix enim discrepat nisi habitu laxiore magis-
que luxuriante pedunculis longitudinaliter quasi sulcatis,

glumâ calycinâ interiore magis aliquantulum conspicuâ.

gigantea. 8. *FESTUCA* paniculâ secundâ nutante ramosâ laxâ, flosculis lanceolatis ventricosis aristatis, foliis ensiformibus nervosis.

Festuca gigantea. *Villars. Dauph. v. 2. 110.*

Bromus giganteus. *Sp. Pl. 114. Hudf. 51.*

With. 162. Relb. 47. Sibth. 48. Curt.

Lond. fasc. 5. t. 7. Schreb. Gram. 88. t. 11.

Fl. Dan. t. 440. Leers. 39. t. 10. f. 1.

B. n. 1510. Hall. Hist. v. 2. 238.

Gramen avenaceum glabrum, paniculâ e spicis raris strigosis compositâ, aristis tenuissimis.

Raii Syn. 415.

G. avenaceum paniculatum Gallo-Provinciale Aquilegiæ semine. Moris. vol. 3. 214 n. 29.
ex *Herb. Bobart. N. 30* in eodem herbario non reperitur.

G. sylvaticum glabrum paniculâ recurvâ. Vaill. Paris. t. 18. f. 3.

G. bromoides aquaticum latifolium, paniculâ sparsâ tenuissimè aristatâ. Scheuchz. Agr. 264. t. 5. f. 17.

Festuca avenacea sterilis elatior, spicis aristatis in gyrum contortis. Rel. Rudb. 17. f. 2.

Angl. Tall Fescue-grass.

In sylvis et sepibus humidiusculis.

Perennis. Fl. Julio, Augusto.

Radix fibrosa. Culmus tri- vel quadripedalis, erectus, teres, lævis, striatus, foliosus. Folia erectiuscula, ensiformia, acuminatâ, pedalia, lata, plana, nervosa, utrinque margineque scabra. Vaginæ longæ, striatæ, glabræ, nec pilosæ. Stipula brevis, erosa, amplexicaulis, purpurascens. Panicula elongata, nutans, ramosa, multiflora, rachi ramulisque angulatis scabris. Spiculæ alternæ,

alternæ, ovato-lanceolatæ, glabræ, nutantes. Glumæ calycinæ in quales, acuminatæ, glabræ, carinatæ; interior latior et trinervis. Flosculi 4 ad 6, imbricati, glabri, ovato-lanceolati, vix carinati, apice quinque-nerves. Arista terminalis, flosculo duplo vel triplo longior, capillaris, scabra, albida, quandoque subflexuosa. Gluma interior longitudine et ferè latitudine exterioris, acuta, concaviuscula, margine primo intuitu glabra, sed oculo armato scabra ut in plurimis Festueis, nec ciliata ut in Bromis.

Gluma interior non pectinato-ciliata, arista, monente Villarsio, terminalis, radix insuper perennis, hanc speciem a Bromis ad Festueas amandari suadent. A Bromo aspero distinctissima est.

9. FESTUCA paniculâ secundâ erectâ ramosissimâ coarctatâ, flosculis oblongis angulatis muticis, foliis ensiformibus striatis. *calamaria.*

Festuca sylvatica. Villars. *Dauph.* v. 1. 271, v. 2. 105?

Poa sylvatica. Pollich. v. 1. 83, ex descr. optimâ; excluso reverà synonymo Halleri.

Gramen paniculatum nemorosum latifolium glabrum, paniculâ nutante non aristatâ. Till. *Pis.* 45. Herb. Sherard. ex Tillio & Michelio.

Angl. Reed-like Fescue-grass.

In sylvis subalpinis Scotiæ.

In a moist wooded valley at the foot of Ben Lawers, 1793. Mr. Mackay.

Perennis. Fl. Julio.

Radix fibrosa, cæspitosa. Culmi plures, erecti; tripedales, arundinacei, stricti, teretes, striati, glaberrimi, foliosi, geniculis constrictis, purpurascentibus. Folia erecta, plana, ensiformia, acuminata, striata, margine præcipuè scabra. Vaginæ longitudine plerumque foliorum, cylindricæ, strictæ, striatæ, scabriusculæ, minimè pilosæ vel hirtæ. Stipula breviuscula, obtusa, erosa. Panicula

nicula pro ratione plantæ parva, erecta, secunda, ramossissima, multiflora. Rami binati vel ternati, angulati, scabri. Spicula parvæ, congestæ, erectæ, ovato-oblongæ, compressiusculæ, albido-virescentes, sæpè subpurpurascens, oculo armato subpubescentes. Glumæ calycinæ setacæ, acuminatæ, carinatæ, læves, parùm inæquales. Flosculi tres ad quinque, distichi, remotiusculi, lineari-oblongi, angulato-teretes, glumis subæqualibus, convexis, lanceolatis, acuminatis, muticis, scariosis, compresso-carinatis, carinâ obsolete scabris. Antheræ lineares, flavæ.—Peculiares sunt glumæ interiores exteriorum modo convexæ, carinatæ, margine attenuatæ nudæ, vix concavæ vel depressæ, margineve incrassato, binervi, pubescente.

Hujusce exemplaria e Michelio missa cum Festuca elatiore confudit Dillenius in Herbario Sherardiano; confer *Raii Syn.* 411. 15.

Loliacea. 10. **FESTUCA** spicâ distichâ nutante, spiculis subsessilibus lineari-oblongis, flosculis cylindricis enervibus muticis.

Festuca loliacea. *Huds. ed.* 1. 38. *With.* 157.

Curt. Lond. fasc. 6. t. 9. *Sibth.* 45.

F. fluitans β. *Huds.* 47.

Poa n. 1452. *Hall, Hist.* v. 2. 219.

Graminis loliacci vulgaris, varietas spicis rariùs dispositis. *Morif. sect.* 8. t. 2. f. 2.

Angl. Spiked Fescue-grass.

In pascuis humidis.

Near London not very uncommon. Perhaps frequently overlooked, being mistaken for *Lolium perenne*.

Perennis. Fl. Junio, Julio.

Radix fibrosa. Culmi erecti, bipedales, simplices, teretes, striati, glabri, foliosi. Folia linearia, acuminata, patentia, plana, striata, lævia. Vagina longa, striata, glabra. Stipula brevissima, amplexicaulis. Spica elongata, nutans, disticha, multiflora, basi quandoque subramosa, rachis

rachi angulatâ, hinc inde excavatâ, angulis asperâ. Spiculæ alternæ, plerumque sessiles, obliquè patentés, lineari-oblongæ, acutæ, multifloræ, glabræ; inferiores sæpius pedunculatæ, et frequenter binæ. Glumæ calycinæ muticæ, inæquales; altera subulato-triquetra, brevior; altera longior multoque latior, multinervosa, vix carinata. Flosculi distichi, remotiusculi, acuti, mutici, vix unquam aristati, cylindrici, ecarinati, omninò læves et enerves. Glumæ interiores haud breviores, complanatæ, margine pubescentes.

Dignoscitur ab omni Poæ fluitantis varietate, glumis acutis, enervibus; a Loliiis, calyce bivalvi. Facies Lolii perennis.

11. FESTUCA paniculâ secundâ erectiusculâ *pratensis*. laxâ, spiculis linearibus compressis obtusiusculis, flosculis cylindricis obsoletè nervosis.

Festuca pratensis. *Huds. ed. 1. 37. Curt.*

Lond. fasc. 6. t. 7. Mart. Fl. Rust. t. 84.

F. elatior. *Lim. Fl. Suec. 32. With. 156.*

var. 2. Lightf. 102?

F. fluitans γ. Huds. 47.

Gramen paniculatum elatius, spicis longis muticis squamosis. *Raii Syn. 411. excluso syn. Barrelieri.*

Angl. Meadow Fescue-grass.

In pratis et pascuis humidioribus.

Perennis. Fl. Junio, Julio.

Radix fibrosa. Culmi plures, erecti, sub-bipedales, simplices, teretes, striati, glabri, foliosi. Folia linearia, acuminata, patentia, striata, superiora utrinque scabra. Vaginæ striatæ, lævissimæ. Stipula brevissima, amplexicaulis. Panicula erectiuscula, ramosa, secunda, patens, ramis binatis, inæqualibus, plerumque simplicibus, subracemosis, scabris. Spiculæ brevius pedicellatæ, alternæ, erecto-divaricatæ, lineari-oblongæ, compressæ, obtusiusculæ, multifloræ, glabræ. Glumæ calycinæ inæquales, acutiusculæ, carinatæ, glabræ; alterâ paulò

paulò latiore, trinervi. Flosculi numerosi, parùm remoti, distichi, leves, cylindrici, acuti, basi cecarinati, apicem versus obsolete quinque nervos et subcarinati, plerumque mutici. Glumæ interiores concavæ, vix breviores, apicè laceræ, margine pubescentes.

clatior. 12. **FESTUCA** paniculâ subdiffusâ nutante ramosissimâ laxâ, spiculis ovato-lanceolatis acutis, flosculis cylindricis obsolete nervosis.

Festuca clatior. *Sp. Pl.* 111. *Huds.* 47. *Sibth.* 45. *With.* 155. *Curt. Lond. fasc.* 6. *t.* 7. *Schreb.* *Gram.* 34. *t.* 2. *Leers.* 35. *t.* 8. *f.* 6.

Gramen paniculatum nemorosum, latiore folio, glabrum, paniculâ nutante non aristatâ. *Dill. in Raii Syn.* 411. nec tamen Michellii; confer *F. calamariam*, p. 121.

G. loliaceum, spicâ divisâ, pratense majus. *Morif.* v. 3. 183. *sect.* 8. *t.* 2. *f.* 15.

G. arundinaceum aquaticum, paniculâ avenaceâ. *Raii Syn.* 411. *ed.* 2. 254?

Angl. Tall Fescue-grass.

In pratis humidis, salicetis, et ad fossarum margines.

Near Dover. *Herb. Sherard.* About Battersea and Chelsea. *Curtis.* Near Hastings, under the rocks to the east of the town. *Rev. Dr. Goodenough.*

Perennis. Fl. Junio, Julio.

Radix fibrosa, subrepens, limum profundè penetrans. Culmi tri- vel quadripedales, striati, glabri, foliosi. Folia ut in præcedente, sed duplo latiora atque longiora, nervosa, omnia utrinque glabra, margine tantum scabra. Panicula nutans, demum suberecta, minùs secunda, decomposita, spiculis quam in præcedente duplo numerosioribus, ovatis, acutis, parùm compressis. Reliqua ut in *F. pratensi*, parùm verò majora.

38. BROMUS. *G. Pl.* 36. *Juss.* 32.

Cal. bivalvis. *Spicula* oblonga, disticha: *arista* infra apicem: *gluma* interior pectinato-ciliata.

1. BROMUS paniculâ patente; pedunculis sub-*secalinus*. simplicibus, spiculis ovatis compressis decemfloris, flosculis distinctis teretiusculis.

Bromus secalinus. *Sp. Pl.* 112. *Tr. of Linn. Soc.* v. 4. 281.

B. polymorphus γ. *Huds.* 49. var. 2. *With.* 159.

B. vitiosus. *Weigel. Obs.* 4. t. 1. f. 2.

Festuca avenacea spicis habitioribus, glumis glabris. *Raii Syn. ed.* 2. 261. *ed.* 3. 414.

F. graminea glumis glabris. *Scheuchz. Agrost.* 251. t. 5. f. 10.

Gramen avenaceum segetale majus, glumâ turgidiore. *Morif. sect.* 8. t. 7. f. 16.

Aegilops, sive *Festuca.* *Camer. Epit.* 927.

Angl. Smooth Rye Brome-grass.

In arvis.

Near Edwinsford, Caermarthenshire, among the winter corn. *Sir J. Banks.* In a meadow below Cook's Folly near Bristol. *Sir T. Frankland.* In Norfolk not rare. *Rev. Mr. Bryant.*

Annua. Fl. Julio.

Radix fibrosa. *Culmus* tripedalis, foliosus, erectus, simplex, teres, glaber. *Folia* linearia; supra, margineque præcipuè, pilosa; subtus scabra; vagina læviuscula. *Stipula* brevis, cerosa, pilosa. *Panicula* vix semipedalis, patens; ramis semiverticillatis, rectiusculis, angulatis, scabris, longitudine inæqualibus, plerumque simplicibus unifloris. *Spiculæ* ovatæ, fere ellipticæ, compresso-planæ, flosculis 9 ad 12, basi remotis, unde rachis flexuosa, glabra, tota in conspectum venit. *Calyx* inæqualis, muticus, glumis ellipticis, glabris, triner-vibus,

vibus, margine membranaceis. Flosculi teretiusculi, glabri, (rarius pubescentes), albido-virescentes, nitidi; gluma exterior elliptica, turgida, nec depressa, obsolete septemnervis, margine apicem versus parum membranacea, sub apice aristata; arista rectiuscula, scabra, glumâ brevior; interior longè angustior & tenuior, concava, mutica, margine pectinato-ciliata. Semen cylindricum, supra fuleo exaratum.

multiflorus. 2. **BROMUS** paniculâ patente; pedunculis simplicibus, spiculis ovato-lanceolatis compressis quindecimfloris, flosculis subimbricatis teretiusculis.

Bromus multiflorus. *Weigel. Obs. 2. t. 1. f. 1. Gmel. Syst. Nat. Linn. 188. Tr. of Lin. Soc. v. 4. 283.*

B. secalinus. *Leers. 36. t. 11. f. 2. Lightf. 1086. Linn. Fl. Suec. ed. 2. 96, var. 2.*

Festuca graminea, glumis hirsutis. Baub. Theatr. 143. Scheuchz. Agr. 250, t. 5. f. 9. Angl. Downy Rye Brome-grass.

In arvis rariùs. *Dill. in Raii Syn. 414.*

Between Edinburgh and New Haven. Observed by Miss Hancock near Norwich.

Annua. Fl. Julio.

Habitus præcedentis. Culmus glaber. Folia inferiora subtùs nuda, supra ad margines subpilosa; superiora subtùs brevi lanugine pubescentia, supra pilosa; omnia margine scabra, vagina glabra. Stipula brevissima, laccera, vix pilosa. Panicula ut in priore. Spiculæ unciales, ovato-lanceolatae, compresso-planæ, flosculis 12 ad 16, imbricatis, demum subremotis. Glumæ omnes plerumque molissimè pubescentes, rariùs glabræ. Affinis præcedenti, at distincta species videtur.

mollis. 3. **BROMUS** paniculâ erectâ coarctatâ; pedunculis ramosis, spiculis ovatis, flosculis imbricatis depressis nervosis pubescentibus.

Bromus

Bromus mollis. *Sp. Pl.* 112. *Tr. of Linn. Soc.* v. 4. 284. *Curt. Lond. fasc.* 1. t. 8. *Mart. Fl. Rust.* t. 99. *Relb.* 44. *Sibth.* 47. *Leers.* 37. t. 11. f. 1. *Weigel. Obs.* 7. t. 1. f. 4. *Schreb. Gram.* 60. t. 6. f. 1, 2.

B. polymorphus α . *Huds.* 48. var. 1. *With.* 159.

B. hordeaceus. *Sp. Pl. ed.* 1. 77.

B. Linn. Hort. Cliff. 25. ex *Herb. Cliff.*

B. n. 1504. *Hall. Hist.* v. 2. 236.

Festuca avenacea hirsuta, paniculis minus sparsis.
Raii Syn. 413.

Gramen avenaceum pratense, glumâ breviorē squamosâ & villosâ. *Moris. sect.* 8. t. 7. f. 18.

G. avenaceum, locustis villosis, angustis, candidantibus & aristatis. *Scheuchz. Agr.* 254. t. 5. f. 12.

β . *Culmo humillimo*, paniculâ pauciflorâ, e solo sterilissimo.

Bromus nanus. *Weigel. Obs.* 8. t. 1. f. 9. *Herb. Linn.*

Angl. Soft Brome-grass.

In muris, aggeribus et pascuis vulgaris.

Biennis. Fl. Junio.

Radix fibrosa, parùm ramosa, albida. Culmus erectus, bipedalis, simplex, teres, striatus, sæpiùs glaber, subindè pubescens, geniculis tumidis, quandoque hirsutis. Folia, ut et vaginæ, striata, pilis mollibus canescentibus villosa, margine vix scabra. Stipula acutiuscula, lacera. Panicula bi- vel triuncialis, erecta, coarctata, demum patentiuscula, decomposita; ramis semiverticillatis, simplicibus racemosisque, longitudine variis, angulatis, pubescentibus. Spiculæ erectiusculæ, ovatæ, acutæ, parùm compressæ, flosculis 5 ad 10, arctè imbricatis. Calyx e glumis duabus inæqualibus, ellipticis, mucronatis, margine scariosis, carinatis, villosis, 7—9 nervibus, nervis viridibus validis exstantibus.

Flosculi

Flosculi calyci conformes, depresso-concavi, aristis scabris longitudine valvularum; glumæ interiores tenuissimæ, longèque angustiores, enerves, margine incrassato, viridi, ciliato. Semen ellipticum, depresso-planum.

*racemo-
sus.*

4. BROMUS paniculâ erectiusculâ diffusa; pedunculis simplicibus, spiculis ovatis sexfloris, flosculis imbricatis depresso-nervosis glabris.

Bromus racemosus. Sp. Pl. 114. Tr. of Linn. Soc. v. 4. 286.

B. polymorphus β et δ . *Huds. 49. var. 3 et 4. With. 160.*

Festuca avenacea, spicis strigosioribus, è glumis glabris compactis. *Raii Syn. ed. 2. 261. ed. 3. 414.*

Gramen avenaccum pratense, gluma tenuiore glabra. *Dill. in Raii Syn. 414.*

G. avenaccum pratense, glumâ squamosâ longiore glabrâ. *Morif. sect. 8. t. 7. f. 19. Herb. Bobart.*

G. avenaceum, hirsutum, annuum, paniculâ ampliore; magisque sparsâ, locustis crassioribus, glabris et aristatis. *Till. Pis. 74. et*

G. loliaceum, alpinum, spicâ exili, rarioribus locustis. *Ponted. Comp. 46. Herb. Sherard. Ang. Smooth Brome-grass.*

In pratis et pascuis.

At Holkham, Norfolk. *Mr. Crowe. Northamptonshire. Mr. Dickson.*

Annua. Fl. Junio.

Præcedente toto habitu gracilior. Culmus glabriusculus. Folia utrinque pilosa, margine parùm scabra. Vaginæ pilis, plerumque deflexis, hirsutæ. Panicula rara, diffusa, nec coarctata, simplex, pedunculis semiverticillatis, scabris, elongatis, indivisis, vix unquam bifloris. Spiculæ præcedentis, at glaberrimæ, vix septemfloræ, nitidæ, viridi alboque vittatæ, nervis minùs prominentibus, aristis scabris longitudine valvularum.

E spe-

¶ specimine minùs luxuriante in Angliâ nato descripsit Linnæus.

†5. **BROMUS** paniculâ nutante ; pedunculis sim-*squarros-*
plicibus, spiculis ovatis duodecimfloris, floscu-*sus.*
lis imbricatis depresso, aristis divaricatis.

Bromus squarrosus. Sp. Pl. 112. Tr. of Linn. Soc. v. 4. 288. Hudf. 49. With. 160. Villars. Dauph. v. 2. 115.

Festuca graminea, glumis vacuis. Scheuchz. Agroft. 251. t. 5. f. 11.

Gramen phalaroides majus acerosum, nutante spicâ. *Barrel. Ic. t. 24. f. 1.*

Angl. Corn Brome-grass.

In arvis Angliæ australioris. *Hudf.*

Near Glastenbury, Somersetshire, and Marshfield, Suffex. *Hudf.*

Annua. Fl. Julio.

Radix fibrosa, parva, annua. Culmus pedalis, simplex, glaber, striatus, foliosus. Folia et vaginæ pubescunt, villis brevissimis deflexis. Panicula pauciflora, pedunculis secundis, simplicibus, compressis, apice incrassatis. Spiculæ omnium maximæ, ovatæ, tumidæ, nitidæ, flosculis 12 ad 16, arctè imbricatis, depresso-concavis, glabris, subinde scabriusculis. Calyx inæqualis, nervosus, muticus. Glumæ corollinæ exteriores calyce parùm majores, multinerves, apice bipartitæ, aristis subulatis, scabris, recurvato-divaricatis, longitudine glumarum ; valvulæ interiores longè minores, ciliis raris pectinatæ. Semen ellipticum, depresso-concavum, corollæ adnatum, apice villosum.

Descriptionem confeci ad exemplaria optima in herbario Linnæano conservata cum plantam ipsam in Angliâ nunquam inveni, nec in ullo plantarum thesauro, purè Britannicarum, adhuc vidi. De Hudsoni auctoritate unicè pendet ; et vereor ne ex errore quodam Bromum fecalinum pro hâc specie assumpserit.

arvensis. 6. *BROMUS* paniculâ patente ; pedunculis ramosis, spiculis lanceolatis octofloris, flosculis ellipticis imbricatis depresso glabriusculis.

Bromus arvensis. *Sp. Pl.* 113. *Fl. Suec.* n. 97. *Tra. of Linn. Soc.* v. 4. 289. *With.* 162. *Leers.* 38. t. 11. f. 3. *Villars. Dauph.* v. 2. 116.

B. erectus β . *Huds.* 50?

Festuca elatior, paniculis minùs sparsis, locustis oblongis strigosis aristatis purpureis splendentibus. *Raii Syn. ed.* 2. 261. *ed.* 3. 414. *Herb. Sherard.*

F. graminea, effusâ jubâ. *Scheuchz. Agr.* 262. t. 5. f. 15.

F. avenacea sterilis elatior. *Rel. Rudb.* 15. f. 2.

Aegilops major, caule et foliis arundinaceis, locustis glabrioribus et angustioribus c fusco xerampelinis. *Dill. Giff.* 130. *App.* 60. *Herb. Sher.*

Gramen bromoides serotinum annuum, paniculâ rariore magisque sparsâ, locustis angustioribus, purpurascentibus, glabris et aristatis. *Till. Pis.* 76. *Ibid.*

Angl. Field Brome-grass.

Inter segetes varius.

Near Southampton. *Sherard.* At Earsham, Norfolk. *Mr. Woodward.*

Annua. Fl. Julio.

Radix fibrosa, parva. Culmus tripedalis, erectus, simplex (basi quandoque ramoso-luxurians), teres, striatus, glaber, foliosus, geniculis quatuor vel quinque. Folia patentia, acuta, striata, utrinque pubescentia, margine parùm scabra, longius vaginata. Vaginæ cylindricæ, nervosæ, mollissimè pubescentes, pilis deflexis. Stipula brevissima, lacera. Panicula erectiuscula, multiflora, ramis patentibus, numerosis, simplicibus et ramosis, scabris. Spiculæ ovato-lanceolatæ, erectiusculæ, de-

mum nutantes, nitidæ; purpureo variæ, plerumque glabræ, subinde oculo armato pubescentes. Glumæ calycinæ valdè inæquales, lanceolatae, acutæ, carinatae, nervosæ, margine membranaceæ. Flosculi 7 ad 10, arctè imbricati, elliptici, depressio-concavi, carinâ scabri, margine membranacei, nervis duobus approximatim longitudinalibus lateralibus; aristis glumâ plerumque longioribus, parùm infra apicem membranaceum e carinâ elongatâ exortis, rectis, subulatis, scabris. Glumæ interiores minores, enerves, obtusæ, muticæ, margine pectinatae. Semen apice villosum.

7. *BROMUS* paniculâ erectâ; pedunculis sub-*erectus*, simplicibus, flosculis lanceolatis teretiusculis, foliis radicalibus angustissimis ciliato-pilosis.

Bromus erectus. *Huds.* 49. *ed.* 1. 39. *Tr. of Linn. Soc.* v. 4. 290. *With.* 160. *Sibth.* 47. *Relb. Suppl.* 2. 8. *Engl. Bot. t.* 471.

B. agrestis. *Allion. Ped.* v. 2. 249.

B. perennis. *Villars. Dauph.* v. 2. 122.

B. n. 1507. *Hall. Hist.* v. 2. 237. *Davall.*

Festuca avenacea sterilis spicis crectis. *Raii Syn. ed.* 2. 261. *ed.* 3. (exclusis synonymis) 413.

Gramen bromoides pratense, foliis præter culmum angustissimis, rarâ lanugine villosis. *Scheuchz. Agr.* 255. *t.* 5. *f.* 13. *Herb. Sher.* ab ipso Scheuchzero missum.

G. bromoides paniculatum, foliis et culmo villosis. *Scheuchz. Agr.* 257. ex Sherardi sententiâ.

G. quod Festuca pratensis lanuginosa C. B. *Vaill. Paris.* 93. *t.* 18. *f.* 2, synonymis pessimè confusis.

G. avenaceum glabrum, paniculâ purpureo-argentea splendente. *Morif.* v. 3. 213. *n.* 20. *Herb. Bobart.*

G. loliaceum, locustis longis aristatis. *Mont. Prod.* 35. f. 2. exclusis omnino synonymis Raii et Morisoni. *Herb. Sher.*

G. sparteum, longâ et spicatâ paniculâ, Lolii utriculis, *Festucæ* potius, majus. *Barr. Ic.* t. 13. f. 1.

Angl. Upright Perennial Brome-grass.

In pascuis arenosis, cretâ substratis.

Ditchley Park, Oxfordshire. *Mr. Woodward.*
Holkham, Norfolk. *Mr. Crowe.* Near Darent Wood, Dartford. *Rev. Dr. Goodenough.*

Perennis. Fl. Julio.

Radix fibrosa, cæspitosa, fibris villosis, fuscis. Culmus ferè tripedalis, erectus, strictus, simplex, teres, glaber, rariùs pubescens, ultra medium foliosus, geniculis quatuor circiter. Folia lineari-lanceolata, acutiuscula, striata, lætè viridia, scabra; radicalia angustissima, pilis longis albis sparsis adscendentibus ciliata, subinde involuta. Vaginæ striatæ, glabræ. Stipula brevissima, erosa. Panicula erecta, ramis erectis, scabris, numero et longitudine variis, simplicibus, rariùs bifloris. Spiculæ lineari-oblongæ, compressæ, erectæ, purpurascens, vel glabræ vel tomentosæ. Glumæ calycinæ lanceolatæ, acutæ, carinatæ; interiore majore trinervi. Flosculi 5 ad 9, imbricati, lanceolati, compresso-teretiusculi, subangulati, carinati, obsoletè nervosi, carinâ scabri; aristis longitudine vix glumarum, parùm infra apicem membranaceum exortis, rectis, scabris. Glumæ interiores apicem versus minutè pectinato-ciliatæ, ciliis adscendentibus, minùs quam in maximâ parte Bromorum conspicuis. Antheræ crocæ. Semen lineari-oblongum, supra sulcatum, apice villosum.

Primus invenit Sherardus et cum Bobarto communicavit. *Herb. Sher.*

Anne ad *Festucas* amandanda, ob valvulam interiorem vix pectinatam, radicemque perennem? Obstat verò affinitas ejus cum sequente.

8. *BROMUS* paniculâ nutante ramosâ, flosculis *asper*. lanceolatis teretiusculis subnervibus, foliis uniformibus: inferioribus hirsutis.

Bromus asper. *Linn. Suppl.* 111. *Tr. of Linn. Soc.* v. 4. 293. *With.* 161. *Mart. Fl. Rust.* t. 126. *Retz. Prod. ed.* 2. 25.

B. ramosus. *Linn. Syst. Veg. ed.* 13. 102, (nec *Mant.* 34.) *Huds. ed.* 1. 40. *Cullum.* 40.

B. nemoralis. *Huds.* 51.

B. nemorosus. *Villars. Dauph.* v. 2. 117.

B. hirsutus. *Curt. Lond. fasc.* 2. t. 8. *Relb.* 48. *Sibth.* 48.

B. montanus. *Pollich.* v. 1. 116. *Retz. Obs. fasc.* 2. 7.

Poa paniculâ nutante, laxâ, spicis oblongis, multifloris, sæpe aristatis. *Gmel. Sib.* v. 1. 110. t. 21. *Herb. Linn.*

Gramen avenaceum dumetorum, paniculâ sparsâ. *Raii Syn.* 415.

G. avenaceum dumetorum, jubâ longiore, spicâ divisâ. *Morif. sect.* 8. t. 7. f. 27.

Angl. Hairy Wood Brome-grass.

In nemorosis et sepibus humidiusculis.

Annua vel *biennis.* Fl. Julio.

Radix fibrosa, fibris crassiusculis, horizontaliter patentibus, lævibus. *Culmus* 4 vel 5 pedalis, erectus, teres, striatus, foliosus, supernè glaber, geniculis tribus circiter. *Folia* patentia, plana, acuta, nervosa, aspera atque pilosa, latitudine omnia ferè æqualia. *Vaginæ* teretes, vix carinatae, nervosae; inferiores pilis deflexis hirsutissimæ. *Stipula* brevis, lacera. *Panícula* pedalis, nutans, multiflora; ramis ramulisque plerumque binatis, elongatis, nutantibus, asperis. *Spiculæ* lineari-oblongæ, pendulæ, fusco-virides, subpubescentes. *Glumæ* calycinæ valdè inæquales, lanceolatae, carinatae, scabrae; majore trinervi. *Flosculi* novem circiter, imbricati, demum laxè patentés, lanceolati, teretiusculi,

carinati, obsoletè trinerves, carinâ præcipuè asperi, margine scariosi, sub apicem aristati, aristâ glumis breviorè ; glumæ interiores crebriùs ciliatæ, margine scarioso, inflexo. Semen fulco exaratum.

Sterilis. 9. BROMUS paniculâ nutante subsimplici, flosculis lanceolatis nervosis sulcatis, foliis pubescentibus.

Bromus sterilis. *Sp. Pl.* 113. *Tr. of Linn. Soc.* v. 4. 295. *Huds.* 50. *With.* 162. *Relb.* 46. *Sibth.* 47. *Curt. Lond. fasc.* 1. t. 9. *Mart. Fl. Ruß. t.* 125. *Leers.* 37. t. 11. f. 4. *Ger. em.* 76.

Festuca avenacea sterilis elatior, seu Bromos Dioscoridis. *Raii Syn.* 412.

F. graminea annua sterilis, spicis dependentibus.

Morif. sect. 8. t. 7. f. 11.

Angl. Barren Brome-grass.

In arvis, ruderatis et sepibus vulgaris.

Annua. Fl. Junio, Julio.

Radix fibrosa, parva, fibris capillaribus. Culmus sesquipedalis, erectus, gracilis, teres, striatus, ad apicem usque foliosus, geniculis quinque vel sex. Folia patentia, plana, subflaccida, angusta, acutiuscula, nervosa, utrinque mollissimè pubescentia, margine scabriuscula. Vaginæ cylindricæ, nervosæ, mollissimè villosæ, pilis parùm deflexis. Stipula oblonga, lacera. Panicula spithamæa, nutans ; ramis sæpiùs ternis, elongatis, nutantibus, asperis, simplicibus, vix unquam bifloris. Spiculæ lanceolatae, pendulæ, fusco-virentes, scabræ. Glumæ calycinæ maximè inæquales, lanceolatae, scabræ ; majori angulatâ, nervosâ. Flosculi 6 ad 8, imbricati, demum patentiusculi, lanceolati, compressi, carinati, asperi, 7 vel 9 nerves, nervis duobus submarginalibus maximis, inter nervos sulcati atque subpubescentes, apice ferè aristati, aristis glumâ longioribus, scabris ; glumæ interiores scitis raris

raris pectinatae, margine scarioso, inflexo. Stamina tria. Semen fulco exaratum.

10. *BROMUS* paniculâ erecto-patente subramo-*diandrus*.
sâ, floſculis lanceolatis nervosis fulcatis dian-
dris.

Bromus diandrus. *Curt. Lond. fasc. 6. t. 5. Tr.*
of Linn. Soc. v. 4. 296.

B. madritensis. *Sp. Pl. 114. With. 161.*

B. muralis. *Hudſ. 50. Sibth. 48.*

B. ciliatus. *Hudſ. ed. 1. 40.*

B. ſterilis, erectâ paniculâ major. *Barr. Ic.*
t. 76. f. 1.

Festuca avenacea ſterilis, paniculis confertis erec-
tioribus, aristis brevioribus. *Raii Syn. ed. 2.*
261. Pluk. Phyt. t. 299. f. 2. Herb. Sher.

F. avenacea ſterilis, pediculis brevioribus & ſpicis
erectis. *Moriſ. v. 3. 212. n. 13. et*

F. graminea annua ſpicis erectis. *Ejuſd. ſect. 8.*
t. 7. f. 13. Herb. Bobart.

Gramen bromoides pumilum, locuſtis erectis,
majoribus, ariſtatis. *Scheuchz. Agr. 260.*
Herb. Sher. a Scheuchzero ipſo.

Angl. Upright Annual Brome-graſs.

In arenosis et ad muros.

Common on the ſandy grounds in Jerſey. *Sherard*
in Herb. At the foot of St. Vincent's Rock,
Bristol, on the farther part near the meadows.
Sir J. Banks. Near Batterſea. Curtis.

Annua. Fl. Junio.

Radix fibroſa, parva. Culmi pedales, vel ſesquipedales,
erecti, ſtriſti, graciles, teretes, ſtriati, glabri, folioſi,
geniculis tribus plerumque. Folia præcedentis, at mi-
nùs pubeſcentia, ſæpè cunino glabra. Vaginæ ner-
voſæ, ſubcarinatae, ſæpiùs glabræ, quandoque piloſæ,
pilis deſlexis. Stipula brevis, eroſa. Panicula vix tri-

uncialis, erecta, fastigiata; ramis ternis, quaternis, quinifve, scabriusculis, plerumque simplicibus, brevibus, unifloris, solo fertiliori tantum elongatis et ramosis, semper tamen erectis vel erectiusculis, nunquam pendulis. Spiculæ lineari-lanceolatae, erectae, sæpius fuscescentes, scabrae. Flosculi præcedentis, sed diandri, minùsque argutè nervosi, nervis duobus lateralibus approximatis præcipuè conspicuis.

Optimè dignoscitur flosculis diandris, monente Curtisio, ut et Witheringio qui in Lusitaniâ invenit. Magnitudine multùm variat. Distinctus est a Bromo rubente Linnæi.

sylvaticus.

11. BROMUS spicâ simplici nutante secundâ, spiculis sessilibus teretiusculis, aristis glumâ longioribus, foliis pilosis.

Bromus sylvaticus. *Pollich. v. 1. 118. Pourret. Aët. Tolos. v. 3. 308. Tr. of Linn. Soc. v. 4. 300.*

B. pinnatus. *Fl. Dan. t. 164.*

B. gracilis. *Weigel. Obs. 15. t. 1. f. 11.*

Festuca sylvatica. *Huds. ed. 1. 38. With. 158.*

Relb. 44. Sibth. 46. Lightf. 103. Mart.

Fl. Rust. t. 114. Dickf. H. Sicc. fasc. 13. 9.

F. pinnata *β. Huds. 48.*

Triticum n. 1432. *Hall. Hist. v. 2. 213.*

Gramen avenaceum dumetorum spicatum. *Raii Syn. 394.*

Angl. Slender Wood Brome-grass.

In dumetis, nemorosis, et scipibus frequens.

Perennis. Fl. Julio.

Radix fibrosa, cæspitosa, fusca. Culmi bipedales, erecti, simplices, teretes, foliosi, glabri, apice attenuati et inclinati. Folia patentia, latiuscula, acuminata, rigidiuscula, lætè viridia, scabra, nervosa, subtùs præcipuè striata, magis vel minùs pilosa. Vaginæ strictæ, striatæ, pilosæ. Stipula brevis, obtusa, lacera. Spica simplex, nutans, serè triuncialis, rachi subflexuosâ, striatâ, scabrâ,

scabrâ, hinc canaliculato-compressâ. Spiculæ 6 vel 7, alternæ, sessiles, secundæ, non divaricatæ, lineari-lanceolatæ, luteolo-virides, plerumque pubescentes. Glumæ calycinæ inæquales, lanceolatæ, brevius aristatæ, nervosæ, apicem versus hirtæ. Flosculi 6 ad 9, imbricati, demum patentiusculi, subcylindrici, apicem versus præcipuè nervosi atque hirti. Arista terminalis, glumis plerumque longior, subulata, scabra. Gluma interior linearis, retusa, setis erectis rigidis ciliata.

Valvula interior pectinato-ciliata mihi certius Bromi indicium præbet quam aristæ insertio, quæ in pluribus speciebus difficillimè determinatur, & sæpe variabilis est.

12. BROMUS spicâ simplici erectâ distichâ, *sp. pinnatus*.
culis sessilibus teretiusculis, aristis glumâ brevioribus, foliis nudiusculis.

Bromus pinnatus. Sp. Pl. 115. Tr. of Linn.

Soc. v. 4. 301. Hudf. ed. 1. 41. Pollich.

v. 1. 117. Weigel. Obs. 14. t. 1. f. 10.

Festuca pinnata. Hudf. 48. Relb. 44. Sibth.

46. Dickf. H. Sicc. fasc. 13. 8.

Avena læta. Salisb. Prod. 22.

Triticum n. 1431. Hall. Hist. v. 2. 212.

Gramen spicâ Brizæ majus. Raii Syn. 392 Rel.

Rudb. 11. f. 2.

G. sparteum, spicatâ Brizæ paniculâ et corniculatâ. Barr. Ic. t. 25.

Angl. Spiked Heath Brome-grass.

In campis et ericetis, solo calcareo.

In Yorkshire, Oxfordshire and Kent not uncommon. On Burford Downs, Oxfordshire. *Rev. Dr. Goodenough.*

Perennis. Fl. Julio.

Radix fibrosa, subrepens. Culmus sesquipedalis, vel bipedalis, erectus, simplex, teres, foliosus, glaberrimus, apice rectus et rigens, parumque attenuatus. Folia erectiuscula, lanceolata, acuminata, rigidula, striata, nervosa,

nervosa, scabra, sæpiùs nuda, rariùs supra pilosa. Vaginæ strictæ, striatæ, glaberrimæ. Stipula brevis, obtusa, ciliata. Spica simplex, erecta, longitudine varia, rachi subflexuosâ, striatâ, angulis scabrâ, hinc complanatâ. Spiculæ 6 ad 10, alternæ, sessiles, distichæ nec secundæ, erectæ, lineari-lanceolatæ, pubescentes. Glumæ calycinæ parùm inæquales, lanceolatæ, vel subovatæ, subaristatæ, multinervosæ, margineque præcipuè hirtæ. Flosculi 6 ad 12 vel etiam plures, arctè imbricati, subcylindrici, apicem versus præcipuè nervosi atque hirti. Arista terminalis, glumis plerumque brevior, quandoque etiam brevissima, subulata, scabra. Gluma interior retusa, setis erectis ciliata ut in præcedente.

39. STIPA. *G. Pl.* 37. *Juss.* 30.

Cal. bivalvis, uniflorus. *Cor.* valvula exterior aristâ terminali, longissimâ, basi articulatâ.

pennata. † 1. STIPA aristis lanatis.

Stipa pennata. *Sp. Pl.* 115. *Huds.* 29. *With.* 163. *S. n.* 1514. *Hall. Hist.* v. 2. 239.

Gramen sparteum pennatum. *Raii Syn.* 393.

G. spicatum aristis pennatis. *Scheuchz. Agr.* 153. *t.* 3. *f.* 13. B.

Spartum austriacum. *Ger. em.* 42.

Avena capillacea austriaca, aristis longissimis pennatis. *Morif. sect.* 8. *t.* 7. *f.* 9. opt.

Angl. Feather-grass.

In rupibus calcareis subalpinis rarissimè.

On the lime-stone rocks hanging over a little valley called Long Sleedale, about 6 miles north of Kendal, Westmoreland; Dr. Richardson and T. Lawson. *Dill. in R. Syn.* Not now to be found.

Perennis. Fl. Julio.

Radix fibrosa, caespitosa. Culmi simplices, pedales, erecti, enodes, teretes, glaberrimi, foliis omnino vestiti. Folia involuto-setacea, mucronata, glauca, marginibus scabri.

scabriuscula. Vaginæ longissimæ atque maximè dilatæ, amplexicaules, striatæ, glabræ, intùs nitidæ. Stipula lanceolata, folio adnata. Panicula simplex, pauciflora, intra vaginam folii supremi latens. Flores maximi, a 4 ad 6. Glumæ calycinæ subæquales, lanceolatæ, scariosæ, concavæ, nervosæ, glabræ, in aristam membranaceam tenerrimam, valvulis longiorem, attenuatæ. Flosculus subulatus, teres, enervis, calyce brevior, basi setoso-sericeus. Arista e geniculo ad apicem glumæ exterioris, omnium longissima, ferè pedalis, nutante-incurva, acuta, angulata, basi torta, per totam ferè longitudinem pilis mollibus albis sericeis patentiusculis pulcherrimè vestita, demum unà cum corollâ persistente induratâ, semen maturum investiente, decidua, et gyratim per campos scopulosque, ad novam propagandam progeniem sparsa.

40. AVENA. *G. Pl.* 37. *Juss.* 32.

Cal. bivalvis, multiflorus. *Cor.* valvula exterior dorso aristata; aristâ contortâ.

1. AVENA paniculata, calycibus subtrifloris, flos *fatua*. culis basi pilosis omnibusque aristatis enervibus.

Avena fatua. *Sp. Pl.* 118. *Huds.* 52. *With.*

164. *Relb.* 49. *Sibth.* 49. *Mart. Fl. Rust.* t. 81. *Leers.* 42. t. 9. f. 4.

A. erratica pilosa, aristis intortis. *Morif. sect.* 8. t. 7. f. 5.

Ægilops quibusdam aristis recurvis, seu *Avena pilosa.* *Raii Syn.* 389.

Æ. bromoides. *Ger. em.* 77.

Angl. Wild Oat, or Haver.

In agris et inter segetes.

At Hafod, Cardiganshire. Common in Norfolk.

Near Weymouth. *Rev. Dr. Goodenough.*

Annua. Fl. Augusto.

Radix fibrosa. Culmus erectus, simplex, tripedalis, subfoliosus, striatus, glaberrimus. Folia linearia, patentia, nervosa, scabra. Vaginæ tenuiores, nervosæ, læves.

Stipula

Stipula obtusa, denticulata, lacera. *Panicula erecta*, ramosissima, patens, ramis ramulisque semiverticillatis, scabris; pedunculis alternis, capillaribus, scabris, apice incrassatis, nutantibus. *Spiculæ* obcuneatæ. *Glumæ* calycinæ æquales, lanceolatæ, acutæ, submembranaceæ, multinervosæ, glabræ, flosculis longiores. *Flosculi* plerumque tres, subremoti, sensim diminuti, tereiusculi, eearinati, serè enerves, læviusculi, basi fetis numerosis, nitidis, hispidi, dorso medium versus aristati; aristâ calyce duplo longiore, genieulatâ, scabrâ, arefactione sub genieulo tortâ. *Gluma* interior mutica, eoneava, margine minutè ac densissimè ciliata. *Semen* fericeum.

Avena nuda minimè inter stirpes indigenas numeranda est, nee a Raio ipso pro indigena traditur.

Avena strigosa *Schreb. Spicil.* 52, *With.* 166, inter *A. fativam* aliquando lecta, nequaquam indigena est.

pubescens.

2. *AVENA* paniculâ erectâ subsimplici, calycibus subtrifloris, receptaculis barbatis, foliis planis pubescentibus.

Avena pubescens. *Sp. Pl.* 1665. *Huds.* 52. *ed.* 1. 42. *With.* 165. *Relb.* 50. *Sibth.* 49. *Lightf.* 105. *Leers.* 43. *t.* 9. *f.* 2.

A. sesquitertia. *Linn. Mant.* 1. 34, excluso syn. *Scheuchzeri.*

A. pratensis. *Gouan. Hort.* 53?

A. n. 1498. *Hall. Hist.* v. 2. 234.

Gramen avenaceum 7, seu glabrum (potiùs hirsutum) panicula purpuro-argentea splendente. *Raii Syn.* 406. *t.* 21. *f.* 2. *ed.* 2. 252, *n.* 3. 262, *n.* 10. conf. etiam p. 345.

G. avenaceum, paniculâ purpuro-argenteâ, splendente. *Scheuchz. Agr.* 226. *t.* 4. *f.* 20.

Angl. Downy Oat-grass.

In pratis et pascuis cretaceis.

At Marham, Norfolk. *Mr. Woodward.* On Swaffham heath, and at Holkham and Carbrooke, Norfolk. *Mr. Crowe.*

Perennis. Fl. Junio.

Radix fibrosa, cæspitosa, subrepens. Culmus pedalis, bipedalis, subindè major, erectus, simplex, teretiusculus, glaber, striatus, foliosus. Folia patentia, brevia, latiuscula, obtusa, plana, utrinque, ut et vaginæ inferiores, molliter pubescentia. Stipula deltoidea, brevis. Panícula erecta, subspicata, parùm ramosa, subsecunda, pedunculis scabris. Spiculæ erectæ, sub anthesin patentes, obcuneatæ. Glumæ calycinæ valdè inæquales, carinâ scabræ, apice acutæ, membranaceæ, muticæ; interior duplo longior, spiculam æquans, trinervis. Flosculi tres, tertio sæpè abortiente, remotiusculi, clavato-cylindrici, nervosi, scabriusculi, diaphani, apice subindè laceri, dorso medium versus aristati; aristâ calyce duplo longiore, scabrâ, sub geniculo tortâ. Gluma interior minor et tenuior, mutica, margine scabra. Receptaculum commune supra flosculos elongatum, pilosum, pilis albis, fursùm longioribus.

3. **AVENA** spicâ erectâ, calycibus subquinqueflo. *pratensis.*
ris, receptaculis pilosis, foliis involutis ferrulatis nudis.

Avena pratensis. *Sp. Pl.* 119. *Huds.* 52. *With.* 165. *Relb.* 50. *Sibth.* 49. *Lightf.* 105. *Leers.* 43. *t.* 9. *f.* 1.

A. bromoides. *Sp. Pl.* 1666. *Gouan. Hort.* 52.

A. n. 1499. *Hall. Hist.* v. 2. 234.

Gramen avenaceum montanum, spicâ simplici, aristis recurvis. *Raii Syn.* 405. *t.* 21. *f.* 1. *ed.* 2. 252. *n.* 2. *et* 345.

G. avenaceum angustifolium alpinum, spicatâ paniculâ, ex purpureo viridi et argenteo variegatâ. *Scheuchz. Agr.* 230.

G. aven. alpinum glabrum angustifolium, locustis aristatis in spicam dispositis. *Scheuchz. Agr.* 228. *t.* 4. *f.* 21, 22.

Angl. Narrow-leaved Oat-grass.

In pratis et pascuis calcareis.

Perennis.

Perennis. Fl. Julio.

Radix fibrosa, cæspitosa. Culmi plures, pedales vel sesquipedales, erecti, stricti, simplices, geniculo solitario prope basin, supernè nudè, striati, scabriusculi. Folia radicalia linearia, acuta, rigida, incurva, involuta, utrinque nuda, margine ferrulato-scabra; caulinum lætius, nervosum, vaginâ longissimâ, nervosâ, glabrâ. Stipula lanceolata. Spica erecta, plerumque simplicissima, spiculis superioribus subsessilibus, inferioribus longius pedunculatis, hisce rariùs binis, alterâ ferè sessili. Spiculæ erectæ, obovatæ. Glumæ calycinæ subæquales, acutæ, trinerves, carinâ parùm scabræ, longitudine flosculi inferioris. Flosculi quatuor vel plures, subremoti, teretiusculi, scabriusculi, nervosi, apice membranacei laceri, dorso supra medium aristati; aristâ flosculo duplo longiore, purpurascente, apice albâ. Gluma interior minor, tenuissima, minutè ciliata. Receptaculum sub flosculis pilosum, pilis quam in præcedente brevioribus.

flaves-
cens. 4. AVENA paniculâ ramosissimâ laxâ, calycibus subtrifloris inæqualibus, receptaculis pilosis, foliis planis subpubescentibus.

Avena flaves cens. Sp. Pl. 118. Hudf. 53. With. 165. Relb. 51. Sibth. 49. Curt. Lond. fasc. 3. t. 5. Leers. 44. t. 10. f. 5. Schreb. Gram. 76. t. 9.

A. n. 1497. Hall. Hist. v. 2. 233.

Gramen avenaceum pratense elatius, paniculâ flavescente, locustis parvis. *Raii Syn. 407. Scheuchz. Agr. 223. t. 4. f. 18.*

G. aven. spicâ sparsâ flavescente, locustis parvis. Moris. sect. 8. t. 7. f. 42.

Angl. Yellow Oat-grass.

In pratis, pascuis, et ad vias.

Perennis. Fl. Junio, Julio.

Radix fibrosa, subrepens. Culmus erectus, basi arcuatus et subramosus, sesquipedalis, foliosus, teres, striatus, triarticulatus,

articulatus, sub geniculis sæpè villosus. Folia plana, acuta, striata, magis vel minus pubescentia, vaginis striatis, tumescentibus, subvillosis. Stipula setosa. Panicula subnutans, effusa, aureo nitens, ramosissima, multiflora; ramis semiverticillatis, ramulis capillaribus, scabris, subflexuosis, pedunculis alternis. Spiculæ parvæ, obovatæ, divaricato-patentes. Glumæ calycinae acutæ, carinatæ, dorso scabræ, alterâ duplo majore trinervi. Flosculi duo vel tres, subremoti, lanceolati, compressi, obsoletè nervosi, dorso ultra medium aristati; aristâ flosculo duplo longiori, scabrâ. Gluma interior angustior, vix brevior. Receptaculum pilosum, pilis erecto-adpressis.

41. LAGURUS. *G. Pl.* 37. *Juss.* 30. *Gærtn. t.* 1.

Cal. bivalvis, uniflorus, aristis villosis. *Cor.* gluma exterior aristis 2 terminalibus, tertiâ dorsali retortâ.

1. LAGURUS spicâ ovatâ.

ovatus.

Lagurus ovatus. Sp. Pl. 119. *With.* 166. *Dicks.*

H. Sicc. fasc. 7. 1. *Schreb. Gram.* 143. *t.* 19. *f.* 3. *Salisb. Prod.* 25.

Alopecuros spicâ rotundiore. Moris. sect. 8. *t.* 4. *f.* 1.

Gramen alopecurum molle, spicâ incanâ. Barrel. Ic. t. 116. *f.* 1, 2.

G. spicatum tomentosum, longissimis aristis donatum. Scheuchz. Agr. 58. *t.* 2. *f.* 4. *B. C.*

Angl. Hare's-tail-grass.

In campis arenosis australioribus.

Found in Guernsey by Mr. Gosselin. *Dickson.*

Annua. Fl. Junio.

Radix fibrosa, parva, glabra. Culmus solitarius, basi subindè ramosus, ascendens, pedalis vel sesquipedalis, geniculis quatuor vel quinque sulcatis donatus, foliolus, teres; supernè nudus, striatus, glaber. Folia lanceolata

lata (superiora ovata) acuta, nervosa, mollissimè pubescentia. Vaginæ foliis longiores, fursùm ventricosæ, nervosæ, tomentosæ. Stipula oblonga. Spica nutans, ovata, obtusiuscula, densissima, multiflora. Flores congesti, ferè sessiles. Glumæ calycinæ æquales, lineari-lanceolatae, acutæ, pilis longis, albis, densis, pectinatosiliatæ. Flosculus solitarius, sessilis, lanceolatus, basi carinâque setosus, aristis duabus terminalibus, rectis, longitudine glumæ, tertiâque dorsali, duplo longiore, geniculatâ, scabrâ. Gluma interior minor, apice bipartita.

42. ARUNDO. *G. Pl.* 38. *Juss.* 32.

Cal. bivalvis. *Flosculi* lanâ persistente cincti.

Phragmites. 1. ARUNDO calycibus quinquefloris, paniculâ laxâ.

Arundo Phragmites. *Sp. Pl.* 120. *Huds.* 53. *With.* 166. *Relb.* 51. *Sibth.* 50. *Engl. Bot.* t. 401. *Leers.* 45. t. 7. f. 1.

A. vallatoria. *Raii Syn.* 401. *Ger. em.* 36.

A. vulgaris palustris. *Morif. sect.* 8. t. 8. f. 1. *Angl.* Common Reed.

In fossis, stagnis, paludibus, et ad fluviorum margines vulgaris.

Perennis. Fl. Julio.

Radix repens, perennis. Culmi annui, erecti, simplices, sexpedales, teretes, geniculati, foliosi, læves, intùs nivi. Folia lanceolata, acuminata, patentia, striata, glabra, margine aspera, subtùs lævissima atque glaucescentia. Vaginæ cylindricæ, striatæ, glabræ. Stipula brevissima, crosa, utrinque pilosa. Panicula erectiuscula, diffusa, ramosissima, ramulis capillaribus, læviusculis. Spiculæ erectæ, ovatæ. Glumæ calycinæ valdè inæquales, lanceolatae, acutæ, carinatæ, læves, majori trinervi. Flosculi a quatuor ad sex, glabri, lanceolati, acuti, mutici, nervosi, basi lanâ sericeâ longitudine glumarum, e receptaculo ortâ, cincti. Gluma interior duplo

duplo brevior, ciliata. Semen corollâ induratâ tectum, lanâque ad basin, demùm clongatâ, evolat.
Variat foliis variegatis. *D. Relban.*

2. ARUNDO calycibus unifloris corollâ longi-*epigejos.*
oribus, paniculâ erectâ, floribus imbricato-
congestis secundis, foliis lanceolatis.

Arundo epigejos. *Sp. Pl.* 120. *Engl. Bot. t.* 402.

A. Calamagrostis. *Huds.* 54. *Relb.* 52. *Lightf.* 106.

Calamagrostis lanceolata. *With.* 122.

Gramen arundinaceum, paniculâ molli spadiceâ,
majus. *Raii Syn.* 401. *Scheuchz. Prod.* 21. *t.* 5.

G. paniculatum, palustre, præaltum, exile, pani-
culâ arundinaceâ. *Ponted. Comp.* 56. *Herb.*
Sherard.

Angl. Wood Reed.

In fossis umbrosis, et nemorosis humidis.

Between Newington and Hornsey. *Hudson.* Near
Kennington Common. *Mr. Sowerby.* In Hethel
and Arminghall woods, Norfolk. *Mr. Crowe.*

Perennis. Fl. Julio.

Radix repens. Culmus altitudine ferè præcedentis, sed
parùm tenuior, basi sæpè ramosus, foliosus, glaber.
Folia lanceolata, acuminata, nervosa, subtùs præcipuè
glauca margineque aspera. Vaginæ strictæ, striatæ,
glabræ. Stipulâ lanceolata, multifida, utrinque nuda.
Panicula erecta, scabra, ramis patentibus. Flores con-
gesti, secundi, nutantes. Glumæ calycinæ subæquales,
lanceolatæ, acutæ, nervosæ, carinâ scabræ. Flosculus
solitarius, calyce multò brevior, albus, membranaceus,
lanâ petalis longiore cinctus, apice sæpiùs lacer, dorso
basin versus aristatus, aristâ geniculatâ, longitudine ferè
lanæ.

Icon in *Engl. Bot. t.* 402, ex errore caret lanâ ad basin
flosculi, et aristâ dorsali.

Arundinis genus maximè naturale nunquam ob numerum
flosculorum, in diversis speciebus diversum, dilaccran-

dum; neque tolerandum est nomen *Calamagrostis*, e Calamo et Agrostide conflatum. Conf. Linn. *Phil. Bot. sect.* 224, 225.

Calama- 3. ARUNDO calycibus unifloris corollâ longiori-
grostis. bus, paniculâ erectâ diffusâ, floribus sparsis erectis, foliis linearibus.

Arundo Calamagrostis. *Sp. Pl.* 121. *Fl. Dan.* t. 180. *Engl. Bot.* t. 403.

A. epigejos. *Huds.* 54. *Relb.* 51.

Calamagrostis epigejos. *With.* 123.

C. minor, glumis ruffis et viridibus. *Dill. in Raii Syn.* 401.

Gramen arundinaecum, paniculatum, montanum, paniculâ spadiceo-viridi, semine papposo. *Scheuchz. Agr.* 124.

Angl. Small Reed.

In nemorosis et sepibus pratisque humidis.

Near Oundle, Leicestershire; Mr. Seampton. *Dillenius.* At Ranaugh, Norfolk. *Mr. Humphrey.* Earsham-wood. *Mr. Woodward.* East Fen near Revesby Abbey, Lincolnshire. *Sir Jos. Banks.* Cambridgeshire. *Rev. Mr. Hemsted.* Near Prickwillow, Isle of Ely. *Rev. Dr. Goodenough.*

Perennis. Fl. Julio.

Radix fibrosa, vix repens. Culmus erectus, tri- vel quadri-pedalis, teres, glaberrimus, foliosus, præcedentibus multò gracilior, quandoque ramosus. Folia linearia, acuta, angusta, subinvoluta, lætè viridia, subtus scabriuscula, suprâ rariùs pilosa. Vaginæ longæ, strictæ, striatæ, ferè glabræ. Stipula lanceolata, sæpè lacera, utrinque glabra, decurrens. Panicula erectiuscula, diffusa, ramosissima. Flores sparsi, erecti. Glumæ calycinæ, spadiceæ vel purpureæ, subæquales, lanceolatæ, acutæ, carinatæ, dorso scabriusculæ, vix nervosæ. Flosculus solitarius, calyce multò brevior, albus, apice lacer, lanâ petalis longiore cinctus, aristâ exiguâ inter lacinias ad apicem petali majoris.

4. ARUN-

4. ARUNDO calycibus unifloris corollam æquantibus, paniculâ erectâ glomeratâ, floribus secundis muticis, foliis planis.

Arundo colorata. *Soland. in Ait. Hort. Kew.* v. 1. 116.

A. n. 1524. *Hall. Hist.* v. 2. 243.

Phalaris arundinacea. *Sp. Pl.* 80. *Huds.* 23.

Relb. 22. *Fl. Dan. t.* 259. *Leers.* 18. t. 7. f. 3.

Calamagrostis colorata. *Sibth.* 37.

C. variegata. *With.* 124.

Gramen arundinaceum acerosâ glumâ nostras.
Raii Syn. 400.

- β. G. arundinaceum acerosâ glumâ Jerscianum.
Raii Syn. 400.

- γ. Phalaris arundinacea β picta. *Sp. Pl.* 80.

Gramen paniculâ torosâ, folio variegato, striatum. *Morif. sect.* 8. t. 6. f. 43.

Angl. Canary Reed-grass.

In stagnis et ad fluviorum ripas.

- β. On the sea coast of Jersey, and near Llanberis.
Raii Syn.

- γ. Common in gardens. Wild near Cambridge.
Rev. Mr. Relban.

Perennis. Fl. Julio.

Radix repens, cæspitosa. Culmus bi- vel quinque-pedalis, erectus, teres, foliosus, striatus, glaber, geniculis pluribus, lævibus. Folia patentia, lanceolata, acutiuscula, striata, utrinque margineque lævia, in β glauca, in γ variegata. Vaginæ nervosæ, vix inflatæ, glabræ. Stipula brevis, obtusiuscula. Panicula erecta, ramosa, lobata, ramulis angulatis, scabris. Flores glomerati, secundi, albo purpureoque variegati. Glumæ calycinæ æquales, compressæ, carinatæ, trinerves, acutiusculæ, carinâ scabræ. Flosculus solitarius, longitudine calycis, lanceolatus, subcompressus, nitidus, muticus, basi penicillis duobus lanæ flosculo longè brevioribus. Glumæ extus pilosæ, longitudine æquales; exterior latior, alteram amplectens.

arenaria. 5. ARUNDO calycibus unifloris corollâ longioribus, paniculâ spicatâ, floribus erectis muticis, foliis involutis pungentibus.

Arundo arenaria. *Sp. Pl.* 121. *Huds.* 54. *Mart.*

Fl. Rusl. t. 32. *Dicks. H. Sicc. fasc.* 12. 5.

Fl. Dan. t. 917.

Calamagrostis arenaria. *Witb.* 123.

Gramen sparteum spicatum, foliis mucronatis longioribus, vel spicâ fecalinâ. *Raii Syn.* 393.

G. sparteum juncifolium non aristatum, spicâ fecalinâ. *Morif. sect.* 8. *t.* 4. *f.* 16.

Spartum anglicanum. *Ger. em.* 42.

Angl. Sea Reed. *Marram.* Sea Mat-weed.

In arenosis maritimis frequens.

Perennis. Fl. Julio.

Radix repens, articulata, longè latèque extensa. Culmus adscendens, tripedalis, rigens, teres, glaber, geniculatus, foliosus. Folia erecto-patentia, stricta, rigida, involuta, mucronato-pungentia, glauca, glabra; suprà fulcata, subtùs lævissima. Vaginæ nervosæ, glabræ. Stipula elongata, acuta, bipartita. Panicula erecta, spiciformis, ramis brevibus, erecto-adpressis. Flores erecti, glaucescentes. Glumæ calycinæ subæquales, lanceolatae, acutæ, compressæ, carinatae, obsolete trinerves, carinâ scabræ. Flosculus solitarius, calyce parùm brevior. Glumæ lanceolatae, inæquales, nervosæ, carinâ scabræ; exterior latior, apice erosa, alteram amplectens. Lana flosculo triplò brevior.

Gramen præcedenti, et inde Phalaridi affine; ad arcuam mobilem firmandam nec cominodius nec aptius quicquam.

43. LOLIUM. *G. Pl.* 38. *Juss.* 31.

Cal. monophyllus, fixus, multiflorus. *Flosculi* distichi.

perenne. 1. LOLIUM, spicâ muticâ, spiculis calyce longioribus, flosculis lanceolatis.

Lolium

Lolium perenne. *Sp. Pl.* 122. *Huds.* 55.
With. 167. *Relb.* 52. *Sibth.* 50. *Engl.*
Bot. t. 315. *Mart. Fl. Rust. t.* 4. *Dicks.*
Hort. Sicc. fasc. 11. 4. *Leers.* 47. *t.* 12.
f. 1.

L. rubrum. *Ger. em.* 78.

Gramen loliaceum, angustiore folio et spicâ.
Raii Syn. 395.

G. loliaceum, spicâ simplici, vulgare. *Morif.*
sect. 8. *t.* 2. *f.* 2.

β. *Lolium tenue.* *Sp. Pl.* 122, spiculis paucifloris.

γ. *Gramen loliaceum paniculatum.* *Dill. in Raii*
Syn. 395. *inmultiplici spicatâ paniculâ.* *Morif.*
loc. cit.

Leers. t. 12. *f.* 1. +.

δ. spicis brevioribus compressis. *Morif. loc. cit.*

G. loliaceum, spicâ latâ, e plurimis spicis duplici versu dense dispositis constante. *Scheuchz.*
Prod. 16. *t.* 2.

Sibth. 50. *With.* 167.

Angl. Perennial Darnel. Red Darnel. Ray-
grafs.

In pratis, pascuis, et ad vias ubique.

Perennis. *Fl. Junio.*

Radix fibrosa, pubescens. Culmus pedalis, erectus, basi geniculatus et flexuosus, foliosus, superne teres, striatus, glaber. Folia linearia, carinata, glabra, saturatè viridia. Vaginæ striatæ, glabræ. Stipula brevis, obtusa, amplexicaulis. Spica erectiuscula, disticha, compressa, rachi flexuosâ, glabrâ, striatâ, alternatim scrobiculatâ ad spiculas recipiendas. Spiculæ alternæ, erectæ, ovatæ, compressæ, multifloræ. Calyx univalvis, lanceolatus, concaviusculus, acutus, muticus, striatus, vix carinatus, glaber, spiculâ duplo ferè brevior. Flosculi distichi, approximati, alterni, lanceolati, depressi, nervosi, mutici, glumâ interiore minore, ciliatâ, concavâ.

vâ. Spiculæ quandoque aristatæ, rariùs viviparæ sunt. β e solo sterili gaudet foliis et spicis tenuioribus, spiculis tri- vel quadri-floris, minùs compressis, calyce tamen longioribus.

γ dignoscitur spicâ basin versus compositâ.

δ spicam habet apice condensatam, e spiculis approximatis ovatam. *Herb. D. Rose.*

temulentum.

2. **LOLIUM** spicâ aristatâ, spiculis calyce brevioribus, flosculis ellipticis, culmo supernè scabro. *Lolium temulentum. Sp. Pl. 122. With. 168. Relb. 53. Leers. 48. t. 12. f. 2. Fl. Dan. t. 160.*

L. album. Ger. em. 78. Raii Syn. 395.

Zizania. Tillands. Ic. 112.

Angl. Bearded Darnel.

In arvis, inter *Hordeum*, *Triticum*, aut *Linum*.

Annua. Fl. Julio.

Radix fibrosa, tomentosa, parva. Culmus plerumque subsolitarius, erectus, bipedalis, firmus, teres, trinodis, foliosus, striatus; infernè glaberrimus, nitidus; supernè scaber. Folia lanceolata, patentia, nervosa, scabra. Vaginæ striatæ, scabriusculæ. Stipula brevissima, obtusa, crenulata. Spica erecta, figurâ prioris, sed major, rachis scabrâ. Spiculæ oblongæ, minùs compressæ. Gluma calycina lineari-lanceolata, acutiuscula, mutica, complanata, striata, scabra, spiculâ plerumque longior; valvula quandoque altera, interna, minuta, elliptica, adest in spiculis inferioribus. Flosculi plurimi, distichi, approximati, alterni, elliptici, ventricosi, vix depressi, obsoletè nervosi, scabri, parùm infra apicem aristati, aristâ flosculo duplo longiore, subulatâ, albidâ, scabrâ, rectiusculâ. Gluma interior mutica, concava, fubeiliata. Semen depresso-planum.

arvense. 3. **LOLIUM** spicâ sub-muticâ, spiculis longitudine calycis, flosculis ellipticis, culmo glaberrimo.

Lolium

Lolium arvense. *With.* 168.

L. temulentum. *Huds.* 55. ex caractere.

L. verum Gefneri. *Morif. sect.* 8. *t.* 2. *f.* 1.

Angl. White Darnel. Annual Beardless Darnel.

In arvis rariùs.

In fields to the north of Forfar, Scotland. *Mr.*
Mackay.

Annua. Fl. Julio.

Habitus præcedentis, sed minor et tota glabra; folia supra tantum quandoque scabra sunt. Culmus, ut et rachis, omnino lævis. Calyx plerumque mihi videtur univalvis, at Cl. Withering (qui hanc speciem primus distinxit) bivalvem descripsit. Spicula calyci æqualis, vel parùm longior. Flosculi præcedentis; primo intuitu mutici, reverà accuratiùs inspecti, sub apicem brevissimè aristati sæpe inveniuntur.

Cum præcedente confudit Linnæus.

44. ROTBOLLIA. *Suppl.* 13. *Juss.* 31.

Cal. fixus, subuniflorus, simplex five bipartitus.

Flores alterni in rachi articulatâ.

1. ROTBOLLIA spicâ tereti subulatâ, glumâ *ca-incurvata*,
lycinâ subulatâ adpressâ bipartitâ. *ta.*

Rotbollia incurvata. *Linn. Suppl.* 114. *With.*

169. *Relb. Suppl.* 2. 8. *Flo. Dan. t.* 938.

Aegilops incurvata. *Sp. Pl.* 1490. *Huds.* 441.

Gramen parvum marinum spicâ loliaceâ. *Raii*

Syn. 395.

G. loliaceum maritimum, spicis articulatis. *Morif. sect.* 8. *t.* 2. *f.* 8.

Angl. Sea Hard-grass.

In pratis et pascuis maritimis.

At Sheringham and Yarmouth, Norfolk. *Mr.*

Crowe. Below King's Weston near Bristol. *Dr.*

Stokes. Near Seaton, Durham. *Mr. Robson.*

At Exmouth, Devonshire. *Rev. Dr. Goodenough.*

Annua. Fl. Augusto.

Radix fibrosa, ramosissima, elongata. Culmi numerosi, basi decumbentes, ramosi, geniculati, teretes, glaberrimi, foliosi. Folia patula, linearia, acuta, plana, striata, supra margineque scabra. Vaginæ foliis breviores, subtumidæ, striatæ, glabræ. Stipula brevis, obtusa. Spicæ terminales, solitariae, filiformes, subaeuatae, teretes, glabræ, multifloræ. Rachis articulata, alternatim scrobiculata, articulis unifloris. Calyx e glumis duabus, lanceolatis, acutis, muticis, glabris, striatis, parallelis, omne scrobiculum arctè cludentibus, sub anthesi vero patentibus. Flosculus solitarius, hermaphroditicus, calyce parùm brevior, glumis membranaceis, albis, muticis, subæqualibus.

45. ELYMUS. *G. Pl.* 39. *Juss.* 31.

Cal. lateralis, bivalvis, aggregatus, multiflorus.

arenari- 1. ELYMUS spicâ erectâ arctâ, calycibus lanceo-
us. latis longitudine spiculæ, foliis mucronato-
pungentibus.

Elymus arenarius. *Sp. Pl.* 122. *Huds.* 56.

With. 170. *Mart. Fl. Rust. t.* 31.

Triticum radice perenni, spiculis binis lanuginosis. *Gmel. Sib. v.* 1. 119. *t.* 25.

Spartum herba 4 Batavicum. *Clus. Hist.* 2. 221.

Angl. Upright Sea Lyme-grass.

In arenosis maritimis.

Isle of Bute, and in Devonshire. *Huds.* At Wells and Happisburgh, Norfolk. *Mr. Crowe.* Scotland. *Rev Dr. Walker.* Sea coast, four miles east of St. Andrews. *Mr. Mackay.*

Perennis. Fl. Julio.

Radix repens, latè extensa. Culmus tri- vel quadri-pedalis, erectus, firmus, teres, parùm striatus, glaberrimus, geniculis

niculis subtribus, basin versus præcipuè foliosus. Folia erecta, culmo laud multum breviora, lincaria, rigida, stricta, glauca, mucronato-pungentia, canaliculata, involuta, ccarinata; subtus nervosa, glaberrima; supra sulcata, sulcis albidis, oculoque armato scabris. Vaginæ striatæ, glaberrimæ. Stipula brevissima, crenulata. Spica erecta, spithamæa, subcompressa, arctè imbricata, multiflora, glauca, rachi tectâ angulatâ, angulis pubescentibus. Spiculæ sessiles, erecto-adpressæ, duplici serie per paria alternatim digestæ, spicæ medium versus ternatæ. Glumæ calycinæ sub singulâ spiculâ binæ, adpressæ, lanccolatæ, subinde ovatæ, mucronatæ, longitudine ferè æquales, carinatæ, extûs plerumque subpubescentes. Spiculæ lanccolatæ, subtrifloræ, longitudine calycis vel parûm longiores. Flosculi mucronati, apicc nervosi, vix carinati, pubescentes, glumâ interiore concavâ, parûm breviora, emarginatâ, minutè ciliatâ. Receptaculum commune flosculorum articulatum, articulis clavatis, pilosis.

Habitus Arundinis arenariæ, at etiam non florens dignoscitur stipulâ exiguâ, nec elongatâ bipartitâ.

2. ELYMUS spicâ inflexo-pendulâ laxâ, calycibus setaceis patulis spiculâ longioribus, foliis mucronato pungentibus. *geniculatus.*

Elymus. *Amæn. Acad.* v. 7. 192. t. 2. f. 5.

With. v. 1. 134. t. 2. f. 26.

E. geniculatus. *Curt. Obs. on Brit. Grasses*, 46.

Angl. Pendulous Sea Lyme-grass.

In littoribus maritimis.

Near Gravesend. *Mr. Dickson.*

Perennis. Fl. Julio.

Habitus præcedentis, quocum a Linnæo confundi videtur; altior tamen et gracilior est, foliis angustioribus, et dum floret facillimè distinguendus. Spica sesquipedalis, vel bipedalis, laxa, floribus distantibus unde rachis in conspectum venit, inferioribus etiam remotissimis. Ad primi vel secundi florum fasciculi inscriptionem rachis, primò erecta, angulo acuto demum inflectitur,

flectitur, et spica dein ad perpendiculum dependet. Glumæ calycinæ spiculis plerumque sesquialongiores, linearisetaceæ, glabræ, patentés, æquales. Spiculæ quadrifloræ. Flosculi remotiusculi, mucronati, nervosi, pubescentes. Receptaculum commune basi quandoque denudatum et elongatum.

europæus. 3. ELYMUS spicâ erectâ, spiculis bifloris aristatis, involucro exstriato.

Elymus europæus. *Linn. Mant.* 35. *With.* 170. *Sibth.* 51.

Hordeum sylvaticum. *Huds.* 57. *Mart. Fl. Rust.* t. 45.

Triticum sylvaticum. *Salisb. Prod.* 27.

Secale villosum. *Huds. ed.* 1. 46.

Gramen fecalinum majus sylvaticum. *Raii Syn.* 392.

G. hordeaceum montanum five majus. *Scheuchz. Prod.* 14. t. 1. f. 1.

Angl. Wood Lyme-grass.

In sylvis solo calcarco.

In Oxfordshire, Berkshire, Derbyshire, and the north of England.

Perennis. Fl. Junio.

Radix fibrosa, subcæspitosa. Culmus bipedalis, erectus, simplex, teres, striatus, glaber, basi præcipuè foliosus. Folia patentia, lanceolata, acuminata, plana, nervosa, utrinque scabra. Vaginæ nervosæ, parùm ventricosæ, magis vel minùs pilosæ, pilis deflexis. Stipula exigua, crenulata. Spica erecta, vel parùm nutans, stricta, arcta, biuncialis, obtusa. Rachis flexuosa, angulata, sulcata, scabra. Spiculæ erecto-adpressæ, triplici serie per paria digestæ. Glumæ calycinæ lanceolato-setaceæ, scabræ, obsoletè trinerves nec striatæ, aristatæ, aristis scabris, rectis, glumâ longioribus. Spiculæ subpedicellatæ, bifloræ, sæpe unifloræ, longitudine calycis. Flosculi omnes ferè perfecti, lanceolati, aristati, apice nervosi, scabri, aristis flosculo quandoque duplò longioribus.

46. HOR-

46. HORDEUM. *G. Pl.* 39. *Juss.* 32. *Gærtn. t.* 81.

Cal. lateralis, bivalvis, ternus, uniflorus.

1. HORDEUM flosculis lateralibus masculis arif- *muri-*
tatis, glumis calycinis intermediis lanceolatis *num.*
ciliatis.

Hordeum murinum. Sp. Pl. 126. *Huds.* 56.

With. 171. *Relb.* 53. *Sibth.* 51. *Curt. Lond.*

fasc. 5. *t.* 9. *Mart. Fl. Rust. t.* 43. *Dicks. H.*

Sicc. fasc. 13. 10. *Fl. Dan. t.* 629.

H. spurium. Ger. em. 73.

Gramen fecalinum et Secale sylvestre. Raii Syn.
391.

G. fecalinum vulgatissimum viarum. Moris.
sect. 8. *t.* 6. *f.* 4.

Angl. Wall Barley. *Moufe Barley.* *Way Bennet.*

In ruderatis et ad vias vulgaris.

Annua. Fl. totâ æstate.

Radix fibrosa, albida. Culmi plures, pedales, erectiusculi, basi decumbentes, patuli, infracti, geniculis tumidis, foliosi, teretes, læves. Folia patentia, laxiuscula, lineari-lanceolata, acuta, striata, scabra, lætè viridia. Vaginæ longitudine ferè foliorum, subventricosæ, striatæ, glaberrimæ. Stipula brevissima. Spica erecta, bi- vel tri-uncialis, arcta, multiflora. Rachis articulata, fragilis, compressa, angulis scabra. Spiculæ distichæ, imbricatæ. Glumæ calycinæ exteriores setacæ; interiores lineari-lanceolatæ, trinerves, pectinato-ciliatæ; omnes aristatæ, aristis rectis, scaberrimis, glumâ duplo longioribus. Flosculi semper solitarii, lanceolati, lævissimi, aristati, aristis validioribus et longioribus quam in calyce; centralis sessilis, hermaphroditicus; laterales pedicellati, minores, masculi. Glumæ interiores obtusè acuminatæ, muticæ, cum setâ ad basin, longitudinaliter fulcatæ; in floribus hermaphroditicis margine læves, masculis ciliatæ.

2. HOR-

pratense. 2. **HORDEUM** flosculis lateralibus masculis brevius aristatis, glumis calycinis omnibus setaceis scabris.

Hordeum pratense. *Huds.* 56. *With.* 171. *Relb.* 54. *Sibth.* 51. *Engl. Bot.* t. 409. *Mart. Fl. Rust.* t. 108.

H. nodosum. *Sp. Pl.* 126, excluso syn. *Raii.*

H. murinum β . *Ibid.*

Gramen fecalinum. *Raii Syn.* 392. *Ger. em.* 29.*
n. 4.

G. fecalinum minus pratense. *Morif. sect.* 8. t. 2.
f. 6.

G. spicatum fecalinum minus. *Scheuchz. Agr.* 17.
Vaill. Paris. t. 17. f. 6.

Angl. Meadow Barley.

In pratis et pascuis, præcipuè humidiusculis.

Perennis. Fl. Junio.

Radix fibrosa, in locis sterilioribus rariùs inundatis bulbosa evadit. Culmus sesquipedalis, erectus, gracilis, foliosus, supernè nudus, teres, tenuissimè striatus, glaber. Folia patentia, linearia, acuta, scabriuscula. Vaginæ parùm ventricosæ, striatæ, glabræ. Stipula exigua. Spica ut in præcedente, sed angustior, brevius aristata, rachi minùs fragili. Glumæ calycinæ omnes æquales, angustissimæ, setaceæ, scabræ, aristatæ. Flosculi ferè ut in priore, laterales verò longiùs pedicellati, multòque brevius aristati, plerumque masculi, subinde abortivi.

Dignoscitur primâ facie culmo altiore, stricto, supernè nudo, spicâ strigiosiori.

maritimum. 3. **HORDEUM** flosculis lateralibus masculis brevius aristatis, glumâ calycinâ internâ flosculorum lateraliuin semi-ovato.

Hordeum maritimum. *With.* 172. *Relb.* 54. *Mart. Fl. Rust.* t. 44. *Fl. Dan.* t. 630. *Pourret. Act. Tolos.* v. 3. 320.

H. ma-

H. marinum. *Hudf.* 57. *Dickf. H. Sicc. fasc.* 5. 4.

Gramen fecalinum palustre et maritimum. *Raii*

Syn. 392.

G. fecalinum maritimum minus. *Morif. sect.* 8.

t. 6. *f.* 5.

Angl. Sea Barley. Squirrel-tail-grass.

In pascuis et arenosis maritimis.

Annua. Fl. Junio, Julio.

Habitus Hordei murini, quocum a Curtifio confunditur.

Dignoscitur foliis lævioribus, glaucescentibus, spiculis subpyramidatis, compactioribus. Aristæ calycinæ et floris centralis longitudine æquales sunt, lateralium verò florum multò breviores. Character essentialis constat e figurâ dilatatâ, semi-ovatâ, glumarum calycinarum internarum flosculorum lateralium, dum reliquæ omnes e basi setaceæ sint.

47. TRITICUM. *G. Pl.* 40. *Juss.* 32. *Gærtn. t.* 81.

Cal. bivalvis, solitarius, multiflorus, in rachi flexuosâ, dentatâ.

1. TRITICUM calycibus truncatis quinquefloris, *juncceum*. foliis involutis mucronato-pungentibus.

Triticum juncceum. *Sp. Pl.* 128. *Hudf.* 58.

With. 173. *Relb.* 55. *Dickf. H. Sicc. fasc.* 6. 9.

Fl. Dan. t. 916.

Gramen maritimum, spicâ loliaceâ, foliis pungentibus, nostras. *Pluk. Phyt. tab.* 33. *f.* 4.

Dill. in Raii Syn. 391.

β. *G. loliaceum maritimum supinum*, spicâ crassiflore. *Dill. in Raii Syn.* 391.

γ. *G. caninum maritimum*, spicâ foliaceâ. *Raii Syn.* 391.

Angl. Sea Rushy Wheat-grass.

In arenosis maritimis frequens.

Perennis. Fl. Julio,

Radix

Radix repens, teres, articulata, glabra, fibris tomentosis. Culmus adscendens, basi sæpè purpurascens, teres, nudus, lævissimus, medium versus foliosus, supernè nudus, striatus. Folia erecto-patentia, linearia, acuminato-pungentia, involuta, striata, glauca; subtus lævissima; suprà inter strias minutè scabra, ut in *Elymo arenario*. Vaginæ strictæ, glabræ. Stipula brevissima. Spica erecta, stricta, glauca, palmaris. Rachis articulata, alternatim hinc complanata, extus teres, striata, glaberrima. Spiculæ solitariae, erectiusculæ, distichæ, remotiusculæ, mutiæ, glaberrimæ. Glumæ calycinæ eymbiformes, sulcatæ, obtusæ. Flosculi 5 vel 6, longitudine et figurâ calycis, subcarinati, apice emarginati vel denticulati, glumis interioribus parùm minoribus, ciliatis.

repens.

2. **TRITICUM** calycibus subulatis, multinervibus, quinquefloris, flosculis acuminatis, foliis planis, radice repente.

Triticum repens. *Sp. Pl.* 128. *Huds.* 57. *Willd.* 173. *Rehb.* 55. *Sibth.* 52. *Leers.* 45. t. 12. f. 3.

Gramen spicâ triticeâ repens vulgare, caninum dictum. *Raii Syn.* 390.

G. caninum vulgatius. *Morif. sect.* 8. t. 1. f. 8.

- β. *Triticum repens.* *Mart. Fl. Rusl.* t. 124. *Fl. Dan.* t. 748. *Willd.* 173. var. 4.

- γ. Gramen caninum maritimum, spicâ triticeâ nostras. *Raii Syn.* 390.

Triticum repens var. 5. *Willd.* 173.

Angl. Creeping Wheat-grass. Couch-grass.

In cultis et ruderalis ubique. γ in maritimis haud infrequens.

Perennis. Fl. a Junio ad Septembrem usque.

Radix valdè repens, articulata, tunicata, fibris tomentosis.

Culmus erectus, bipedalis, gracilis, teres, striatus, foliosus. Folia patentissima, subsecunda, lineari-lanceolata, acuminata, plana, striata, supra margineque scabra. Vaginæ strictæ, nervosæ, glabræ. Stipula brevissima.

Spica

Spica erectiuscula, bi- vel tri-uncialis, priori similis sed tenuior, spiculisque duplò minoribus. Glumæ omnes lanceolato-subulatæ, acuminatæ, sæpè aristatæ, aristis longitudine variis. Rachis subinde hirsuta. Glumæ interiores ciliatæ. Flosculi numero a tribus ad octo variant.

Folia in γ mucronato-pungentia, glauca, rigida, calycis carinâ seabrâ. Flores subinde vivipari. *Herb. Lightf.*

3. TRITICUM calycibus acuminatis subquinque-*caninum*
nervibus quadrifloris, flosculis aristatis, foliis
planis, radice fibrosâ.

Triticum caninum. Hudf. 58. With. 173. Relb.
56 ? Sibth. 52.

T. n. 58. Gmel. Sib. v. 1. 122. t. 27.

T. n. 1429. Hall. Hist. v. 2. 212.

Elymus caninus. Sp. Pl. 124.

Gramen caninum aristatum, radice non repente.

Morif. sect. 8. t. 1. f. 2. —sylvaticum. Raii

Syn. 390.

Angl. Fibrous or Bearded Wheat-grass.

In sylvis et ad sepes umbrosas, solo calcareo.

Perennis. Fl. Julio.

Radix fibrosa, tomentosa, nequaquam repens. Culmi pluri-
mi, erecti, stricti, bipedales, foliosi; teretes, glaberrimi,
minutè striati. Folia erecto-patentia, lineari-lanceolata,
acuta, nervosa, viridia, plana, utrinque aspera, quandoque
hirsuta. Vaginæ striatæ, glabræ. Stipula obsoletæ. Spica 3-4
uncialis, erectiuscula, demùm nutans, laxiuscula. Spiculæ inferiores
sæpè geminæ, cæteræ alternæ. Glumæ calycinæ æquales,
lanceolatæ, brevius aristatæ, 3- vel 5-nerves, oppositæ, nec, ut in
Elymo Hordeoque, laterales. Flosculi lanceolati, nervosi,
aristati, aristis glumâ duplò longioribus, rectis. Glumæ
interiores ciliatæ.

4. TRITICUM calycibus obtusis multifloris, spicâ *loliace-*
simplici secundâ, flosculis muticis, culmo ra-*um.*
mofo.

Triticum loliaceum. *Engl. Bot.* t. 221. *With.* 174.

T. unilaterale. *Ait. Hort. Kew.* v. 1. 122.

Poa loliacea. *Huds.* 43. *Relb.* 37.

Gramen pumilum, loliaceo simile. *Raii Syn.* 395.

G. loliaceum exile durius. *Rel. Rudb.* 13.

G. exile duriusculum maritimum, foliolis circumvolutis veluti junceis brevibus. *Pluk. Phyt.* t. 32. f. 7.

G. parvum marinum, spicâ loliaceâ. *Morif. sect.* 8. t. 2. f. 6.

Angl. Dwarf Sea Wheat-grafs.

In arenosis maritimis.

On the north coast of Norfolk. *Rev. Mr. Bryant.*

At Lowestoft, Suffolk. Near Landguard Fort.

Rev. Dr. Goodenough. Near South End, Essex, and in the Isle of Wight. *Mr. E. Forster jun.*

Annua. Fl. Junio, Julio.

Radix fibrosa, sæpiùs pubescens. Culmi plures, erecto-patentes, ramosi, rigidi, subcompressi, glabri, foliosi, palmares. Folia lineari-lanceolata, acutiuscula, plana, dorso scabriuscula, exsiccatione involuta. Vaginæ parùm inflatæ, substriatæ, glabræ. Stipula brevis, obtusa, erosa. Spica erecta, rigida, disticha, secunda, rachi flexuosâ, angulatâ, marginatâ, glabrâ, spiculis lanceolatis, alternis, plerumque solitariis; inferioribus sæpè binatis, ternis, vel subinde quaternis. Glumæ calycinæ æquales, obtusæ, carinatæ, obsoletè nervosæ. Flosculi numerosi, imbricati, distichi, glabri, obtusiusculi, mutici, magnitudine valvularum calycinarum. Glumæ interiores ciliatæ. Rachis partialis flexuosa. Habitu, et etiam structurâ florum, *Poa* rigidæ valdè affine est, quâcum genere naturali forsitan conveniat. Variat colorem fusco-violaceo in apricis.

III. TRIGYNIA.

48. MONTIA. *G. Pl.* 41. *Juss.* 313. *Gærtn. t.* 129.

Cal. diphyllus. *Cor.* monopetala, irregularis.

Capf. unilocularis, trivalvis, trisperma.

1. MONTIA.

fontana.

Montia fontana. Sp. Pl. 129. *Huds.* 60. *With.* 175.

Relb. 56. *Sibth.* 53. *Curt. Lond. fasc.* 3. *t.* 8.

Fl. Dan. t. 131.

M. aquatica minor. Mich. Gen. 18. *t.* 13. *f.* 2.

Alfine parva palustris tricoccus. Raii Syn. 352.

Alfineformis paludosa tricarpus, flosculis albis in-
apertis. Pluk. Phyt. t. 7. *f.* 5.

Cameraria arvensis et minor. Dill. Giff. 46. *ap-*
pend. 114. *t.* 6.

Angl. Water Chickweed. Blinks.

Ad fontes, scaturigines, et in campis humidis.

Annua. *Fl.* Aprili, Maio.

Radix fibrosa. Herba glabra, succulenta. Caulis ramo-
fissimus, ramis decumbentibus, radicanibus, foliosis,
rubicundis. Folia opposita, spatulata, integerrima.
Flores pedunculati, albi, sole acriori tantummodo ex-
panfi, post anthesin nutantes. Fructus erectus. Se-
mina tria, reniformia, scabra. Variat partium numero.

49. HOLOSTEUM. *G. Pl.* 42. *Juss.* 299.

Gærtn. t. 130.

Cal. pentaphyllus. *Pet.* 5, erosa. *Capf.* unilo-
cularis, subcylindrica, apice dehiscens.

1. HOLOSTEUM floribus umbellatis.

umbella-
tum.

Holosteum umbellatum. Sp. Pl. 130. *With.* 175.

Engl. Bot. t. 27. *Rose Elem. append.* 445. *t.* 2.

f. 4.

VOL. I.

M

Holo-

Holosteum, quæ Alfine verna glabra, floribus umbellatis. *Dill. Giff. 41. append. 130. t. 6.*

Cerastium umbellatum. *Huds. 201. Dickf. H. Sicc. fasc. 2. 5.*

Caryophyllus Holostius arvensis. *Ger. em. 595. Angl. Umbelliferous Chickweed.*

In muris antiquis, rariùs.

On the walls of Norwich. *Mr. Pitchford.* On walls about Bury. *Sir T. G. Cullum, Bart.*

Annua. Fl. Aprili.

Herba lævis, subglauca, caule basi ramoso, apicem versus hirsuto et viscido. Folia ovata, acuta. Umbella simplex, terminalis, pedunculis bracteatis, maturescente fructu divaricatis. Flores albi, vel rubicundi. Petala variè incisa. Capsula sexdentata. Stamina numero variant,

50. POLYCARPON. *G. Pl. 42. Juss. 299. Gærtn. t. 129.*

Cal. pentaphyllus. Pet. 5, ovata, minima. Caps. unilocularis, trivalvis. Sem. plurima.

tetra- 1. POLYCARPON.
phyllum.

Polycarpon tetraphyllum. *Sp. Pl. 131. Huds. 60. With. 176. Gærtn. t. 129. Dickf. H. Sicc. fasc. 17. 6.*

Paronychia altera Matthioli. *Dalech. Hist. 1213. f. 2.*

Anthyllis marina incana alfinifolia. *Ger. em. 622.*

A. alfinifolia polygonoides major. *Barr. Ic. t. 534.*

Angl. Four-leaved All-seed.

In ruderalis et arenosis ad maris littora australia.

On various parts of the coast of Devonshire and Dorsetshire. On the neck of the Isle of Portland, close to the shingly beach. *Rev. Dr. Goodenough.*

Annua.

Annua. Fl. Maio—Augusto, etiam hyeme.

Caulis ramosissimus, diffusus, divaricatus. Folia opposita vel quaterna, obovata, integerrima, subcarnosa, glabra. Stipulæ et bractæ acuminatæ, scariosæ, albæ. Paniculæ terminales dichotomæ. Flores numerosi, parvi. Calyx albo marginatus. Petala exigua, alba, fugacia, subinde emarginata. Valvulæ capsularum lanceolatæ, margine inflexæ.

Classis IV.

TETRANDRIA.

Stamina 4.

MONOGYNIA.

Stylus 1.

* *Fl. monopetali, monospermi, superi.*

51. DIPSACUS. *Cal. communis polyphyllus, foliaceus; proprius superus. Pappus cyathiformis.*
52. SCABIOSA. *Cal. communis polyphyllus; proprius superus, duplex.*
60. SANGUISORBA. *Cal. uniflorus, inferus. Cor. supera.*

* * *Fl. monopetali, dispermi, superi.*

56. RUBIA. *Cor. campanulata. Fructus baccatus.*
55. GALIUM. *Cor. plana. Fructus ficcus.*
54. ASPERULA. *Cor. tubulosa. Fructus muticus.*
53. SHERARDIA. *Cor. tubulosa. Fructus coronatus. Sem. tridentata.*

* * * *Fl. monopetali, polyspermi, inferi.*

57. EXACUM. *Cor.* hypocrateriformis. *Capsf.* bilocularis, apice dehiscens.
58. PLANTAGO. *Cor.* refracta. *Capsf.* bilocularis, circumscissa.
59. CENTUNCULUS. *Cor.* tubulosa. *Capsf.* unilocularis, circumscissa.
Gentianæ quadrifidæ.

* * * * *Fl. tetrapetali.*

61. EPIMEDIUM. *Nectaria* 4, petalis incumbentia, faccata. *Siliqua* polysperma, super.
62. CORNUS. *Nectaria* nulla. *Drupa* infra, bilocularis.

Cardamine hirsuta. Coronopus didyma.

* * * * * *Fl. apetali.*

63. PARIETARIA. *Cal.* quadrifidus. *Semen* calyce tectum.
64. ALCHEMILLA. *Cal.* octofidus. *Semen* nudum.

DIGYNIA.

Styli 2.

65. BUFFONIA. *Cor.* tetrapetala. *Capsf.* bivalvis, disperma.
Gentianæ quadrifidæ.

TETRAGYNIA.

Styli 4.

66. ILEX. *Cor.* monopetala. *Bacca* tetrasperma.
69. SAGINA. *Pet.* 4. *Caps.* unilocularis.
71. RADIOLA. *Pet.* 4. *Cal.* multifidus. *Caps.*
 octolocularis, octovalvis.
70. TILLÆA. *Pet.* 3—5. *Capsulæ* plures.
67. POTAMOGETON. *Pet.* 4. *Cal.* nullus.
 Sem. 4, nuda.
68. RUPPIA. *Pet.* nulla. *Cal.* nullus. *Sem.* 4,
 pedicellata.

Cerastium tetrandrum.

IV. TETRANDRIA.

I. MONOGYNIA.

51. DIPSACUS. *G. Pl.* 48. *Juss.* 194. *Gærtn. t.* 86.

Cal. communis polyphyllus; *proprius* superus, monophyllus. *Receptaculum* paleaceum. *Pappus* cyathiformis.

fullo-
num.

* 1. DIPSACUS foliis connatis ferratis, paleis recurvis, involucris reflexis.

Dipsacus fullonum. *Linn. Syst. Veg. ed.* 14. 143.
With. 182. *Hudsf. ed.* 1. 49. *Sp. Pl.* 140. β.
Hudsf. 61. β.

D. sativus. *Ger. em.* 1167. *Raii Syn.* 192.

D. albus. *Fuchsf. Hist.* 224. *Id.* 127.

Carduus fullonum. *Lob. Id.* v. 2. 17.

Angl. Manured or Fuller's Teasel.

Ad sepes. *Hudson.* Vix spontaneus.

Biennis. Fl. Julio.

Caulis 5-pedalis, rectus, ramosus, angulatus, aculeatus. Folia radicalia obovata, in petiolum attenuata; caulina connato-perfoliata; omnia glabra, inæqualiter ferrata. Capitulum ovatum, multiflorum, involucre e foliolis pluribus, inæqualibus, lanceolatis, rigidis, mucronatis, recurvis. Flores dilutè purpurei. Paleæ rigidæ, acuminatæ, aduncæ, ad lanas carminandas aptæ, ad quem usum capitula matura frigile lignea disponuntur.

sylvestris. 2. DIPSACUS foliis oppositis ferratis, paleis rectis, involucris inflexis capitulo longioribus.

Dipsacus sylvestris. *Linn. Syst. Veg. ed.* 14. 143.
With. 182. *Relb.* 58. *Sibth.* 54. *Hudsf. ed.* 1. 49.
Curt. Lond. fasc. 3. t. 9. *Jacq. Austr. t.* 402.
Ger. em. 1167. *Raii Syn.* 192. *Cord. Hist.*
105. 2.

Dipsacus

Dipsacus fullonum α . *Sp. Pl.* 140. *Huds.* 61.

D. purpureus. *Fuchs. Hist.* 225. *Ik.* 128.

Labrum Veneris. *Lob. Ic.* v. 2. 19.

Angl. Wild Teasel.

Ad sepes humidiusculas et ad vias.

Biennis. Fl. Julio.

Habitus prioris, sed minùs robustus. Folia radicalia crenata; caulina ferrata, sessilia, minùsque connatò-perfoliata; omnia magis vel minùs aculeata. Involucri foliola inæqualia; majora capitulo longiora, inflexa, quandoque apice foliaceo-dilatata. Paleæ rectæ, flores multùm excedentes, pilosæ.

3. *DIPSACUS* foliis petiolatis appendiculatis, in-*pilosus*,
volucris deflexis capitulo æqualibus.

Dipsacus pilosus. *Sp. Pl.* 141. *Huds.* 61.

With. 182. *Relb.* 59. *Sibth.* 54. *Abbot.* 29.

Curt. Lond. fasc. 1. t. 10. *Jacq. Austr.* t. 248.

D. minor, seu *Virga pastoris.* *Ger. em.* 1168.

Raii Syn. 192.

D. sylvestris capitulo minore. *Morif. sect.* 7. t. 36.

f. 5.

Angl. Small Teasel. Shepherd's Staff.

In humidis, rudерatis, et ad vias, solo calcareo.

Out of St. Benedict's gates, Norwich, and in several other parts of Norfolk. At Matlock bath. Near Deptford, in the London road. *Rev. Dr. Goodenough.*

Perennis. Fl. Augusto.

Radix fusiformis. Caulis ramosus, erectus, sulcatus, aculeatus. Folia petiolata, ovata, ferrata, basi auriculata. Capitula subrotunda, involucris æqualibus, foliaceis, planis, deflexis. Flores albi. Paleæ rectæ, pubescentes, setis ciliatæ.

52. SCABIOSA. *G. Pl.* 48. *Juss.* 194. *Gærtn. t.* 86.

Cal. communis polyphyllus; *proprius* duplex, superus. *Recept.* paleaceum, five nudum.

succisa. 1. SCABIOSA corollulis quadrifidis æqualibus, foliis caulinis dentatis, floribus subglobosis. *Curtis.*

Scabiosa succisa. *Sp. Pl.* 142. *Huds.* 62.

With. 183. *Relb.* 59. *Sibth.* 55. *Curt.*

Lond. fasc. 3. *t.* 10. *Fl. Dan. t.* 279.

S. radice succisa, flore globofo. *Raii Syn.* 191.

Succisa. *Fuchsf. Hist.* 715.

Morsus Diaboli. *Ger. em.* 726.

Angl. Devil's-bit Scabious.

In pascuis copiosè.

Perennis. Fl. Augusto ad Octobrem.

Radix præmorsa, nigricans. Caulis erectus, supernè ramosus, ramis elongatis, unifloris. Folia radicalia ovata, integerrima, obtusiora; caulina lanceolata, plerumque dentata, superiora verò integerrima. Flores violacei, subinde albi.

arvensis. 2. SCABIOSA corollulis quadrifidis radiantibus, foliis pinnatifidis incis, caule hispido.

Scabiosa arvensis. *Sp. Pl.* 143. *Huds.* 62.

With. 183. *Relb.* 60. *Sibth.* 55. *Curt.*

Lond. fasc. 4. *t.* 13. *Dicks. H. Sicc. fasc.* 12. 6.

Fl. Dan. t. 447.

S. major communior, folio laciniato. *Raii Syn.* 191.

S. major vulgaris. *Ger. em.* 719.

Scabiosa. *Fuchsf. Hist.* 716.

Angl. Field Scabious.

In arvis et pratis.

Perennis. Fl. Julio.

Radix

Radix fusiformis, longissima, subramosa. Caulis bi- vel tri-pedalis, ramosus, hispidus, pilis deflexis. Folia sæpius pubescentia; radicalia lanceolata, ferrata; caulina pinnatifida, sæpè incisa; summa integra. Rami patentes, uniflori. Corollæ dilutè purpureæ. Flosculi radii majores, irregulares.

Flores tabaci fumo infecti pulchrè viridescunt.

3. SCABIOSA corollulis quinquefidis radiantibus, *columbaria*. foliis radicalibus ovatis lyratifve crenatis; caulinis pinnatifidis: laciniis linearibus.

Scabiosa columbaria. *Sp. Pl.* 143. *Huds.* 62.

With. 184. *Relb.* 60. *Sibth.* 55. *Fl. Dan.*

t. 314.

S. minor vulgaris. *Raii Syn.* 191.

S. minor five *Columbaria.* *Ger. em.* 719.

Angl. Small Scabious.

In pascuis glareosis vel calcareis.

Perennis. *Fl.* Junio, Julio.

Facile dignoscitur corollulis quinquefidis, maximè radiantibus. Habitu gaudet præcedentis, sed humilior atque omnibus partibus tenuior est. Folia caulina quandoque bipinnatifida. Pappus setis 5, fuscis, longitudine corollæ.

53. SHERARDIA. *G. Pl.* 50. *Juss.* 196. *Gærtn.*

t. 24.

Cor. monopetala, infundibuliformis, supera.

Semina duo, tridentata.

1. SHERARDIA foliis omnibus verticillatis, floribus terminalibus.

Sherardia arvensis. *Sp. Pl.* 149. *Huds.* 66.

With. 185. *Relb.* 63. *Sibth.* 57. *Curt.*

Lond. fasc. 5. *t.* 13. *Dicks. H. Sicc. fasc.*

15. 1. *Fl. Dan. t.* 439.

Sherardia. *Dill. Giff. append.* 96. *t.* 3. *Blair*

Bot. Eff. 155. *t.* 4. *f.* 6. *Hall. Hist. n.* 734.

v. 1. 321.

Asperula

Asperula flore carneo, acuto folio. *Barr. Ic.*
t. 541.

Rubeola arvensis repens cærulea. Raii Syn. 225.
Angl. Blue Shcrardia, or Little Field Madder.

In arvis, præsertim requietis, et inter scgetes.

Annua. Fl. totâ æstatc.

Herba ramosa, diffusa, glabra vel pilosa. Folia plerumque
fena, ovata, acuminata, marginibus carinâque aspera.
Umbella terminalis, sessilis, summo foliorum verticillo
octophyllo involucrata. Calyx superus, diphyllus, fo-
liolis tridentatis, persistentibus, fructum coronantibus.
Flores purpuro-cærulei, tubo gracili.

54. ASPERULA. *G. Pl.* 50. *Juss.* 196.

Cor. monopetala, infundibuliformis, supera. *Se-
mina* duo, globosa.

odorata. 1. ASPERULA foliis octonis lanceolatis, florum
fasciculis pedunculatis, fructu hispido.

Asperula odorata. Sp. Pl. 150. *Huds.* 66.
With. 185. *Relb.* 64. *Sibth.* 57. *Curt. Lond.*
fasc. 4. *t.* 15. *Dicks. H. Sicc. fasc.* 8. 7. *Fl.*
Dan. t. 562.

Asperula. Ger. em. 1124. *Raii Syn.* 224.

Angl. Sweet Woodruff or Woodroof.

In sylvis et umbrosis, præsertim montosis, copiosè.

Perennis. Fl. Maio.

Radix repens. Caules erecti, simplices, læves. Folia
7—9, plerumque 8, in omni verticillo, elliptico-lanceo-
lata, margine scabra. Paniculæ dichotomæ, termina-
les, sæpius ternæ. Flores nivei, interdum suaveolen-
tes. Fructus scetis erecto-patentibus hispidus. Herba
recens inodora, exsiccatione fragrantissima, ut Antho-
xanthum odoratum.

*cynan-
chica.* 2. ASPERULA foliis linearibus quaternis: supe-
rioribus maximè inæqualibus, floribus omni-
bus quadrifidis, fructu glabro.

Asperula

Asperula cynanchica. *Sp. Pl.* 151. *Huds.* 66.
With. 186. *Relb.* 64. *Sibth.* 58. *Engl. Bot.*
t. 33. *Dicks. H. Sicc. fasc.* 17. 5.

A. n. 730. *Hall. Hist.* v. 1. 320.

Galium tetraphyllum montanum cruciatum.
Column. Ecphr. 297. *f.* 1.

*Rubeola vulgaris quadrifolia lævis, floribus pur-
 purascentibus.* *Raii Syn.* 225.

Synanchica. *Ger. em.* 1120.

Angl. Squinancy-wort, or Small Woodruff.

In apricis calcareis.

Perennis. Fl. Junio.

Radix fibrosa, longa. Caules numerosi, basi procumben-
 tes, apice erecti. Folia inferiora quaterna, in ramulis
 obovata; superiora linearia, caulium apicem versus
 valdè inæqualia, ut par intermedium in stipulas dimi-
 nutum videatur. Umbellæ terminales. Flores carnei,
 rubro tri-lineati, suaveolentes. Fructus glaber, nec ut
 in icone Columnæ hispidus.

55. GALIUM. *G. Pl.* 52. *Juss.* 196. *Gærtn. t.* 24.

Cor. monopetala, plana, supera. *Semina* duo,
 subrotundâ.

* *Fructu* glabro.

1. GALIUM foliis quaternis ovatis hirsutis, caule *crucia-*
supernè simplici piloso, pedunculis lateralibus tum.
diphyllis.

Galium cruciatum. *Engl. Bot. t.* 143. *With.* 186.

G. cruciata. *Scop. Carn. v.* 1. 100.

G. n. 709. *Hall. Hist.* v. 1. 314.

Valantia Cruciata. *Sp. Pl.* 1491. *Huds.* 441.

Relb. 377. *Sibth.* 60.

Cruciata. *Ger. em.* 1123. *Raii Syn.* 223.

Angl. Cross-wort. Mug-weed.

In dumetis et ad sepes.

Perennis. Fl. Maio.

Herba latè virens, Caules supernè simplices, elongati, debiles, foliosi. Folia latè ovata, molliuscula, pilosa. Flores numerosi, aurei, corymbosi, axillares, hermaphroditici, intermixti masculis, omnes plerumque quadrifidi, rariùs trifidi, quandoque quinquesidi. Fructus foliis deflexis contactus.

palustre. 2. **GALIUM** foliis quaternis obovatis inæqualibus obtusis, caulibus diffusis supernè ramosis.

Galium palustre. *Sp. Pl.* 153. *Huds.* 67.

With. 187. *Relb.* 65. *Sibth.* 58. *Fl. Dan.* t. 423.

G. n. 719. *Hall. Hist. v. 1.* 317.

G. album. *Ger. em.* 1126.

Molluginis vulgationis varietas minor. *Raii Syn.* 224. *Herb. Buddl.*

Angl. White Water Bed-straw.

In pratis humidis, rivulis et fossis.

Perennis. Fl. Julio.

Caules diffusi, ramosissimi, læves. Folia obovata, semper obtusa, margine sæpiùs scabra, magnitudine inæqualia, subinde fœna. Pedunculi terminales, terni, umbellati, triflori. Flores nivei. Fructus parvus, lævis.

Witheringii. 3. **GALIUM** foliis quinis reflexis lanceolatis aristatis ciliatis, caule erectiusculo simplici scabro. *With.*

Galium montanum. *With.* 187. t. 28, nec *Lin.*

Angl. Rough Heath Bed-straw.

In ericctis montosis subhumidis.

On high but boggy parts of Handsworth heath.
Withering.

Perennis? Fl. Julio.

Caulis spithamæus, debilis, subsimplex, scaber. Folia quina, reflexa, lanceolata, setis ciliata, pilo terminata.
Paniculæ

Paniculæ terminales, trichotomæ. Flores albi, ante expansionem subpurpurei. Antheræ rufæ. Stylus brevissimus, bipartitus. *With.*

Plantam ipse non nisi exsiccatum vidi. Omnino diversa a *G. montano* *Herb. Linn.* proculdubio est.

4. *GALIUM* foliis senis obovatis obtusis mucro-*saxatile*.
natis, caule ramosissimo prostrato glabro,
fructu granulato.

Galium saxatile. Sp. Pl. 154.

G. montanum. Hudf. 67. Relb. 66. Cullum. 55.

G. procumbens. With. 187. Sibth. 59.

G. album minus. Petiv. Herb. Brit. t. 30. f. 6.

Mollugo montana minor, Galio albo similis.

Raii Syn. 224. Herb. Buddl.

Angl. Smooth Heath Bed-straw.

In ericetis et montosis frequens.

Perennis. Fl. Julio, Augusto.

Radix repens. Caules ramosissimi, diffusi, procumbentes, glabri. Folia sena, patentia, obovata, mucronata, margine sæpius aspera. Flores lactei, copiosi. Paniculæ terminales, trichotomæ. Fructus granulatus.

Variat foliis ferè linearibus.

5. *GALIUM* foliis senis lanceolatis retrorsum fer-*uligino-*
rato-aculeatis mucronatis rigidis, corollis fruc-*sum.*
tu majoribus.

Galium uliginosum. Sp. Pl. 153. Hudf. 68.

With. 188. Relb. 65. Sibth. 58.

Aparine palustris minor Parisiensis, flore albo.

Dill. in Raii Syn. 225.

Rubia quædam minor. Baub. Hist. v. 3. 716?

Angl. Rough Marsh Bed-straw.

In aquosis, et pratis humidis.

Perennis. Fl. Augusto.

Caules ramosi, debiles, scabri. Folia sena, variè patentia, lanceolata, basi attenuata, apice aristata, margine carinaque

carinâque retrorsum aculeata. Pedunculi terminales, sæpius terni, glabri, bifidi vel trifidi. Flores albi. Fructus lævis, parvus.

Folia prope marginem sæpè aculeis antrorsum porrectis, simplici serie digestis, aspera sunt, dum ipse margo semper retrorsum aculeatus invenitur.

erectum. 6. **GALIUM** foliis suboclonis lanceolatis antrorsum aculeato-ferrulatis mucronatis, paniculis trichotomis, caule læviusculo flaccido.

Galium erectum. *Huds.* 68. *With.* 189. *Dicks.* *H. Sicc. fasc.* 17. 2.

Angl. Upright Marsh Bed-straw.

In pratis et pascuis subhumidis.

Perennis. Fl. Junio, Julio.

Caules ramosissimi, flaccidi, erectiusculi, angulis subinde scabriusculi, sæpius omninò læves, apice paniculati, multiflori. Folia sena, septena vel octona, elliptico-lanceolata, aristata, glabra, margine aculeato-ferrulata. Paniculae terminales et laterales, trichotomæ, foliolosæ, multifloræ. Corolla alba, acuminata. Fructus lævis, parvus.

Galium austriacum, a Jacquino missum, omninò diversum est, foliis linearibus, duplò longioribus, margine proculdubiò lævibus.

tricorne. 7. **GALIUM** foliis suboclonis lanceolatis margine, cauleque, retrorsum aculeatis, pedunculis axillaribus trifloris, fructu granulato nutante.

Galium tricorne. *With. ed.* 2. 153.

G. spurium. *Huds.* 68. *With.* 190. *Relb.* 65. *Sibth.* 59.

G. n. 725. *Hall. Hist.* v. 1. 319.

Valantia Aparine. *Mart. Fl. Rust.* 1. 122.

Aparine femine læviore. *Raii Syn.* 225.

A. femine lævi. *Vaill. Paris.* 1. 4. f. 3. a. a.

A. foliis brevioribus et femine læviore. *Morif. sect.* 9. 332.

Angl.

Angl. Corn Bed-straw. Three-flowered Goose-grass.

In arvis, haud frequens.

In the isle of Thanet, in Surrey and near Stamford.

Huds. Oxfordshire. *Morif. and Sibth.* Yorkshire. *Mr. Woodward.* Gloucestershire. *Rev. Mr. Swayne.* Near Barton, Norfolk. *Mr. D. Turner.*

Annua. Fl. Julio.

Radix parva. Caulis simplicissimus, foliosus, tetragonus, angulis retrorsum scaber. Folia plerumque octona, patentia, lanceolata, mucronata, margine, subinde carinâ, retrorsum cartilagineo-ferrata. Pedunculi axillares, folitarii, alterni, longitudine vix foliorum, quadranguli, angulis asperis, apice trifidi, triflori, plerumque diphylli. Flores parvi, albidî, rarius omnes fertiles. Pedunculi fructiferi arcuato-reflexi. Fructus majusculus, granulatus, vel tuberculis parvis asperatus, nec hispidus.

A Valantiâ Aparine Linnæi (Aparine semine Coriandri Parkinsonii. *Vaill. t. 4. f. 3. b.*) abundè differt foliis retrorsum ferratis, nec antorsum margine scabris, fructisque tuberculis longè minoribus. Galium spurium *Herb. Linn.* etiam, ni fallor, a nostrâ plantâ discrepat, foliis tantum senis, pedunculis fructiferis erecto-patentibus folio duplò longioribus, fructu lævi vix granulato vel exasperato.

8. GALIUM foliis octonis lineari-lanceolatis mucronatis integerrimis subpubescentibus, pedunculis dichotomis, fructu lævissimo. *pusillum.*

Galium pusillum. *Sp. Pl.* 154. *Huds.* 69.

With. 188. *Engl. Bot. t.* 74.

G. scabrum. *Jacq. Fl. Austr. t.* 422.

G. album supinum multicaule. *Dill. in Raii Syn.*

224?

Angl. Least Mountain Bed-straw.

In montosis calcareis.

Near Kendal. *Huds.* At Matlock-bath, Derbyshire, in great plenty.

Perennis. Fl. Julio, Augusto.

Caules plurimi, spithamæi, ramosi, quadranguli, sæpiùs pilis patentibus brevibus hispidi, interdum glabri; basi decumbentes. Folia plerumque octona, lineari-lanceolata, mucronata, integerrima, subrevoluta, sæpiùs pilis brevibus scabra; superiora omninò glabra; infima parva, imbricata, sena, obovata. Pedunculi paniculati, dichotomi. Corolla alba, mutica. Fructus omninò lævis et glaber, stylo persistente, semibifido, coronatus.

verum. 9. **GALIUM** foliis octonis linearibus sulcatis integerrimis scabris, floribus paniculatis congestis.

Galium verum. Sp. Pl. 155. *Huds.* 69. *With.* 191.

Relb. 66. *Sibth.* 59. *Curt. Lond. fasc.* 6. t. 13.

Mart. Fl. Rust. t. 54. *Dicks. II. Sicc. fasc.* 17. 4.

Gallium. Fuchs. Hist. 196.

G. luteum. Raii Syn. 224. *Ger. em.* 1126.

Angl. Yellow Bed-straw. Cheese-rennet.

In collibus, dumetis, et ad agrorum margines:

Perennis. Fl. Julio, Augusto.

Radix repens, fulva. Caules erectiusculi, sesquipedales, basi ramosi. Folia linearia, mucronata, subreflexa, integerrima, supra scabra. Panicula ramosissima, ramulis coarctatis, corymbosis, multifloris. Flores aurei, odore melleo. Antheræ defloratæ fuscæ. Fructus lævis.

Mollugo. 10. **GALIUM** foliis octonis ellipticis obtusiusculis mucronatis margine scabris, floribus paniculatis divaricatis.

Galium Mollugo. Sp. Pl. 155. *Huds.* 68.

With. 189. *Relb.* 67. *Sibth.* 59. *Fl. Dan.*

t. 455.

Mollugo vulgator. Raii Syn. 223.

Rubia

Rubia sylvestris. *Ger. em.* 1118. *Fuchsf. Hist.* 281.

β. *Galium scabrum.* *With.* 190. nec Jacq. Ex ipso *Witheringio* accepi.

Angl. Great Hedge Bed-straw.

In sepibus et dumetis.

β. Hedge-row in a marly soil, on the side of the Red-house lane near Worcester. *Dr. Stokes.* *With.*

Perennis. Fl. Julio, Augusto.

Caules 2—4-pedales, flaccidi, læviusculi, supra genicula incrassati. Folia elliptico-lanceolata, subinde obovata, mucronulata, plerumque glabra, margine (in nostris ut et Linnæanis) ferrulato-scabra. Panicula ramosissima, dichotoma, sparsa, multiflora. Flores albi.

Folia ut et caulis in β piloso-scabra.

§ 1. **GALIUM** foliis subsenis lanceolatis mucronatis margine, cauleque, scabris, pedunculis bifidis, fructu granulato. *anglicum.*

Galium anglicum. *Huds.* 69. *With.* 191. *Engl. Bot. t.* 384.

G. parisiense. *Relb.* 67.

Aparine minima. *Raii Syn.* 225. t. 9. f. 1.

Angl. Wall Bed-straw.

In muris, et arenosis aridis.

In Kent. *Huds.* Binham Priory, Norfolk. *Mr. Crowe.* Carrow Abbey near Norwich. *Mr. Pitchford.* At Mildenhall, Suffolk. *Rev. C. Sutton.*

Annua. Fl. Junio, Julio.

Radix fibrosa, parva. Caules palmares, vel spithamæi, ramosi, curvi, retrorsum scabri. Folia sena vel septena, parva, deflexa, lanceolata, margine ferrulato-scabra, interdum supra exasperata. Rami floriferi alterni, subdichotomi, pauciflori. Flores luteo-virescentes. Fructus granulato-scabriusculus.

A *G. parisiensi* Linn. vix differt, nisi floribus luteo-virescentibus nec atro-purpureis, fructu granulato nec setis hispido; fortè mera varietas est.

** *Fructu hispido.*

boreale. 12. **GALIUM** foliis quaternis lanceolatis trinervi-
bus glabris, caule erecto, fructibus hispidis.

Galium boreale. *Sp. Pl.* 156. *Huds.* 70.

With. 192. *Dicks. H. Sicc. fasc.* 3. 5. *Engl.*

Bot. t. 105. *Fl. Dan. t.* 1024.

Mollugo montana erecta quadrifolia. *Raii Syn.*
224.

Angl. Cross-leaved Bed-straw.

In petrosis subalpinis.

In the mountainous parts of the north of England,
as well as Wales and Scotland, not unfrequent.

Just above Kirkby-Lonsdale bridge plentifully.

Perennis. Fl. Julio.

Radix repens, ramosa, rubro-fusca. Caules sesquipedales, erecti, scabriusculi. Folia quaterna, inæqualia, ovato-lanceolata, obtusa, mutica, integerrima, trinervia, margine præcipuè scabra. Panicula trichotoma. Pedunculi subcorymbosi. Flores lactei. Fructus fetis albis muricatus.

Aparine. 13. **GALIUM** foliis octonis lanceolatis carinatis
scabris retrorsùm aculeatis, caule flaccido,
fructibus hispidis.

Galium Aparine. *Sp. Pl.* 157. *Huds.* 70.

With. 192. *Relb.* 67. *Sibth.* 60. *Curt.*

Lond. fasc. 2. t. 9. *Mart. Fl. Rust. t.* 104.

Fl. Dan. t. 495. *Woodv. Suppl. t.* 269.

Dicks. H. Sicc. fasc. 17. 3.

Aparine. *Raii Syn.* 225. *Ger. em.* 1122.

Angl. Goose-grass, or Cleavers.

Ad fepes vulgaris.

Annua. Fl. Maio, Augusto.

Radix fibrosa. Caulis tripedalis, flaccidus, fragilis, ramosus, retrorsum scaber. Folia obovato-lanceolata, mucronata, integerrima, scabra, margine carinâque retrorsum aculcata. Pedunculi axillares et terminales, simplices vel bifidi, scabri. Flores parvi, albidî. Fructus magnus, fetis uncinatis hispidus.

56. RUBIA. *G. Pl.* 52. *Juss.* 197.

Cor. monopetala, campanulata, supera. *Baccæ peregrinæ*, monospermæ.

1. RUBIA foliis subquaternis ellipticis suprà nitidis lævibus, floribus quinquefidis.

Rubia peregrina. *Sp. Pl.* 158. *Huds.* 65. *Culm.* 56.

R. anglica. *Huds. ed.* 1. 54.

R. tinctorum. *With.* 193.

R. sylvestris aspera, quæ sylvestris Dioscoridis. *Raii Syn.* 223. *ed.* 1. 317. *Morif. sect.* 9. t. 21. f. 2.

Angl. Wild Madder. *Petiv. Herb.* t. 30. f. 3.

In dumetis et ad sepes anguli Britanniae australioris.

On St. Vincent's rock, and in Devonshire. *Ray.*

In Cornwall. *F. Borone.* On Tunbridge rocks.

Rev. Dr. Goodenough. At Chepstow.

Perennis. Fl. Junio, Julio.

Radix rubra. Caulis ramosus, diffusus, angulis asper, perennis. Folia quaterna, rarius sena, perennantia, elliptica, acuminata, nitida, lævia, nervo utrinque, margineque, denticulato-asperissima. Paniculae dichotomæ, terminales. Flores flavescentes, semper quinquefidi, pentandri, atque calyce orbatî, optimè monente Cl. Swaynco apud Witheringium.

Rubia Halleri n. 708. ab amicissimo Davallio missa, est omnino R. tinctorum Linnæi, cui folia ovato-lanceolata supra scabra sunt, flores sæpius, nec semper, quadrifidi, calyce etiam destituti.

57. EXACUM. *G. Pl.* 57. *Juss.* 142. *Gærtn.*
t. 114.

Cal. quadrifidus. *Cor.* hypocrateriformis, tubo
ventricoso. *Caps.* bisulca, bilocularis, poly-
sperma, apice dehiscens. *Stigma* capitatum.

filiforme. 1. EXACUM foliis sessilibus, caule filiformi di-
chotomo, pedunculis elongatis.

Exacum filiforme. *Engl. Bot.* t. 235. *With.* 194.

Gentiana filiformis. *Sp. Pl.* 335. *Huds.* 103.

Fl. Dan. t. 324.

Centaurium palustre luteum minimum nostras.

Raii Syn. 235. *Vaill. Paris.* t. 6. f. 3.

Angl. Least Gentianella.

In palustribus spongiosis vel arenosis.

In Dorsetshire. *Dr. Pulteney.* Cornwall and De-
vonshire. *Mr. Stackhouse.* On Wareham com-
mon, by the sides of the road as you go to Corfe
Castle. *Rev. Dr. Goodenough.*

Annua. Fl. Julio.

Radix fibrosa. Herba lævis, amara. Caulis dichotomus,
tenellus, vix triuncialis, teretiusculus, basi præcipue
foliosus. Folia opposita, sessilia, spatulato-lanceolata,
integerrima. Pedunculi longissimi, tetragoni, uniflori,
erecti. Flores aurei, non nisi soli meridiano expansi.
Laciniae calycis acutæ, margine membranaceæ. Sta-
mina parùm inæqualia, fauci corollæ inserta, incurva.
Stigma capitatum, vix emarginatum.

58. PLANTAGO. *G. Pl.* 57. *Juss.* 90. *Gærtn.*
t. 51.

Cal. quadrifidus. *Cor.* quadrifida, infera; limbo
reflexo. *Stam.* longissima. *Caps.* bilocularis,
circumscissa.

maj. 1. PLANTAGO foliis ovatis glabriusculis petiolo
brevioribus, seapo tereti, spica floribus imbri-
cata, seminibus plurimis.

Plantago

Plantago major. *Sp. Pl.* 163. *Huds.* 63.
With. 195. *Relb.* 61. *Sibth.* 56. *Curt.*
Lond. fasc. 2. t. 11. Fl. Dan. t. 461. Dreves.
Bilderb. t. 25.

P. latifolia vulgaris. *Raii Syn.* 314.

P. latifolia. *Ger. em.* 419.

P. rubea. *Brunf. Herb. v. 1. 25.*

β. *P. latifolia glabra minor.* *Dill. in Raii Syn.* 314.
Huds. 63. *With.* 195. *var. 1.*

γ. *P. major, paniculâ sparsâ.* *Raii Syn.* 314.
Huds. 63. *With.* 195. *var. 5.*

P. panniculis sparsis. *Ger. em.* 420.

δ. *P. rosea spicata.* *Ger. em.* 420.

Huds. 63 δ. *With.* 195. *var. 4.*

Angl. Greater Plantain.

In pratis, pascuis, et ad vias ubique.

Perennis. *Fl. totâ æstate.*

Folia longiùs petiolata, ovata, sæpè subdentata, 7—9 nervia, glabriuscula. Spica longissima, linearis, multiflora, arctè imbricata. Semina, monente Curtisio, plurima.

Folia in β trinervia, spica brevior, pauciflora. γ spica paniculata, abortiva, foliolosa. δ spica abortiva, e foliolis composita, vel rosaceè expansis, vel in pyramidem imbricatis.

2. *PLANTAGO* foliis ovatis pubescentibus *pe-media.*
 tiolo longioribus, scapo tereti, spicâ cylindricâ,
 seminibus solitariis.

Plantago media. *Sp. Pl.* 163. *Huds.* 63.

With. 196. *Relb.* 61. *Sibth.* 56. *Curt.*

Lond. fasc. 4. t. 14. Fl. Dan. t. 581.

P. major incana. *Raii Syn.* 314.

P. incana. *Ger. em.* 419.

P. major. *Brunf. Herb. v. 1. 23.*

Angl. Hoary Plantain.

In pratis et pascuis solo calcareo vel glareoso passim.

Perennis. Fl. totâ æstate.

Folia pubescentia, minùs obtusa, plerumque 5-nervia, depressa, petiolis dilatatis, brevissimis. Spica brevis, densa, staminibus dilutè purpureis, antheris albis.

lanceolata. 3. **PLANTAGO** foliis lanceolatis utrinque acutis, spicâ ovatâ nudâ, scapo angulato.

Plantago lanceolata. *Sp. Pl.* 164. *Huds.* 64.

With. 196. *Relb.* 62. *Sibth.* 56. *Curt.*

Lond. fasc. 2. *t.* 10. *Mart. Fl. Rust.* *t.* 67.

Fl. Dan. *t.* 437. *Engl. Bot.* *t.* 507.

P. quinquenervia. *Ger. em.* 422. *Raii Syn.* 314.

P. minor. *Brunf. Herb.* *v.* 1. 24.

Angl. Ribwort Plantain.

In pratis et pascuis vulgaris.

Perennis. Fl. Junio, Julio.

Folia elongata, erecta, 3—5-nervia, saturatè viridia. Spica ovata, brevis, albo nigroque varia. Variat spicâ foliolis obvallatâ. *D. Woodward.* Quandoque paniculata, vel rosacea, modo *P. majoris* evadit.

maritima. 4. **PLANTAGO** foliis linearibus, subintegerrimis canaliculatis basi lanatis, spicâ cylindricâ, scapo tereti.

Plantago maritima. *Sp. Pl.* 165. *Huds.* 64.

With. 197. *Relb. Suppl.* 2. 9. *Engl. Bot.*

t. 175. *Fl. Dan.* *t.* 243. *Lightf.* 118.

P. marina. *Raii Syn.* 315.

P. an alpina angustifolia J. B. ? *Raii Syn.* 315.

P. montana. *Huds. ed.* 1. 53.

Coronopus. *Ger. em.* 425.

β. *Plantago Loeslingii.* *Huds. ed.* 1. 52. *nec Linn.*

P. marina. *Ger. em.* 423.

Angl. Sea Plantain.

In maritimis limosis, etiâ in fissuris rupium alpinarum.

Perennis. Fl. Augusto.

Magnitudine insigniter variat. Folia canaliculata, ecarinata, sæpiùs integerrima, subinde irregulariter dentata, præcipuè in maritimis.

5. **PLANTAGO** foliis linearibus pinnato-dentatis, scapo tereti. *Coronopus.*

Plantago Coronopus. *Sp. Pl.* 166. *Huds.* 64.

With. 198. *Sibth.* 56. *Fl. Dan. t.* 272.

P. foliis laciniatis, Coronopus dicta. *Raii Syn.* 315.

Cornu Cervinum. *Ger. em.* 427.

β. *Plantago gramineo folio hirsuto, minor, capitulo rotundo brevi.* *Dill. in Raii Syn.* 316.

Angl. Bucks-horn Plantain. *Star of the Earth.*

In arenosis, glareosis, et maritimis passim.

β. In Portland Island. *Mr. Yalden.*

Annua. Fl. totâ æstate.

Folia stellatim depressa, pilosa, pinnatifida, subinde bipinnatifida, in β ferè integra. Scapi adscendentes, teretes. Spica cylindrica, uncialis, in arenosis quandoque brevissima, pauciflora, et ferè capitata. Antheræ, monente Witheringio, membranâ lanceolatâ terminatæ. Distinctissima a *P. Loeßlingii*.

59. **CENTUNCULUS.** *G. Pl.* 58. *Juss.* 95.

Gærtn. t. 50.

Cal. quadrifidus. *Cor.* quadrifida, infera, tubulosa, limbo patente. *Stam.* brevia, nuda.

Capf. unilocularis, circumscissa, polysperma.

1. **CENTUNCULUS.**

minimus.

Centunculus minimus. *Sp. Pl.* 169. *Huds.* 63.

With. 198. *Abbot.* 30. *Curt. Lond. fasc.* 3.

t. 11. *Dicks. H. Sicc. fasc.* 7. 2. *Fl. Dan.*

t. 177. *Engl. Bot. t.* 531.

Centunculus. *Dill. in Raii Syn.* 1. *Dill. Giff.* 161.

app. 111. *t.* 5.

Anagallis,

Anagallis, quæ *Alfine palustris minima*, flosculis albis, fructu *Coriandri* exiguo. *Mentz. Pugill. t. 7. Vaill. Paris. t. 4. f. 2.*

Anagallidastrum exiguum, foliis lanceolatis, alternis, flore albo, fugaci, et vix conspicuo. *Mich. Gen. 14. t. 18. f. 2.*

Angl. Bastard Pimpernel. Small Chaff-weed.

In uliginosis, vel ericetis inundatis, rariùs.

On Hounslow Heath. *Sir William Watfon.* Near Hampton Court. *Hudson.* Ashford Common. *Curtis.* Iwer Heath in abundance. *Rev. Dr. Goodenough.*

Annua. Fl. Junio, Julio.

Radix fibrosa. Caulis basi parùm ramosus, adscenders, uncialis vel biuncialis, foliosus, subangulatus, glaber. Folia alterna, sessilia, patentia, ovata, integerrima, glabra. Flores solitarii, axillares, sessiles, albi vel rubicundi, fugaces, ardentissimis diei horis expansi. Corolla regularis, persistens. Capsula globosa, stylo mucronata, glabra.

60. SANGUISORBA. *G. Pl. 58. Juss. 336.*

Cal. quadrifidus. Germen inter calycem corollamque. Semen unicum.

officina- 1. SANGUISORBA spicis ovatis.
lis.

Sanguisorba officinalis. Sp. Pl. 169. Hudf. 65. With. 199. Relb. 63. Sibth. 57. Abbot. 37. Mart. Fl. Rust. t. 142. Fl. Dan. t. 97.

S. major flore spadiceo. Raii Syn. 203.

Pimpinella sylvestris. Ger. em. 1045.

Angl. Great Burnet.

In pratis et pascuis calcareis humidiusculis Angliæ borealis.

Perennis. Fl. Junio, Julio.

Folia impari-pinnata, foliolis cordatis, acutè crenatis, glabris.

glabris. Caulis bipedalis, erectus, nudiusculus, ramosus. Spicæ ellipticæ, obtusæ, atro-sanguinæ, floribus superioribus primùm expansis. Corolla extus basi pilosa. Germen tunicâ glabrâ, e basi corollæ ortâ, arctè investitur.

61. EPIMEDIUM. *G. Pl.* 59. *Juss.* 287.

Petala 4. *Nectaria* 4, saccata, petalis incumbentia. *Cal.* petalis oppositus, caducus. *Siliqua* supera, unilocularis, polysperma.

1. EPIMEDIUM.

Epimedium alpinum. *Sp. Pl.* 171. *Cullum.* 60.

With. 199. *Engl. Bot. t.* 438. *Ait. Hort.*

Kew. v. i. 157. *Allion. Ped. v. i.* 238.

Epimedium. *Ger. em.* 480. *Lind. Alsat.* 136. *t. 6.*

Angl. Barren-wort.

alpinum.

In dumetis alpinis.

On Carrock Fell, Cumberland. *Mr. Thomas Hutton.* Bingley woods, Yorkshire. *Dr. Richardson.*

Perennis. Fl. Maio.

Radix repens. Caulis nullus. Folium erectum, solitarium, irregulariter ternato-supradecompositum, floriferum. Foliola petiolata, opposita, cordata, dependentia, acuta, serrata, glabra, submembranacea, lætè virentia, subtus glauca. Petioli teretes, glabri. Panicula e geniculo medium versus petioli communis orta, erecta, parùm ramosa, pedunculis alternis, rubentibus, pilosis, pilis viscidis. Flores nutantes. Calyx tetraphyllus, virens, adhuc inaperto flore caducus. Petala ovata, æqualia, concava, atro-sanguinea. Nectaria flavo-hyalina, obtusa. Antheræ filamentis longitudinaliter adnatæ, operculatæ, ut in *Leontice* et *Lauro*. Germen ovatum, superum. Siliqua bivalvis, polysperma.

62. CORNUS. *G. Pl.* 59. *Juss.* 214. *Gærtn. t.* 26.

Cal. quadridentatus. *Petala* 4. *Drupa* infera, nuce biloculari.

1. COR-

- sanguinea.* 1. CORNUS ramis rectis, foliis ovatis utrinque viridibus, cymis nudis planis.
 Cornus sanguinea. *Sp. Pl.* 171. *Huds.* 70. *With.* 200. *Relb.* 68. *Sibth.* 61. *Engl. Bot.* t. 249. *L'Herit. Corn.* 5. *Fl. Dan.* t. 481.
 C. fœmina. *Raii Syn.* 460. *Ger. em.* 1467.
Angl. Wild Cornel-trec. Dog-wood.

In sylvis et sepibus.

Frutex. Fl. Junio.

Lignum durum. Rami oppositi, teretes, sanguinei. Folia opposita, petiolata, ovata, integerrima, venosa, glabra, opaca, utrinque viridia, subtus pallidiora, decidua, fero autumnio rubescentia. Flores albidī, graveolentes, Drupæ globosæ, atro-purpureæ, amaræ, nauscosæ.

- Suecica.* 2. CORNUS herbacea, ramis binatis, umbellâ axillari pedunculatâ involucreatâ, foliorum nervis omnibus subdistinctis.

Cornus Suecica. *Sp. Pl.* 171. *With.* 200. *Engl. Bot.* t. 310. *Fl. Dan.* t. 5. *Linn. Fl. Lapp.* t. 5. f. 3. *Lightf.* 119. *Pennant. Tour in Scotl.* t. 39.

C. herbacea. *Huds.* 71.

C. pumila herbacea, Chamæpericlymenum dicta, *Dill. Elth.* 108. t. 91.

Chamæpericlymenum. *Raii Syn.* 261. *Ger. em.* 1296. *Park.* 1461.

Angl. Dwarf Cornel.

In pratis alpinis.

On the Cheviot hills, and in some parts of Yorkshire; also on the mountains of Scotland.

Perennis. Fl. Junio, Julio.

Radix repens. Caules herbacei, spithamæi, foliosi, erecti, quadranguli, apice bifidi. Umbella interramea, pedunculata, erecta, simplex, involuero tetraphyllo, albo vel rubicundo, persistente, demum viridescente et ampliato,

pliato. Flores parvi, atro-purpurei, brevius pedicellati. Drupæ globosæ, rubræ, subdulces. Folia ovata, opposita, sessilia, nervis omnibus e basi ferè ortis, nec, ut in Corno canadensi, costâ centrali ramos multifariam exferenti, monente amicissimo Salisburio.

63. PARIETARIA. *G. Pl.* 544. *Juss.* 404.
Gærtn. t. 119.

Cal. quadrifidus, inferus. *Cor.* nulla. *Stamina* elastica. *Semen* 1, calyce elongato tectum. Flores aliquot fœminei, calyce immutato.

1. PARIETARIA foliis lanceolato-ovatis basi *officina-*
uninervibus, involucris laciniis ovatis, caule *lis.*
erectiusculo.

Parietaria officinalis. Sp. Pl. 1492. *Huds.* 442.
With. 201. *Relb.* 378. *Sibth.* 62. *Curt.*
Lond. fasc. 4. *t.* 63. *Fl. Dan. t.* 521.

Parietaria. Ger. em. 331. *Raii Syn.* 158.

Parietaria, herba. Pharmac. Lond.

Angl. Pellitory of the wall.

In ruderatis et muris.

Perennis. Fl. Junio ad Septembrem.

Caulis ramosus, patens, rubens, nitidus, foliosus. Folia elliptico-lanceolata, obliqua, acuminata, subtus pallida, nervis lateralibus non in petiolum decurrentibus, sed longè supra basin e costâ centrali enatis, quâ notâ e *P. judaicâ* et *indicâ* dignoscitur. Involucrum septemfidum, triflorum, floribus omnibus fertilibus, centrali fœmineo. Frigoris impatiens herba.

64. ALCHEMILLA. *G. Pl.* 64. *Juss.* 337.
Gærtn. t. 73.

Cal. octofidus, laciniis alternis minoribus, inferus. *Cor.* nulla. *Semen* 1, nudum.

1. ALCHEMILLA foliis plicatis lobatis.

vulgaris.

Alchemilla vulgaris. Sp. Pl. 178. *Huds.* 71.
With.

- With.* 204. *Relb.* 68. *Sibth.* 61. *Fl. Dan.*
t. 693. *Abbot.* 36. *t.* 1. *Engl. Bot. t.* 597.
Alchemilla. *Ger. em.* 949. *Raii Syn.* 158.
Stellaria. *Camer. Epit.* 908.
β. A. minor. *Hudsf. ed.* 1. 59.
A. alpina pubescens minor. *Pluk. Phyt. t.* 240.
f. 2.
Angl. Common Ladies' Mantle.

In pratis et pascuis submontosis.

Perennis. *Fl.* Junio, Julio.

Caules adscendentes, paniculati, floribus confertis. Folia plicata, 7—9-loba, lobis obtusis, crenato-ferratis; caulina minora, subsessilia, stipulis maximis, incis. Omnia in *β* magis sericea atque concinniora.

alpina. 2. *ALCHEMILLA* foliis digitatis ferratis subtus sericeis.

Alchemilla alpina. *Sp. Pl.* 179. *Hudsf.* 71.

With. 205. *Engl. Bot. t.* 244. *Fl. Dan.*

t. 49. *Dicksf. H. Sicc. fasc.* 15. 2.

A. Alpina pentaphyllos. *Raii Syn.* 158.

Pentaphyllum petrosum, Heptaphyllum Clusii.

Ger. em. 988.

Stellaria argentea. *Camer. Epit.* 909.

Angl. Alpine Ladies' Mantle.

In rupibus alpinis, præcipuè micaceis.

In Cumberland, Westmoreland, North Wales, and the Highlands of Scotland.

Perennis. *Fl.* Julio.

Folia quinquepartita, lobis apice ferratis, subtus pulcherrimè sericeo-argentea. Caulis paniculatus, erectus. Flores extus sericei.

arvensis. 3. *ALCHEMILLA* foliis planis trilobatis incis. *Alchemilla Aphanes.* *Leers.* 54. *Sibth.* 61. *Abbot.* 36.

A. n.

A. n. 1569. *Hall. Hist.* v. 2. 262.

Aphanes arvensis. *Sp. Pl.* 179. *Huds.* 72.

Relb. 69. *With.* 4. *Fl. Dan.* t. 973.

Percepier Anglorum. *Ger. em.* 1594. *Raii*

Syn. 159.

Angl. Field Ladies' Mantle. Parsley Piert.

In arvis, præsertim requietis, sterilibus.

Annua. Fl. totâ æstate.

Caulis ramosissimus, diffusus, foliosus. Folia alterna, petiolata, palmato-tripartita, crenata et incisa, pilosa, lineata, plana. Stipulæ petiolo adnatæ, incisæ. Calyx urceolatus, angulatus, certè octofidus, laciniis alternis minimis. Semina quandoque bina.

II. DIGYNIA.

65. BUFFONIA. *G. Pl.* 65. *Juss.* 300.

Gærtn. t. 129.

Cal. tetraphyllus. *Cor.* tetrapetala. *Caps.* unilocularis, disperma.

† 1. BUFFONIA. *With.* 205.

Bufonia tenuifolia. *Sp. Pl.* 179. *Huds.* 72. *tenuifolia.*

Alfine polygonoides tenuifolia, flosculis ad longitudinem caulis velut in spicam dispositis nostra. *Raii Syn.* 346. *Pluk. Phyt.* t. 75. f. 3.

Herniaria angustissimo gramineo folio erecta.

Magnol. Hort. Monsp. 97. tab. 15.

Polygonum

Polygonum angustissimo gramineo folio erectum! *Magnol. Bot. Monsp.* 211.
Angl. Slender Buffonia.

In maritimis rarius.

Near Boston, Lincolnshire. *Ray's Syn.* On Hounslow heath. *Doody.* No other botanist has found it on Hounslow heath; and it is the opinion of Sir Jos. Banks, that Bupleurum tenuissimum was carelessly mistaken for it at Boston.

Annua. Fl. Junio.

Caules plurimi, erecti, stricti, foliis vestiti oppositis, subulatis, adpressis, basi membranaceâ connatis. Flores apicem versus caulis, pedunculati, albi. Foliola calycina acuta, margine membranacea. Exemplaria spontanea nondum vidi.

Capsula obovata, seminibus binis, compressis, tuberculatis, basi capsulæ insertis. *Gærtner.*

III. TETRAGYNIA.

66. ILEX. *G. Pl.* 67. *Juss.* 379.

Cal. 4- vel 5-dentatus. *Cor.* rotata. *Styl.* nulli.
Bacca tetrasperma.

Flores aliquot masculi, quadrifidi.

Aquifolium. 1. ILEX foliis ovatis acutis spinosis.

Ilex Aquifolium. *Sp. Pl.* 181. *Huds.* 446.
With. 210. *Relb.* 382. *Sibth.* 64. *Fl. Dan.*
t. 508. *Engl. Bot.* t. 496.

Agri-folium.

Agrifolium. *Ger. em.* 1338. *Raii Syn.* 466.

β. *A. baccis luteis* nondum descriptum. *Raii Syn.* 466.

Angl. Holly-tree. *Hunt. Evel. Sylv. tab. ad* p. 383.

In sepibus et frutetis.

Arbor. Fl. Maio.

Arbor formosa, longæva, sempervirens, lucida, cortice lævi, plumbeo, maceratione glutinoso, ad viscum conficiendum usitatissimo. Folia rigida, nitida, lobis spinosis; in ramis antiquioribus integra. Flores axillares, fasciculati, albi; primordiales sæpius imperfecti, masculis immixti; posteriores omnes hermaphroditici. Baccæ puniceæ, diutius persistentes, quandoque luteæ.

67. POTAMOGETON. *G. Pl.* 67. *Juss.* 19.
Gærtn. t. 84.

Cal. nullus. *Petala* 4. *Stylus* nullus. *Semina* 4.

1. POTAMOGETON foliis superioribus oblongo-natans, ovatis petiolatis natantibus.

Potamogeton natans. *Sp. Pl.* 182. *Huds.* 74.

With. 211. *Relb.* 70. *Sibth.* 64. *Fl. Dan.* t. 1025. *Mill. Illustr. t.* 11.

P. rotundifolium. *Raii Syn.* 148.

P. latifolium. *Ger. em.* 821.

Potamogetum. *Fuchs. Hist.* 651.

Angl. Broad-leaved Pond-weed.

In fluviis et stagnis frequens.

Perennis. Fl. Julio.

Radix e fibris longis simplicibus. Caules longissimi, teretes, foliosi. Folia inferiora submersa, linearia; superiora natantia, ovali-oblonga, nervosa, longius petiolata. Stipulæ intrafoliaceæ, lineares, acute, petiolo ferè æquantes. Pedunculi solitarii, bractea ad basin stipulis conformi, simplices, crassi, teretes. Spica simplex, ex aquis emersa. Flores fusco-virides.

*perfoli-
atum.*

2. POTAMOGETON foliis cordatis amplexicaulis omnibus immerfis.

Potamogeton perfoliatum. Sp. Pl. 182. Hudf. 74.

With. 212. Relb. 70. Sibth. 65. Engl.

Bot. t. 168. Fl. Dan. t. 196. Raii Syn. 149.

P. tertium Dodonæi. Ger. em. 822.

Angl. Perfoliate Pond-weed.

In fluviis, fossis et stagnis frequens.

Perennis. Fl. Julio, Augusto.

Herba tota, præter spicas, immersa. Caulis alternatim ramosus. Folia pellucida, fragilia, nervosa, glaberrima, cordato-ovata, sessilia, amplexicaulia, omnia ferè uniformia. Spicæ erectæ, solitariae, breves, fuscae. Petala inflexo-conniventia, basi distantia.

- densum.* 3. POTAMOGETON foliis ovatis acuminatis oppositis confertis, caule dichotomo, spicâ quadriflorâ.

Potamogeton densum. Sp. Pl. 182. Hudf. 75.

With. 212. Relb. 72. Sibth. 65. Engl. Bot.

t. 397.

P. seu Fontinalis media lucens. Raii Syn. 149.

Baub. Hist. v. 3. 769.

Angl. Close-leaved Pond-weed.

In fossis, stagnis, et rivulis lentè fluentibus.

Perennis. Fl. Junio.

Herba immersa. Caules dichotomi, densè foliati. Folia opposita, acuta, patentissima. Pedunculi e dichotomiâ caulis, solitarii, recurvi. Spica brevissima, quadriflora, subinde biflora, pallidè virens.

Variat magnitudine pro aquarum profunditate. *D. Good-enough.*

- lucens.* 4. POTAMOGETON foliis ovato-lanceolatis planis in petiolos attenuatis, spicâ multiflorâ coarctatâ.

Pota-

Potamogeton lucens. *Sp. Pl.* 183. *Huds.* 74.
With. 212. *Relb.* 71. *Sibth.* 65. *Engl. Bot.*
t. 376. *Fl. Dan. t.* 195.

P. aquis immersum, folio pellucido, lato, ob-
 longo, acuto. *Raii Syn.* 148.

P. longis acutis foliis. *Ger. em.* 822.

β. P. folio angustio, pellucido, ferè gramineo. *Raii*
Syn. 148?

Angl. Shining Pond-weed.

In fossis, fluviis, lacubus et stagnis frequens.

Perennis. Fl. Junio, Julio.

Herba immersa, spicâ tantùm exsertâ.

Variat foliis in petiolum dilatatum decurrentibus ut in
Fl. Dan., vel basi rotundatis, petiolo vix marginato,
 ut in *Engl. Bot.* delineatum videri est. Stipulæ intra-
 foliaceæ, amplexicaules, magnæ, sub pedunculo oppo-
 sitæ. Spica densa, multiflora, fusco-virens.

5. POTAMOGETON foliis lanceolatis alternis *crispum*,
 undulatis serratis.

Potamogeton crispum. *Sp. Pl.* 183. *Huds.* 75.
With. 213. *Relb.* 71. *Sibth.* 65. *Curt. Lond.*
fasc. 5. *t.* 15. *Fl. Dan. t.* 927.

P. seu Fontinalis crispa. *Raii Syn.* 149.

Tribulus aquaticus minor, quercûs floribus.
Ger. em. 824.

β. Potamogeton ferratum. *Huds.* 75.

Angl. Curled Pond-weed.

In stagnis, vel rivulis lentè fluentibus vulgaris.

Perennis. Fl. Junio, Julio.

Caulis angulatus. Folia immersa, obtusa, plerumquē
 alterna; in β superiora verò opposita. Pedunculi
 emerfi. Spica pauciflora, floribus rubicundis, remo-
 tiusculis.

6. POTAMOGETON folis linearibus obtusis, *compressum*,
 caule compresso.

Potamogeton compressum. *Sp. Pl.* 183. *Huds.* 75. *With.* 213. *Relb.* 72. *Engl. Bot.* t. 418. *Fl. Dan.* t. 203.

P. caule compresso, folio graminis canini. *Raii Syn.* 149.

Angl. Flat-stalked Pond-weed.

In rivulis lentè fluentibus, et fossis.

Perennis. Fl. Junio, Julio.

Herba immerfa. Caulis flexuosus, compressus. Foliæ pallidè lætèque virentia, alterna, (sub pedunculo, ut in aliis, opposita,) linearia, angusta, obtusa, integerrima. Stipulæ foliis latiores, complanatæ vel angulato-depressæ, sæpiùs obtusæ, et subinde bifidæ. Pedunculi complanati, emersi, floribus circiter quatuor vel sex, per paria distantibus.

gramineum. 7. POTAMOGETON foliis lineari-lanceolatis alternis sessilibus stipulâ latioribus, caule tereti subdichotomo.

Potamogeton gramineum. *Sp. Pl.* 184. *Huds.* 76. *With.* 214.

P. gramineum latiusculum, foliis et ramificationibus densissimè stipatis. *Dill. in Raii Syn.* 149. t. 4. f. 3.

Angl. Grassy Pond-weed.

In fossis et fluviis. *In Herb. D. Rose.*

Near Deptford. *Dillenius.*

Perennis. Fl. Julio.

Herba immerfa. Caulis gracilis, teres, flexuosus, ramissimus, subdichotomus, immersus. Folia alterna, patentia, approximata, linearia, obtusiuscula, vix serrulata. Stipulæ involuto-tubulosæ, et inde, inferiores saltem, foliis angustiores. Spicæ ovatæ, paucifloræ, densæ, solitariae; primordialis e dichotomiâ caulis. Pedunculi longitudine spicarum.

Fl. Dan. t. 222, icon à Lightfootio laudata, mihi videtur diversa species, cujus exemplaria in Britannia nondum inveni.

8. POTAMOGETON foliis linearibus oppositis *pufillum*.
alternisque stipulâ angustioribus basi patentibus, caule tereti, pedunculis axillaribus.

Potamogeton pufillum. *Sp. Pl.* 184. *Huds.* 77.

With. 214. *Relb.* 73. *Sibth.* 66. *Engl. Bot.*

t. 215.

P. pufillum, graminco folio, caule tereti. *Raii Syn.* 150.

P. pufillum, gramineo folio breviori. *Vaill. Paris. t.* 32. *f.* 4.

Angl. Small Pond-weed.

In fossis et stagnis, præcipuè argillaceis.

Pernnis. Fl. Julio.

Herba immerfa, spicæ sub florecentiâ emersæ, ut in aliis.

Caulis alternatim ramosus, tenuis, striatus. Folia linearia, angustissima, basi patentissima, sub spicis opposita. Stipulæ amplexicaules, foliis multò latiores. Pedunculi axillares, solitarii, spicâ longiores. Spica ovata, parva, densa, pauciflora.

9. POTAMOGETON foliis setaceis parallelis *pectinatum*.
approximatis distichis basi vaginantibus.

Potamogeton pectinatum. *Sp. Pl.* 183. *Huds.*

76. *With.* 213. *Relb.* 72. *Sibth.* 66. *Engl.*

Bot. t. 323. *Fl. Dan. t.* 186.

P. millefolium, seu foliis graminicis, ramosum.

Raii Syn. 150.

β. P. marinum. *Sp. Pl.* 184. *Huds.* 76. *With.* 213.

Relb. Suppl. 2. 9.

P. maritimum ramosissimum, grandiusculis capitulis, capillaceo folio, nostras. *Pluk. Phyt.*

t. 216. *f.* 5. *Dill. in Raii Syn.* 150.

P. ingens, graminco folio longiori. *Vaill. Pa-*

ris. t. 32. *f.* 3.

Angl. Fennel-leaved Pond-weed. β Sea Pond-weed.

In fluviis et stagnis. β In fossis aquâ marinâ imbutis.

Perennis. Fl. Julio.

Radix repens, primo ortu tuberosa. Herba immerfa, non nisi in aquis stagnantibus flores ferens. Caulis teres, ramosissimus, ramis alternis, approximatis. Folia approximata, disticha, setacea, basi vaginantia. Stipula folio adnata, margine membranacea, apice bifida, acuta. Spicæ exsertæ, interruptæ, sæpè longius pedunculatæ, floribus fasciculatis, inferioribus remotissimis.

Quibus notis *P. marinum* a pectinato differat, ut etiam vel varietas censeatur, me, neque sanè cum et alios doctissimos etiam fefellerit, pudet confiteri, fugit.

setaceum. † 10. POTAMOGETON foliis lanceolatis oppositis acuminatis.

Potamogeton setaceum. *Sp. Pl.* 184. *Huds.* 76. *With.* 214.

Angl. Setaceous Pond-weed.

In fossis turfosis in comitatu Lancastriensi. *Huds.*

Perennis. Fl. Julio, Augusto.

Hanc speciem nec in Herb. Linn. nec alibi vidi. Locus in Florâ Britannicâ ex fide Hudsoni pendet.

Dignoscenda foliis setaceis, oppositis.

68. RUPPIA. *G. Pl.* 68. *Juss.* 19. *Gærtn. t.* 84.

Cal. nullus. *Ccr.* nulla. *Semina* 4, pedicellata.

maritima. 1. RUPPIA.

Ruppia maritima. *Sp. Pl.* 184. *Huds.* 77. *With.* 215. *Engl. Bot. t.* 136. *Lightf.* 124. *t.* 8. *f.* 1.

Potamogeton maritimum, gramineis longioribus foliis, fructu ferè umbellato. *Raii Syn.* 134. *t.* 6. *f.* 1.

P. maritimum pusillum alterum. *Pluk. Phyt.* *t.* 248. *f.* 4.

Bucca-

Buccaferrea maritima, foliis acutissimis. *Mich.*

Gen. 72. *t.* 35.

Fucus ferulaceus. *Ger. em.* 1573.

Angl. Sea Ruppia. Taffel Pond-weed.

In fossis marinis salis, haud rara.

Perennis? Fl. Julio.

Habitus Potamogetonis pectinati vel marini, at folia vix disticha, nec stipulae apice productae, acutae, bifidae, sed retusae. Spica subquadriflora, emersa. Pedunculus longitudine valde varius. Germina ovata, post impregnationem pedicellis propriis elevata.

Valisneriae modo planta haec pedunculos convolutos ad aquae profunditatem laxat et contrahit. *D. Goodenough.*

69. SAGINA. *G. Pl.* 68. *Juss.* 300. *Gærtn.* *t.* 129.

Cal. tetraphyllus. *Petala* 4. *Caps.* unilocularis.

1. SAGINA caulibus procumbentibus glabris, petalis brevissimis. *procumbens.*

Sagina procumbens. *Sp. Pl.* 185. *Huds.* 73.

With. 215. *Relb.* 73. *Sibth.* 66. *Curt. Lond.*

fasc. 3. *t.* 12. *Arduin. Spec.* 2. 23. *t.* 8. *f.* 2.

Alfinella muscoso flore repens. *Raii Syn.* 345.

Angl. Procumbent Pearl-wort.

In arenosis, muris, arcis, et ambulacris hortorum neglectis frequens.

Perennis. Fl. Maio—Augusto.

Radix fibrosa. Caules plurimi, procumbentes, radican-
tes, foliosi, glabri. Folia linearia, mucronata, glabra,
connata; radicalia caespitosa. Pedunculi axillares, so-
litarii, foliis longiores, uniflori. Flores subnutantes,
herbaeci. Calyx patens, foliolis obtusis. Petala calyce
duplò breviora, integra. Capsula ovata.

2. SAGINA caulibus erectiusculis pubescentibus, *apetala*,
petalis obsoletis.

Sagina apetala. *Linn. Mant.* 2. 559. *With.* 216.
Sibth. 67. *Abbot.* 39. *Curt. Lond. fasc.* 5.
t. 14. *Dickf. H. Sicc. fasc.* 3. 6. *Arduin,*
Spec. 2. 22. *t.* 8. *f.* 1.

Saxifraga Anglica Alfinefolia annua. *Raii Syn.*
 345.

Angl. Annual Small-flowered Pearl-wort.

In arenosis, muris, arcisque vulgaris.

Annua. Fl. Maio, Junio.

Radix fibrosa, parva. Caules plurimi, pubescentes, erectiusculi, minimè procumbentes, nec radicales. Folia præcedentis. Pedunculi elongati, pilosi. Petala alba, emarginata, minutissima, sæpè abortiva. Semina lineâ nigrâ cincta, ut in *Spergula subulatâ*.

crecta.

3. *SAGINA* caule erecto subunifloro, foliolis calycinis acutis, petalis integris.

Sagina crecta. *Sp. Pl.* 185. *Huds.* 73. *With.*
 216. *Kelb.* 74. *Sibth.* 67. *Abbot.* 40.
Curt. Lond. fasc. 2. *t.* 12. *Engl. Bot. t.* 609.
Dickf. H. Sicc. fasc. 6. 6. *Dryandr. Cat. v.* 3.
 244.

Mœnchia quaternella. *Ehrh. Beitr. fasc.* 2. 177.

Alfinella foliis caryophylleis. *Raii Syn.* 344.
t. 15. *f.* 4.

Angl. Upright Pearl-wort.

In pascuis et ericetis glareosis sterilibus.

Annua. Fl. Maio.

Radix parva. Caules vel paucissimi erecti stricti, vel plurimi divaricati. Herba tota glauca, pulchella, glabra. Folia lanceolata, acuta. Pedunculi solitarii, longissimi, vix duo in caule. Flores albi. Foliola calycina lanceolata, acuta, margine scariosa. Petala calyce parum breviora, integra. Germen ovatum. Stigmata ferè sessilia, villosa. Capsula cylindrica, indivisa, octo vel decem-dentata, unilocularis, polysperma. Semina reniformia, scabra,

Fruetus,

Fruſtus, optimè monente Curtiſio, eſt omninò Ceraſtii, at petala integra, numeruſque floris partium, cum hoc genere nequaquam conveniunt; neque cum Ceraſtio, neque cum aliis Saginis, convenit habitus. Etiamnum in discrimine manet, fortè ſui generis herba.

70. TILLÆA. *G. Pl.* 68. *Juff.* 307. *Gærtn.*
t. 112.

Cal. 3, 4, 5-fidus. *Petala* 3, 4 vel 5. *Nectarium* nullum. *Capsulæ* 3, 4 vel 5, polyspermæ, introrsum dehiscentes.

Non niſi defectu nectarii a Crassulâ differt.—
Gærtner.

1. TILLÆA caulibus procumbentibus, floribus *muscosa*.
sessilibus plerumque trifidis.

Tillæa muscosa. Sp. Pl. 186. *Huds.* 73. *With.*
175. *Rose Elem. append.* 448. t. 2. f. 2.

T. muscosa, annua, perfoliata, flore albo. *Mich.*
Gen. 22. t. 20.

Sempervivum omnium minimum, repens, muscosum, Polygoni facie. *Bocc. Mus.* 36. t. 22.

Polygonum muscosum minimum. *Bocc. Sic.*
56. t. 29.

Crassula foliis sessilibus connatis, floribus aggregatis in foliorum alis. *Guett, Obs.* v. 2. 97.

Angl. Mossy Tillæa.

In ericetis sterilissimis arenosis.

Frequent in Norfolk and Suffolk.

Annua. Fl. Maio. Junio.

Radix fibrosa, parva. Herba succulenta, glabra, rubicunda. Folia perfoliata, obtusa, gibba, subinde terna. Flores subsolitarii, sessiles, plerumque trifidi, subinde quadrifidi, et etiam, monente Gærtnero, quinquedidi. Petala calyce minora, subulata. Capsulæ dispermæ,

71. RADIOLA. *Gmel. Syst.* v. 2. 289.

Linum. G. Pl. 153. *Juff.* 303.

Cal.

Cal. multifidus. *Petala* 4. *Capsula* supera, octovalvis, octolocularis. *Semina* solitaria.

millegrana- I. **RADIOLA.**

na.

Radiola linoides. *Gmel. Syst.* v. 2. 289.

R. vulgaris serpyllifolia. *Dill. in Raii Syn.* 345. t. 15. f. 3.

Linum Radiola. *Sp. Pl.* 402. *Syst. Nat. ed.* 12. v. 3. 225. *Huds.* 134. *With.* 323. *Lightf.* 174. *Dicks. H. Sicc. fasc.* 10. 7. *Fl. Dan.* t. 178.

Linocarpum Serpylli folio, multicaule et multiflorum. *Mich. Gen.* 23. t. 21.

Millegrana minima. *Raii Syn. ed.* 2. 207. *Ger. em.* 569.

Chamælinum vulgare. *Vaill. Paris.* t. 4. f. 6. *Angl. All-feed.*

In inundatis fabulosis.

Annua. Fl. Julio, Augusto.

Radix parva, fibrosa. *Caulis* uncialis vel biuncialis, dichotomus, ramosissimus, corymbosus, foliosus, teres, glabriusculus. *Folia* opposita, sessilia, ovata, integerrima, glabra. *Flores* e dichotomiâ caulis, pedunculati, solitarii, erecti, exigui, albi. *Calyx* monophyllus, semiquadrifidus, laciniis trifidis. *Petala* obovata. *Stamina* æqualia. *Styli* breves. *Stigmata* capitata. *Capsula* subrotunda, fructura Lini, sed valvulis loculamentisque tantum octo. *Semina* solitaria, elliptica.

Genus hocce, olim a Dillenio constitutum, recensioribus verò scriptoribus neglectum, nuper, nec ut mihi videtur inconsultò, instauravit Gmelinus; at ex sequentibus videri potest annon charactere generico, nomine specifico, et citatione Linnæi erraverat. *Calyx* enim multifidus, nec tetraphyllus est, quâ notâ a Lino præcipuè discrepat; habitus Lini non est, ut *linoides* dici possit; nec ut genus novum, *Radiolæ* sub nomine, a Linnæo unquam descriptum fuit.

Classis V.

PENTANDRIA.

Stamina 5.

MONOGYNIA.

Stylus 1.

* *Fl. monopetali, inferi, tetraspermi. Asperifoliæ.*

81. ECHIUM. *Cor. fauce nudâ, irregularis. Stigma bipartitum.*
76. PULMONARIA. *Cor. fauce nudâ, infundibuliformis. Cal. prismaticus.*
73. LITHOSPERMUM. *Cor. fauce nudâ, infundibuliformis. Cal. quinquepartitus.*
77. SYMPHYTUM. *Cor. fauce dentatâ, ventricosa.*
78. BORAGO. *Cor. fauce dentatâ, rotata.*
80. LYCOPSIS. *Cor. fauce fornicatâ, infundibuliformis, tubo curvato.*
79. ASPERUGO. *Cor. fauce fornicatâ, infundibuliformis. Fructus compressus.*
75. CYNOGLOSSUM. *Cor. fauce fornicatâ, infundibuliformis. Semina depressa, latere affixa.*
74. ANCHUSA. *Cor. fauce fornicatâ, infundibuliformis, basi prismaticâ. Sem. basi insculpta.*
72. MYOSOTIS. *Cor. fauce fornicatâ, hypocrateriformis, lobis submarginatis.*

* * *Fl. monopetali, inferi, angiospermi.*

87. ANAGALLIS. *Capsf.* unilocularis, circumscissa. *Cor.* rotata: *Stam.* hirsuta.
86. LYSIMACHIA. *Capsf.* unilocularis, decemvalvis. *Cor.* rotata.
83. CYCLAMEN. *Capsf.* unilocularis, intùs pulposa. *Cor.* rotata, reflexa. *Stigma* acutum.
82. PRIMULA. *Capsf.* unilocularis, ore decemfido. *Cor.* fauce pervia, tubo cylindrico. *Stigma* globosum.
85. HOTTONIA. *Capsf.* unilocularis. *Cor.* hypocrateriformis. *Stam.* fauce inposita. *Stigma* globosum.
84. MENYANTHES. *Capsf.* unilocularis. *Cor.* hirsuta. *Stigma* bifidum.
98. DATURA. *Capsf.* bilocularis, quadrivalvis, *Cor.* infundibuliformis. *Cal.* deciduus.
99. HYOSCYAMUS. *Capsf.* bilocularis, operculata. *Cor.* infundibuliformis. *Stigma* capitatum.
97. VERBASCUM. *Capsf.* bilocularis. *Cor.* rotata. *Stigma* simplex. *Stam.* declinata, barbata.
102. CHIRONIA. *Capsf.* bilocularis. *Cor.* hypocrateriformis. *Antheræ* demum spirales.
89. CONVULVULUS. *Capsf.* bi- vel tri-locularis. *Sem.* bina. *Cor.* campanulata, plicata. *Stigmata* duo.
90. POLEMONIUM. *Capsf.* trilocularis. *Cor.* quinquepartita. *Stam.* valvis inposita.
88. AZALEA. *Capsf.* quinquelocularis. *Cor.* campanulata. *Stigma* obtusum.
112. VINCA. *Folliculi* duo, erecti. *Cor.* hypocrateriformis,

101. SOLANUM. *Bacca* bilocularis. *Cor.* rotata. *Antheræ* biporosæ.
100. ATROPA. *Bacca* bilocularis. *Cor.* campanulata. *Stam.* distantia, incurvata.

*** *Fl. monopetali, superi.*

103. SAMOLUS. *Capsf.* unilocularis, apice quinquevalvis. *Cor.* hypocrateriformis.
93. JASIONE. *Capsf.* semibilocularis, apice dehiscens. *Cor.* rotata, quinquepartita. *Stigma* clavatum.
92. PHYTEUMA. *Capsf.* bi- five tri-locularis, perforata. *Cor.* rotata, quinquepartita. *Stigma* bi- five tri-fidum.
94. LOBELIA. *Capsf.* bi- five tri-locularis. *Cor.* irregularis, fissâ. *Stigma* simplex.
91. CAMPANULA. *Capsf.* tri- five quinquelocularis, perforata. *Cor.* campanulata. *Stigma* trifidum.
104. LONICERA. *Bacca* bilocularis, polysperma. *Cor.* irregularis.

Rubia.

**** *Fl. pentapetali, inferi.*

105. RHAMNUS. *Bacca* trilocularis. *Cal.* urceolatus, corollifer.
106. EUONYMUS. *Capsf.* quinquelocularis. *Sem.* calyptrata. *Cal.* planus.
95. IMPATIENS. *Capsf.* quinquelocularis, elastica. *Cal.* diphyllus. *Cor.* irregularis.
96. VIOLA. *Capsf.* unilocularis, trivalvis. *Cal.* pentaphyllus, calcaratus. *Cor.* irregularis.

***** *Fl.*

* * * * * *Fl. pentapetali, superi.*

107. RIBES. *Bacca* polysperma. *Cal.* corollifer. *Stylus* bifidus.
 108. HEDERA. *Bacca* pentasperma, calyce cincta. *Stylus* simplex.

* * * * * *Fl. incompleti.*

110. GLAUX. *Caps.* pentasperma, supera. *Cal.* coloratus, monophyllus.
 109. ILLECEBRUM. *Caps.* monosperma, supera. *Cal.* cartilagineus, pentaphyllus.
 111. THESIIUM. *Sem.* unicum, corticatum, inferum. *Cal.* staminifer.

Polygonum amphibium.

DIGYNIA.

Styli 2.

* *Fl. monopetali, inferi.*

119. SWERTIA. *Caps.* unilocularis. *Cor.* rotata, poris nectariferis ad basin laciniarum.
 120. GENTIANA. *Caps.* unilocularis. *Cor.* basi tubulosa, poris nectariferis destituta.
 118. CUSCUTA. *Caps.* bilocularis. *Cor.* campanulata.

* * *Fl. pentapetali, inferi.*

Staphylea pinnata.

* * * *Fl. incompleti.*

116. SALSOLA. *Sem.* unicum, cochleatum, tectum.

114. CHE-

114. CHENOPODIUM. *Sem.* unicum, lenticulare, superum.
115. BETA. *Sem.* unicum, reniforme, calyce carnosio immersum.
113. HERNIARIA. *Caps.* monosperma, corticata. *Filam.* quinque sterilia.
117. ULMUS. *Caps.* monosperma, membranacea, compressa.

**** *Fl. pentapetali, superi, dispermi. Umbellatæ.*

A. Involucro universali partialique.

121. ERYNGIUM. *Fl.* capitati. *Recept.* paleaceum.
122. HYDROCOTYLE. *Umbella* simplex. *Sem.* compressa.
123. SANICULA. *Umb.* subcapitatae, disco abortiente. *Sem.* uncinato-muricata.
135. HERACLEUM. *Fl.* radiati, abortivi. *Invol.* deciduum. *Sem.* membranacea.
141. CENANTHE. *Fl.* radiati; disci steriles. *Sem.* coronata, sessilia.
125. ECHINOPHORA. *Fl.* radiati, abortivi; centralis foemineus. *Sem.* involucello immersa.
127. CAUCALIS. *Fl.* radiati, abortivi. *Invol.* simplex. *Sem.* muricata.
128. DAUCUS. *Fl.* radiati, abortivi. *Invol.* pinnatum. *Sem.* muricata.
126. TORDYLIUM. *Fl.* radiati, fertiles. *Invol.* simplex. *Sem.* margine crenata.
142. CORIANDRUM. *Fl.* radiati, abortivi. *Involucella* dimidiata. *Frucl.* sphaericus.
133. PEUCEDANUM. *Fl.* flosculosi, abortivi. *Invol.* simplex. *Sem.* compressa, alata, striata.

130. **CONIUM.** *Fl.* flosc. fertiles. *Pet.* cordata.
Sem. gibba, costato-fulcata. *Involucella*
dimidiata.
129. **BUNUM.** *Fl.* flosc. fert. *Pet.* cordata.
Involucella setacea.
132. **ATHAMANTA.** *Fl.* flosc. fert. *Pet.* cor-
data. *Sem.* convexa, striata.
124. **BUPLEURUM.** *Fl.* flosc. fert. *Pet.* in-
voluta. *Sem.* compressa, striata.
139. **SIUM.** *Fl.* flosc. fert. *Pet.* cordata. *Sem.*
subovata, striata.
131. **SELINUM.** *Fl.* flosc. fert. *Pet.* cordata.
Sem. compressa, medio striata.
134. **CRITHMUM.** *Fl.* flosc. fert. *Pet.* basi
lata. *Invol.* horizontale. *Cal.* integr.
137. **LIGUSTICUM.** *Fl.* flosc. fert. *Pet.* in-
voluta. *Invol.* membranacea. *Cal.* quin-
quedentatus.
136. **MEUM.** *Fl.* flosc. fert. *Pet.* inflexa. *In-*
vol. incisa. *Cal.* obsoletus.
138. **ANGELICA.** *Fl.* flosc. fert. *Pet.* incurva.
Umbellulæ globosæ. *Sem.* hemisphærica,
trialata.
140. **SISON.** *Fl.* flosc. fert. *Pet.* inflexa. *Um-*
bella depauperata. *Sem.* ovata, striata.

B. *Involucris partialibus, universali nullo.*

145. **ÆTHUSA.** *Fl.* subradiantes, fertiles. *In-*
volucella dimidiata.
147. **CHÆROPHYLLUM.** *Fl.* subradiantes,
abortivi. *Involuc.* reflexa, concava, penta-
phylla.
146. **SCANDIX.** *Fl.* radiati, abortivi. *Fruct.*
subulatus.

143. **PHEL-**

143. PHELLANDRIUM. *Fl.* flosculosi, fertiles. *Fruet.* coronatus.
148. IMPERATORIA. *Fl.* flosc. fert. *Umb.* expanso plana. *Fruet.* compressus, marginatus.
144. CICUTA. *Fl.* flosc. fert. *Fruet.* subovatus, fulcatus.

Bupleurum rotundifolium. Caulis infesta. Angelica sylvestris. Sium nodiflorum. Oenanthe fistulosa et peucedanifolia.

C. *Involucro nullo; vix universalis, nunquam partialibus.*

154. APIUM. *Involucrum* monophyllum. *Fl.* flosc. fert. *Fruet.* costatus. *Pet.* inflexa.
153. PIMPINELLA. *Fl.* flosc. fert. *Fruet.* striatus. *Stigmata* subglobosa.
155. ÆGOPODIUM. *Fl.* flosc. fert. *Fruet.* ovatus, striatus. *Stigm.* simplicia.
151. ANETHUM. *Fl.* flosc. fert. *Fruet.* subcompressus, utrinque tricostratus.
149. PASTINACA. *Fl.* flosc. fert. *Fruet.* compresso-planus.
150. SMYRNIUM. *Fl.* flosc. abortivi. *Semina* reniformia, angulata.
152. CARUM. *Fl.* flosc. abort. *Sem.* gibba, striata.

TRIGYNIA.

Styli 3.

* *Flores superi.*

156. VIBURNUM. Cor. quinquefida. *Bacca*
monosperma.
157. SAMBUCUS. Cor. quinquefida. *Bacca*
trisperma.

* * *Fl. inferi.*

158. STAPHYLEA. Cor. pentapetala. *Caps.*
2 vel 3, inflatæ.
159. TAMARIX. Cor. pentapetala. *Caps.* tri-
valvis. *Sem.* papposa.
160. CORRIGIOLA. Cor. pentapetala. *Sem.*
unicum, triquetrum.

Stellaria media.

TETRAGYNIA.

Styli 4.

161. PARNASSIA. *Nectaria* ciliata fetis globi-
feris.

PENTA-

PENTAGYNIA.

Styli 5.

163. LINUM. *Cor.* pentapetala. *Caps.* decemlocularis.
164. SIBBALDIA. *Cor.* pentapetala. *Sem.* quinque, nuda. *Cal.* decemfidus.
162. STATICE. *Cor.* pentapetala. *Sem.* unicam, calyce infundibuliformi vestitum.

Cerastium semidecandrum. Spergula arvensis, pentandra, et subulata.

HEXAGYNIA.

Styli 6.

165. DROSERA. *Cor.* pentapetala. *Caps.* trivalvis, polysperma.

POLYGYNIA.

Styli plurimi.

166. MYOSURUS. *Pet.* quinque, unguibus tubulosis, nectariferis. *Sem.* nuda.

Ranunculus hederaceus.

V. PENTANDRIA.

I. MONOGYNIA.

72. MYOSOTIS. *G. Pl.* 73. *Juss.* 131.
Gærtn. t. 68.

Cor. hypocrateriformis, quinquefida, sub-emarginata : fauce clausâ fornicibus.

1. MYOSOTIS feminibus nudis, foliis elliptico-lanceolatis, racemis ebracteatis multifloris.
scorpioides. *Engl. Bot. t.* 480.

α. *Myosotis scorpioides arvensis.* *Sp. Pl.* 188.
Huds. 78. *Relb.* 75. *hirsuta.* *Ger. em.* 337.
Fl. Dan. t. 583.

M. arvensis. *Sibth.* 68. *With.* 225.

M. scorpioides hirsuta. *Raii Syn.* 229.

β. *M. scorpioides γ.* *Sp. Pl.* 189. *Huds.* 78. β.
Relb. 75.

M. scorpioides hirta minor. *Raii Syn.* 229.

Echium Scorpioides minus, flosculis luteis.
Baub. Prod. 119. *Pin.* 254.

γ. *M. scorpioides latifolia hirsuta.* *Dill. in Raii Syn.* 229. *t.* 9. *f.* 2.

δ. *M. scorpioides palustris.* *Sp. Pl.* 188. *Huds.* 78.
Curt. Lond. fasc. 3. *t.* 13. *Ger. em.* 337.
Raii Syn. 229.

M. palustris. *With.* 225. *Relb.* 76. *Sibth.* 68.
Angl. Mouse-ear Scorpion-grass.

In arvis aridis. β in arenosis et muris. γ in umbrosis. δ in fossis, fluviis, et ad scaturigines.

α, β, γ, *annuæ*, δ *perennis.* Fl. totâ æstate.

Planta polymorpha, magis vel minus hirsuta. Radix fibrosa. Caulis teres, alternatim ramosus, foliosus. Folia obtusa. Racemi simplices, terminales, reflexo-convoluti, multiflori, sub florescentiâ expansi et postea elongati. Pedicelli ebracteati. Semina glabra.

Var.

Var. α gaudet herbâ hirsutissimâ, floribus parvis cæruleis; β floribus exiguis, luteis; γ floribus majoribus, cæruleis, habitu toto laxiore; δ floribus majoribus, cæruleis, herbâ vegetiore, foliis glabrioribus. Flores expansi pulcherrimè incarnati sunt.

73. LITHOSPERMUM. *G. Pl.* 74. *Juss.* 130.
Gærtn. t. 67.

Cor. infundibuliformis; fauce perviâ, nudâ.

Cal. quinquepartitus.

1. LITHOSPERMUM seminibus lævibus, co-officina-
rollis calycem vix superantibus, foliis lanceo-
latis acutiusculis venosis. *le.*

Lithospermum officinale. *Sp. Pl.* 189. *Huds.* 79.

With. 226. *Relb.* 76. *Sibth.* 69. *Abbot.* 41.

Engl. Bot. t. 134. *Dicks. H. Sicc. fasc.* 12. 7.

L. minus. *Ger. em.* 609.

L. f. Miliun Solis. *Raii Syn.* 228.

Angl. Common Gromwell. *Gray mill.* Grey
Millet.

In aridis glareosis vel calcareis.

Perennis. Fl. Maio.

Herba scabra, rigens, ramosa. Folia lanceolata, venosa, integerrima. Spicæ terminales, simplices, recurvæ, foliosæ. Flores parvi, lutei, ore gibbis quinque, obtusis, circumdato. Semina ovata, margaritacea, nitida.

2. LITHOSPERMUM seminibus rugosis, corol-
lis vix calycem superantibus, foliis obtusis
aveniis. *arvense.*

Lithospermum arvense. *Sp. Pl.* 190. *Huds.* 79.

With. 226. *Relb.* 76. *Sibth.* 69. *Engl. Bot.*

t. 123. *Dicks. H. Sicc. fasc.* 12. 9. *Fl. Dan.*

t. 456.

Buglossum arvense annuum, *Lithospermi folio.*

Raii Syn. 227.

Anchusa degener, facie *Milii Solis*. *Ger. em.* 610.

Angl. Corn Gromwell. Bastard Alkanet.

In arvis haud infrequens.

Annua. Fl. Maio, Junio.

Radix rubra, tinctoria. Herba scabra. Folia lingulata, obtusa, uninervia, avenia. Flores lactei. Semina scabra, fusca.

*purpuro-
cærule-
um.*

3. **LITHOSPERMUM** seminibus lævibus, corollis calycem longè superantibus, foliis lanceolatis acutis aveniis.

Lithospermum purpuro-cæruleum. *Sp. Pl.* 190. *Huds.* 79. *With.* 226. *Engl. Bot.* t. 117. *Jacq. Austr.* t. 14.

L. majus Dodonæi, flore purpureo, semine *Anchusæ*. *Raii Syn.* 229.

L. majus. *Ger. em.* 609.

Angl. Creeping or Purple Gromwell.

In dumetis, solo cretaceo, rariùs.

On the top of a bushy hill on the north side of Denbigh, Wales; also near Taunton, Somersetshire. *Ray*. Near Greenhithe, Kent. *Mr. Latham*.

Perennis. Fl. Maio.

Radix lignosa. Caules steriles procumbentes, radican-tes; floriferi erecti. Folia lanceolata, apice attenuata, uninervia, avenia, scabra. Racemi gemini, breves, foliosi. Flores violacei.

74. **ANCHUSA**. *G. Pl.* 74. *Juss.* 131.

Buglossum. *Gertn.* t. 67.

Cor. infundibuliformis; fauce clausâ fornicibus, *Sem.* basi insculpta.

*efficina-
lis.*

1. **ANCHUSA** spicis imbricatis secundis, bracteis ovatis, foliis lanceolatis.

Anchusa

Anchusa officinalis. *Sp. Pl.* 191. *Fl. Dan.*
t. 572.

Buglossa. *Brunf. Herb.* v. 1. 112.

B. italica. *Trag. Hist.* 232.

B. longifolia. *Cord. Hist.* 135.

B. vulgaris. *Ger. em.* 798.

Buglossum vulgare. *Matth. Valgr.* v. 2. 528.

Echium italicum spinosum. *Fuchs. Ic.* 150.

Angl. Common Alkanet.

In ruderatis maritimis.

On the links near Hartley Pans, Northumberland.

Rev. Thomas Butt.

Perennis. Fl. Junio, Julio.

Radix fusiformis, extus nigra. Herba piloso-scabra.

Caulis bipedalis, erectus, angulatus, foliosus, subramosus, paniculatus. Folia lanceolata, acuta, vix repanda; superiora basi subovata. Racemi plerumque gemini, revoluti. Bractæ ovatæ, nec, ut in *A. angustifoliâ*, linearilanceolatæ. Flores purpurei, infundibuliformes.

2. *ANCHUSA* pedunculis axillaribus diphyllis *semper-virens,*
capitatis, foliis ovatis.

Anchusa sempervirens. *Sp. Pl.* 192. *Huds.* 80.

With. 227. *Relb.* 77. *Engl. Bot. t.* 45. *Dicks.*

II. Sicc. fasc. 8. 8.

Buglossum latifolium sempervirens. *Dill. in Raii*

Syn. 227.

Borago sempervirens. *Ger. em.* 797.

Angl. Evergreen Alkanet.

In ruderatis et ad vias.

About Norwich in several places. At Haddiscoe, Suffolk. *Mr. Woodward.* Near Birmingham, Worcester, and Sidmouth. *With.* On the ruins of Maes Glas monastery, Flintshire. *Mr. Bingley.*

Perennis. Fl. Maio, Junio.

Radix crassa, extus nigra. Herba sempervirens, hispida. Caules angulato-teretes, foliosi. Folia alterna, ovata. Pedunculæ axillares, solitarii, capitati, apice diphylly. Corollæ limbus cæruleus, planiusculus. Structura floris ad Myosotidem accedit.

75. CYNOGLOSSUM. *G. Pl.* 75. *Juss.* 131.
Gærtn. t. 67.

Cor. infundibuliformis ; fauce clausâ fornicibus.
Sem. depressâ, interiore tantum latere stylo affixa.

officina- 1. CYNOGLOSSUM flaminibus corollâ brevioribus, foliis lato-lanceolatis tomentosis sessilibus.
le.

Cynoglossum officinale. *Sp. Pl.* 192. *Huds.* 80.
With. 227. *Relb.* 77. *Sibth.* 69. *Curt. Lond. fasc.* 4. *t.* 16.

Cynoglossum. *Raii Syn.* 226.

C. majus vulgare. *Ger. em.* 804.

Angl. Common Hounds-tongue.

In ruderatis et ad vias passim.

Biennis. Fl. Junio.

Radix fusiformis. Caulis erectus, teres, villosus, foliosus. Folia sparsa, amplexicaulia, lanceolata, obtusiuscula, integerrima, fusco-viridia, utrinque mollissimè pubescentia. Racemi paniculati, terminales, reflexo-convoluti. Calyx villosus. Flores sordidè rubri. Tota planta odore murino foetens.

sylvaticum. 2. CYNOGLOSSUM flaminibus corollâ brevioribus, foliis spatulato-lanceolatis lucidis nudiusculis subtus scabris.

Cynoglossum sylvaticum. *Jacq. Coll. v.* 2. 77, (ex descr.) *Sibth.* 70. *With.* 228. *Symons Syn.* 51.

C. virginianum. *Syst. Veg. ed.* 13. 157. nec *Sp. Pl.*
C. offi-

C. officinale γ. *Sp. Pl.* 193. β. *Huds.* 80.

Cynoglossa folio virenti. *Raii Syn.* 226.

C. media altera virente folio, rubro flore, montana frigidarum regionum. *Column. Ecphr.* 176. t. 175.

Angl. Green leaved Hounds-tongue.

Ad vias in umbrosis rariùs.

In Essex, Worcester-shire, and Surrey. *Ray.*

Out of St. Benediçt's gates, Norwich. *Andr. Caldwell, Esq.*

Biennis. Fl. Junio.

Caulis pilosus, foliosus. Folia lætè-virentia, lucida, nec tomentosa, subtùs præcipuè scabra, basin versus subpilosa. Racemi sparsi. Calyx hirtus. Flores cæruleo-rufescentes. Herba vix fœtet.

76. PULMONARIA. *G. Pl.* 75. *Juss.* 130.

Cor. infundibuliformis, fauce perviâ. *Cal.* prismatico-pentagonus.

1. PULMONARIA calyce longitudine tubi, foliis ovatis hirsutis. *officinalis.*

Pulmonaria officinalis. *Sp. Pl.* 194. *Huds.* 81.

With. 228. *Abbot.* 42. *Engl. Bot. t.* 118.

Fl. Dan. t. 482. *Woodv. Suppl. t.* 212.

P. foliis Echii. *Raii Syn.* 226.

P. maculosa. *Ger. em.* 808.

Pulmonaria. *Tillandsf. Ic.* 62.

Angl. Common Lungwort,

In nemorosis rariùs.

In a wood by Holbury-house, in the New Forest, Hampshire. *Ray.* Cliff wood, six miles west of Darlington. *Mr. E. Robson.* Between Thurleigh and Milton-Ernys, Bedfordshire. *Abbot.*

Perennis. Fl. Maio,

Folla

Folia albo maculata, utrinque hirsuta; radicalia petiolata, ovata vel subcordata; caulina ovata, sessilia, amplexicaulia. Racemi gemini, breves, terminales. Flores cærulei, ante expansionem carnei. Calyx hirtus.

Pulmonaria angustifolia Britanniae dubia civis est. Folium e Walliâ, ab amicissimo D. Robson missum, ad *Symphytum tuberosum* potius pertinere olim mihi visum est.

maritima.

2. **PULMONARIA** calyce abbreviato, foliis ovatis glaucis, caule ramoso procumbente.

Pulmonaria maritima. *Sp. Pl.* 195. *Huds.* 81. *With.* 229. *Lightf.* 134. *t.* 7. *Engl. Bot.* *t.* 368. *Curt. Lond. fasc.* 6. *t.* 18. *Fl. Dan.* *t.* 25. *Dicks. H. Sicc. fasc.* 1. 3.

Echium marinum. *Raii Syn.* 228.

Cerinthe maritima procumbens, foliis et floribus cæruleis. *Dill. Hort. Elth.* 75. *t.* 65.

Sea Bugloss. *Pet. Herb. t.* 29. *f.* 3.

Angl. Sea Lungwort.

In maritimis arenosis Scotiae atque Angliae borealis.

Perennis. *Fl.* Julio.

Radix lignosa, nigra. Caules procumbentes, ramosissimi. Folia ovata, basi attenuata, glauca, tuberculis minutis scabriuscula. Racemi foliosi, terminales. Flores purpureo-cærulei, amœni. Calyx ad basin ferè divinus, tubo brevior.

77. **SYMPHYTUM.** *G. Pl.* 76. *Juss.* 131. *Gertn. t.* 67.

Cor. limbus tubulato-ventricosus, fauce clausâ radiis subulatis. *Cal.* quinquepartitus.

officinale.

1. **SYMPHYTUM** foliis ovato-lanceolatis decurrentibus.

Symphytum officinale. *Sp. Pl.* 195. *Huds.* 81. *With.* 230. *Relb.* 78. *Sibth.* 70. *Curt. Lond. fasc.* 4. *t.* 18. *Fl. Dan. t.* 664.

S. magnum. *Raii Syn.* 230,

Consolida major. *Ger. em.* 806.

β. *Symphytum patens.* *Sibth.* 70.

Angl. Common Comfrey.

In aquosis, ad rivos et fossas.

Perennis. Fl. Maio, Junio.

Radix carnosa, mucilaginosa, extus nigra. Caulis bi-
vel tri-pedalis, erectus, foliosus, alatus, pilis deflexis
hispidus. Folia scabra, decurrentia, ovato-lanceolata,
radicalia petiolata, latiora. Racemi gemini, hispidi,
basi dichotomi. Flores luteo-albidi;—in β “rubri,
calycibus patentibus.” *Sibth.*

2. SYMPHYTUM foliis ovatis semi-decurrenti-*tubero-*
bus: summis oppositis. *sum.*

Symphytum tuberosum. *Sp. Pl.* 195. *Huds.* 648.

Lightf. 1091. *With.* 230. *Jacq. Obs. fasc.* 3.

12. t. 63. *Ger. em.* 806.

Angl. Tuberous-rooted Comfrey.

In umbrosis humidis.

Opposite the new well at the Water of Leith, but
more plentifully in Dr. Robertson's walks at
North Marchiston, near Edinburgh. *Mr. Yalden.*

Perennis. Fl. Julio.

Præcedente humilior. Radix extus alba. Folia caulina
ovata, petiolis alatis, vix decurrentia. Flores luteoli,
radiis a basi ad apicem denticulatis, ut in priore.

78. BORAGO. *G. Pl.* 77. *Juss.* 131.

Cor rotata, faucē radiis clausâ.

* 1. BORAGO foliis omnibus alternis, calycibus *officina-*
patentibus. *lis.*

Borago officinalis. *Sp. Pl.* 197. *Huds.* 82.

With. 230. *Relb.* 78. *Sibth.* 71. *Engl. Bot.*

t. 36.

B. hortensis. *Raii Syn.* 228. *Ger. em.* 797.

Buglossa

Buglossa urbana. *Cord. Hist.* 109. 2.

Angl. Common Borage.

In rudratis et ad vias haud rara, nec tamen indigena.

Biennis. Fl. Junio, Julio.

Herba mucilaginosâ, valdè hispida. Folia ovata, amplexicaulia, denticulato-repanda. Flores formosi, cyanei, staminibus radiisque nigris.

79. ASPERUGO. *G. Pl.* 77. *Juss.* 131.

Cal. fructûs compressus, lamellis plano-parallelis, sinuatis.

procumbens. 1. ASPERUGO calycibus fructûs complanatis. Asperugo procumbens. *Sp. Pl.* 198. *Huds.* 82.

With. 231. *Relb.* 79. *Fl. Dan.* t. 552.

A. vulgaris. *Raii Syn.* 228.

Aparine major Plinii. *Ger. em.* 1122.

Angl. German Madwort. *Pet. H. Brit.* t. 29. f. 12.

In rudratis solo pinguiore.

By Boxley in Suffex, and in Holy Island. *Ray.*
Near Purfleet, Effex, Mr. Alchorne. *Huds.* About
the church at Newmarket. *Relban.*

Annua. Fl. Aprili, Maio.

Radix parva, attenuata. Caules procumbentes, flagelliformes, scabri, foliosi. Folia opposita, adscendentia, oblonga, hispida. Flores axillares, solitarii, pedunculati, parvi, cærulei. Calyces post florescentiam aucti, compressi, clausi, reticulati, margine setosi, femina occultantes.

80. LYCOPSIS. *G. Pl.* 78. *Juss.* 131.

Gærtn. t. 67.

Cor. tubo incurvato, fauce clausâ squamis convexis.

1. **LYCOPSIS** foliis lanceolatis hispidis, calycibus *arvensis*.
florescentibus erectis.

Lycopsis arvensis. *Sp. Pl.* 199. *Huds.* 82.

With. 231. *Relb.* 79. *Sibth.* 71. *Curt. Lond.*

fasc. 5. *t.* 17. *Fl. Dan.* *t.* 435.

Buglossa sylvestris minor. *Raii Syn.* 227. *Ger.*
em. 799.

Angl. Small Bugloss.

In arvis et ad vias passim.

Annua. Fl. Junio, Julio.

Herba tota hispida. Caulis teres, erectus, ramosus. Folia cordato-oblonga, amplexicaulia. Racemi gemini, terminales, foliosi. Flores parvi, lætè cyanei, umbilico niveo. Semina angulata, rugosa, tuberculata.

31. **ECHIUM**. *G. Pl.* 78. *Juss.* 130. *Gærtn.* *t.* 67.

Cor. irregularis, fauce nudâ. *Stigma* bipartitum.

1. **ECHIUM** caule erecto piloso, spicis lateralibus *italicum*?
hirsutissimis, erecto-patentibus, corollis sub-
æqualibus, staminibus longissimis.

Echium italicum. *Sp. Pl.* 200. *Huds.* 83. *With.*
232. *Dicks. H. Sicc. fasc.* 14. 8.

E. flore albo. *Camer. Epit.* 738.

Lycopsis. *Raii Syn.* 227.

Angl. White Viper's Bugloss.

In arenosis insulæ Jerseianæ.

On the sandy grounds near St. Hilary plentifully.

Ray.

Biennis? Fl. Julio.

Caules virgati, hispidi, paniculato-ramosi. Spicæ parvæ, erecto-patentes, compactæ, post florescentiam elongatæ. Folia lingulata, tuberculata, hispida. Calyx fetis validis, densissimis, sæpè flavicantibus, muricato-hispidus. Corolla albida, tubulosa, parùm irregularis. Stamina longissimè exserta.

2. ECHIUM

vulgare. 2. **ECHIUM** caule tuberculato hispido, foliis caulinis lanceolatis hispidis, spicis lateralibus pilosis deflexis.

Echium vulgare. *Sp. Pl.* 200. *Huds.* 83. *With.* 232. *Relb.* 80. *Sibth.* 71. *Engl. Bot. t.* 181. *Mart. Fl. Rust. t.* 136. *Fl. Dan. t.* 445. *Raii Syn.* 227. *Ger. em.* 802.

β. *Huds.* 83.

E. alterum five *Lycopsis anglica*. *Dill. in Raii Syn.* 228.

E. violaceum. *With.* 233 ? nec Linn.

Lycopsis anglica. *Ger. em.* 802.

Angl. Common Blue Viper's Bugloss.

In arvis et ruderalis vulgaris.

Biennis. Fl. Junio, Julio.

Caulis tuberculis callosis, setiferis, hispidus, erectus, teres; in β ramosior, patens et sæpè decumbens. Spicæ pyramidatæ, inapertis adhuc floribus deflexæ, exinde sursum elongatæ. Folia tuberculata, hispida. Calyx minùs quam in priore muricatus. Corolla supernè ampliata, cærulea, ante explicationem rosea. Stamina longitudine ferè corollæ; in β longissimè exserta.

82. **PRIMULA**. *G. Pl.* 80. *Juss.* 96.

Gærtn. t. 50.

Capf. unilocularis, ore decemfido. *Corollæ* tubus cylindricus, fauce perviâ. *Stigma* globosum.

vulgaris. 1. **PRIMULA** foliis dentatis rugosis, scapis unifloris, corollæ limbo plano.

Primula vulgaris. *Huds.* 83. *With.* 233. *Sym. Syn.* 53. *Relb.* 81. *Engl. Bot. t.* 4.

P. acaulis. *Curt. Lond. fasc.* 6. t. 16.

P. veris γ. *Sp. Pl.* 205. *Fl. Dan. t.* 194.

P. veris vulgaris. *Raii Syn.* 284.

P. veris minor. *Ger. em.* 781.

Angl. Common Primrose.

In nemorosis, sepibus et dumetis ubique.

Perennis. Fl. Aprili.

Folia obovato-oblonga. Scapi numerosi, uniflori, longitudine foliorum, bracteis subulatis ad basin. Flores erecti, pallidè sulphurei, suaveolentes. Corollæ limbus quinquefidus, laciniis obcordatis, planis. Variat floribus rubicundis.

2. PRIMULA foliis dentatis rugosis medio con-*elator.*
tractis, scapo multifloro, corollæ limbo plano.

Primula elator. *With.* 234. *Sibth.* 72. *Syn.*

Syn. 53. *Engl. Bot. t.* 513. *Abbot.* 44. *Cul-*
lum. 73.

P. veris β. *Sp. Pl.* 204.

P. vulgaris β. *Huds.* 84.

P. inodora. *Relb.* 81.

P. pratensis inodorata lutea. *Raii Syn.* 284.

Angl. Oxlip. Great Cowslip.

In pascuis, dumetis, et nemorosis rariùs.

Perennis. Fl. Aprili.

Folia medium versus contracta ferè ut in sequente. Scapi pauci, erecti, foliis longiores, multiflori. Flores umbellati, pedicellati, exteriores sæpiùs nutantes, præcedenti formâ similes, sed parùm minores, corollæ limbo plano. Hybrida proles videtur e matre *Primulâ vulgaris*, patre *P. veris*.

3. PRIMULA foliis dentatis rugosis medio con-*veris.*
tractis, scapo multifloro, corollæ limbo con-
cavo.

Primula veris. *Huds.* 84. *Relb.* 81. *Engl. Bot.*
t. 5. α *Sp. Pl.* 204. *Fl. Dan. t.* 433.

P. officinalis. *With.* 234. *Curt. Lond. fasc.* 6.
t. 15. *Sibth.* 72.

P. veris major. *Raii Syn.* 284. *Ger. em.* 780.

Angl. Common Cowslip, or Paigele.

In pratis et pascuis, præcipuè solo argillaceo vel calcareo, vulgaris.

Perennis. Fl. Aprili.

Folia obovato-oblonga, medium versus repentè contracta; vel potiùs ovata, petiolata, petiolo alato. Scapi pauci, multiflori, foliis longiores, erecti. Flores umbellati, pedicellati, nutantes, suavcolentes, citrini, subtùs pallidiores. Corollæ limbus parvus, concavus.

farinosa. 4. PRIMULA foliis crenatis glabris subtùs pulverulentis, corollæ limbo plano.

Primula farinosa. *Sp. Pl.* 205. *Huds.* 84.

With. 235. *Engl. Bot. t.* 6. *Curt. Lond.*

fasc. 6. *t.* 14. *Dicks. H. Sicc. fasc.* 1. 4. *Fl.*

Dan. t. 125.

P. veris flore rubro. *Ger. em.* 783.

Verbasculum umbellatum alpinum minus. *Raii*

Syn. 285.

Angl. Bird's-eye Primrose.

In paludibus montosis copiosè.

Frequent in Westmoreland and the north-west part of Yorkshire.

Perennis. Fl. Junio, Julio.

Folia obovato-lanceolata, lætè viridia, lævia, glabra; subtùs venosa, pulverulento-alba. Scapus folia longè superans, erectus, strictus. Umbella multiflora, erecta. Corollæ limbus parvus, planus, ruber; subtùs pallidus; laciniis lineari-obcordatis.

83. CYCLAMEN. *G. Pl.* 82. *Juss.* 97.

Cor. rotata, reflexa; tubo brevissimo; fauce prominente. *Bacca* tecta capsulâ.

europæum. * 1. CYCLAMEN corollâ retroflexâ, foliis cordatis angulatis denticulatis.

Cyclamen europæum. *Sp. Pl.* 207. *With.* 235.

Sym. Syn. 53. *Engl. Bot. t.* 548.

C. ver-

C. vernum. *Ger. em.* 843.

Angl. Spring Cyclamen, or Sow-bread.

In nemorosis et dumetis, vix indigena.

On a bank in the parish of Bramfield, Suffolk, apparently wild. *D. E. Davy, Esq.*

Perennis. Fl. Aprili.

Radix tuber subrotundus, magnus, fibras plurimas, ramosas, exserens. Folia radicalia, longius petiolata, cordato-subrotunda vel oblonga, acuta, angulato-repanda, crenata vel denticulata, venosa, glabra; suprà maculis sinuosis albis picta; subtùs purpurascens. Petioli teretes, flexuosi, basi attenuati. Scapi petiolis conformes, sed altiores, uniflori, nudi. Flores nutantes, albi, ore purpurascens, corollæ laciniis erectis, tortis. Stamina brevissima, inclusa. Antheræ subulatæ, croceæ. Germen superum. Stylus subulatus. Stigma acutum. Scapi germine impregnato fœti spiritaliter contorquentur, fructumque in terrâ recondunt.

84. MENYANTHES. *G. Pl.* 82. *Juss.* 98.

Gærtn. t. 114.

Cor. hirsuta. *Stigma* bifidum. *Capsula* unilocularis.

1. MENYANTHES foliis ternatis, corollâ *super-* *trifoliata.*
pernè villosissimâ.

Menyanthes trifoliata. *Sp. Pl.* 208. *Huds.* 85.

With. 236. *Relb.* 82. *Sibth.* 73. *Abbot.* 44.

Curt. Lond. fasc. 4. *t.* 17. *Engl. Bot. t.* 495.

Fl. Dan, t. 541. *Woodv. Med. Bot. t.* 2.

M. palustre triphyllum latifolium et angustifolium. *Raii Syn.* 285.

Trifolium paludosum. *Ger. em.* 1194. *herba.*
Pharm. Lond.

Angl. Buckbean. Marsh Trefoil.

In aquosis, et palustribus spongiosis frequens.

Perennis. Fl. Junio, Julio.

Radix teres, flagelliformis, nigra. Caulis foliosus, hori-
Vol. I. Q zontaliter

zontaliter patens, ramosus. Folia ternata, petiolata, glabra, dentato-repanda. Flores spicati, bracteolati, pedicellati, incarnati, formosi, suprâ fibris densissimis, carnosis, obtusis, villosi. Stylus exertus. Herba amara, fudorem ciens.

nymphæ- 2. MENYANTHES foliis cordatis repandis, co-
aides. rollâ ciliatâ.

Menyanthes nymphæoides. *Sp. Pl.* 207. *Huds.* 85. *With.* 235. *Relb.* 82. *Sibth.* 73. *Engl. Bot. t.* 217. *Fl. Dan. t.* 339.

Nymphæa lutea minor, flore fimbriato. *Raii Syn.* 368.

Waldschmidia nymphoides. *Wigg. Fl. Hol- sat.* 20.

Angl. Fringed Buckbean. Lesser Yellow Water-lily.

In fluviis et stagnis majoribus, minùs frequens.

In the Thames at Ankerwyck near Windfor. *Rev. Dr. Goodenough.* Near Walton bridge. *Lord Viscount Lewisham.* Lake at Castle Howard, Yorkshire. *Mr. Teesdale.*

Perennis. Fl. Augusto.

Caules longissimi, teretes, apice foliosi atque floriferi. Folia opposita, simplicia, petiolata, natantia, cordato-subrotunda, repanda, glabra. Flores axillares, conferti, pedunculati, foli meridiano expansi, aurei, margine fimbriati, limbo supernè glabri. Stylus brevissimus. Stigma quinquefidum. Capsula compressa.

85. HOTTONIA. *G. Pl.* 82. *Juss.* 95.

Cor. hypocrateriformis. *Stam.* fauce inserta, lobis opposita. *Caps.* unilocularis. *Stigma* globosum. *Cal.* quinquepartitus.

palustris. 1. HOTTONIA caule multifloro, pedunculis verticillatis.

Hottonia palustris. *Sp. Pl.* 208. *Huds.* 85.
With. 236. *Relb.* 83. *Sibth.* 73. *Abbot.* 45.
Curt. Lond. fasc. 1. *t.* 11. *Dickf. H. Sicc.*
fasc. 7. 7. *Fl. Dan. t.* 487. *Engl. Bot.*
t. 364.

Hottonia. *Raii Syn.* 285.

Viola palustris. *Ger. em.* 826.

Angl. Featherfoil. Water Violet.

In fossis et stagnis purioribus, rivulisque tardè fluentibus.

Perennis. Fl. Junio, Julio.

Radix farmentosa. Caules erecti, nudi, multiflori, basi foliosi. Folia pectinato-pinnatifida, integerrima, glabra, immersa. Thyrsus erectus, verticillatus, longè supra aquam elatus. Pedicelli uniflori, basi bracteati. Flores incarnati, elegantes, laciniis sæpè 6, 7 vel 8, cum totidem staminibus. Capsula globosa.

Varietatem pulcherrimam, floribus saturatè roseis, invenit D. Hanbury, Soc. Linn. Socius.

86. *LYSIMACHIA.* *G. Pl.* 83. *Juss.* 95.

Gærtn. t. 50.

Cor. rotata. - *Caps.* globosa, mucronata, decemvalvis.

* *Pedunculis multifloris.*

1. *LYSIMACHIA paniculata*, racemis termina-*vulgaris!* libus, foliis ovato-lanceolatis acutis.

Lysimachia vulgaris. *Sp. Pl.* 209. *Huds.* 86.

With. 237. *Relb.* 84. *Sibth.* 74. *Curt. Lond.*
fasc. 5. *t.* 19. *Fl. Dan. t.* 689.

L. lutea. *Raii Syn.* 282. *Ger. em.* 474.

Angl. Yellow Loosestrife.

In aquosis, præsertim umbrosis, et ad rivos.

Perennis. Fl. Julio.

Radix repens. Caules erecti, tripedales, foliosi, apice
Q 2 pani-

paniculati, multiflori. Folia brevius petiolata, opposita, sæpè terna vel quaterna, patentia, ovato-lanceolata, acuta, repanda, venosa, glabra, subtus quandoque tomentosa. Racemi paniculati, erecti; pedicelli sparsi, graciles, teretes, bracteis setaceis solitariis ad basin. Calycis laciniae lanceolatae, acuminatae, fubeiliatae. Corolla aurea, patens. Stamina basi coalita, glabra.

thyrsiflora.

2. **LYSIMACHIA** racemis lateralibus pedunculatis.

Lysimachia thyrsiflora. *Sp. Pl.* 209. *Huds.* 86. *With.* 237. *Engl. Bot. t.* 176. *Fl. Dan.* t. 517.

L. lutea, flore globoso. *Raii Syn.* 283. *Ger. em.* 475.

Angl. Tufted Loosestrife.

In aquosis rarissimè.

In the east riding of Yorkshire; by King's Langley in Hertfordshire; and in Anglesea. *R. Syn.* Formerly found by Dr. White in a bog by Severus's hills near York, which is now drained, and the plant destroyed.—On the edge of a lake to the north of Glasgow; by the side of Loch Lomond; in a bog close by the town of Forfar; and in great plenty by the side of a lake about four miles east of Forfar. *Mr. John Mackay.*

Perennis. *Fl. Julio.*

Radix repens. Caules erecti, sesquipediales, simplicissimi, teretes, foliosi, glabri, subinde lanati. Folia opposita, sessilia, linear-lanceolata, acuta, integerrima, glabra. Racemi axillares, erecti, solitarii, oppositi, pedunculati, densi, subramosi. Flores parvi, pedicellati, bracteolati, flavi, rubro punctati, corollae laciniis lanceolatis, basi distantibus dente interjecto. Stamina corollâ paulò longiora.

* * *Pedunculis unifloris.*

nemorum.

3. **LYSIMACHIA** foliis ovatis acutis, floribus solitariis,

folitariis, caule procumbente, flaminibus lævibus.

Lyfimachia nemorum. Sp. Pl. 211. Hudf. 86.

With. 237. Relb. Suppl. 3. 1. Sibth. 74.

Abbot. 45. Curt. Lond. fasc. 5. t. 18. Fl.

Dan. t. 174. Engl. Bot. t. 527.

Anagallis lutea. Raii Syn. 282. Ger. em. 618.

Angl. Wood Loofestriſe. Yellow Pimpernel.

In nemoribus et umbrosis humidis.

Perennis. Fl. Maio—Septembr.

Caules procumbentes, radicanſes, ramoſi, tetragoni, rubicundi, nitidi, folioſi. Folia oppoſita, petiolata, ovata, integerrima, glabra. Pedunculi axillares, folitarii, uniflori, graciles. Flores tenelli, aurei. Stamina glaberrima, corollâ breviora. Calycis laciniaë ſubulataë.

4. *LYSIMACHIA* foliis ſubcordatis, floribus foli- *Nummularia.*
tariis, caule repente, flaminibus glanduloſis.

Lyfimachia Nummularia. Sp. Pl. 211. Hudf.

87. With. 238. Relb. 84. Sibth. 74. Curt.

Lond. fasc. 3. t. 14. Dickſ. H. Sicc. fasc. 12.

10. Fl. Dan. t. 493. Engl. Bot. t. 528.

Nummularia. Raii Syn. 283. Ger. em. 630.

Angl. Creeping Loofestriſe. Money-wort, or Herb Twopence.

In pratis et paſcuſ humidis, et umbroſis.

Perennis. Fl. Junio, Julio.

Caules proſtrati, repentes, quadranguli, compreſſi, plerumque ſimplices. Folia oppoſita, petiolata, cordatoſubrotunda, repanda, pallidè viridia. Pedunculi axillares, foliis breviores. Calycis laciniaë ovataë. Corolla citrina, oculo armato glandulis pedicellatis teſta et ciliata. Stamina etiam glandulis veſtita.

87. *ANAGALLIS. G. Pl. 83. Juſſ. 95.*

Gærtn. t. 50.

Cor. rotata. Capſ. circumſciſſa. Stam. hiriſuta.

arvensis. 1. *ANAGALLIS* foliis ovatis subtùs punctatis, caule procumbente.

Anagallis arvensis. *Sp. Pl.* 211. *Huds.* 87.

With. 238. *Relb.* 85. *Sibth.* 74. *Curt. Lond.*

fasc. 1. t. 12. *Dicks.* II. *Sicc. fasc.* 12. 11.

Fl. Dan. t. 88. *Engl. Bot.* t. 529.

A. flore phœniceo. *Raii Syn.* 282.

A. mas. *Ger. em.* 617.

β. *A. phœnicea*, foliis amplioribus ex adverso quaternis. *Raii Syn.* 282.

γ. *A. foemina.* *Raii Syn.* 282. *Ger. em.* 617.

A. cærulea. *Abbot.* 46.

δ. *A. terrestris* flore albo. *Raii Syn.* 282.

Angl. Scarlet Pimpernel. γ Blue Pimpernel.

In arvis et hortis frequens.

γ. Between Stockwell and Camberwell. *Huds.*

Worcestershire. *Nasb.* Devonshire and near

Bath, Mr. Martyn. *With.*

δ. In Cowley field. *Bobart.*

Annua. Fl. Junio, Julio.

Radix parva. Caulis procumbens, basi ramosissimus, tetragonus, glaber, foliosus. Folia opposita, (in β e solo pingui quaterna,) sessilia, ovata, integerrima, glabra, subtùs purpureo punctata. Pedunculi axillares, solitarii, uniflori, foliis longiores. Laciniae calycis lanceolatae, acuminatae, margine membranaceae. Corolla profundè quinquesida, minutè ciliata, plerumque punicea, ore violaceo. Stamina hirsuta, violacea. Antherae flavæ. Stigma obtusum, crenatum. Capsula globosa, diaphana. Semina angulosa, congesta.

Flores in γ cærulei, sed vix aliam inveni differentiam.

Forma calycis in omnibus nostris eadem est.

Corolla dum fudum non est arcè complicatur.

tenella. 2. *ANAGALLIS* foliis subrotundis acutiusculis petiolatis, caule repente, stigmate acuto.

Anagallis tenella. *Syst. Veg. ed.* 14. 196. *With.*

239. *Relb.* 86. *Sibth.* 75. *Abbot.* 46. *Curt. Lond. fasc.* 3. *t.* 15. *Engl. Bot. t.* 530. *Cullum.* 76.

Lyfimachia tenella. *Sp. Pl.* 211. *Huds.* 87. *Dicksf. H. Sicc. fasc.* 2. 12.

Nummularia minor, flore purpurascente. *Raii Syn.* 283. *Ger. em.* 630.

Angl. Bog Pimpernel.

In ericetis inundatis, et uliginosis.

Perennis. Fl. Julio, Augusto.

Caules prostrati, repentes, tenues, angulati, glabri, foliosi. Folia opposita, petiolata, patentia, cordato-subrotunda, parva, integerrima, glabra, lætè viridia, Pedunculi axillares, solitarii, erecti, foliis duplò vel triplò longiores, uniflori. Calycis laciniaè lanceolatae, acutae. Corolla rosea, venulis saturationibus picta, basi tubulosâ, limbo erecto-patente. Stamina villis articulatis vestita. Stylus subulatus. Stigma simplex, acutum. Capsula circumscissa.

Habitus *Lyfimachia*è, corolla ferè *Centunculi*, stamina verò fructusque *Anagallidis* sunt.

88. AZALEA. *G. Pl.* 85. *Juss.* 158.

Gærtn. t. 63.

Cor. campanulata. *Stam.* receptaculo inserta.

Caps. quinquelocularis.

1. AZALEA ramis diffuso-procumbentibus, foliis *procumbens.* oppositis revolutis glaberrimis.

Azalea procumbens. *Sp. Pl.* 215. *Fl. Lapp.*

ed. 2. 60. *t.* 6. *f.* 2. *Huds.* 88. *With.* 239.

Fl. Dan. t. 9. *Lightf.* 139. *Dicksf. H. Sicc. fasc.* 12. 12.

Chamæcistus serpyllifolius. *Ger. em.* 1284.

Angl. Trailing Azalea.

In ericetis clatis alpinis.

On most of the high mountains of Scotland.

Frutex. Fl. Julio.

Caules lignosi, procumbentes, implexi, ramosissimi. Rami foliosi, teretes, glabri. Folia opposita, petiolata, patentissima, elliptica, obtusa, revoluta, integerrima, glabra. Petioli canaliculati, ciliati. Pedunculi terminales, plerumque gemini, uniflori, rubicundi, basi bracteati. Flores erecti, saturatè rosei, regulares, campanulati, limbo quinquesidi. Stamina inclusa, æqualia, nec declinata. Capsula subrotundo-ovata, acuta, quinquelocularis, valvularum marginibus inflexis.

89. CONVULVULUS. *G. Pl.* 86. *Juss.* 133.
Gærtn. t. 134.

Cor. campanulata, plicata. *Stigmata* duo. *Caps.* bi- vel tri-ocularis; loculis dispermis.

arvensis. 1. CONVULVULUS foliis sagittatis utrinque acutis, pedunculis subunifloris, bracteis minutis a flore remotis.

Convolvulus arvensis. *Sp. Pl.* 218. *Huds.* 88.
With. 239. *Relb.* 86. *Sibth.* 76. *Curt.*
Lond. fasc. 2. *t.* 13. *Fl. Dan. t.* 459. *Engl.*
Bot. t. 312. *Mart. Fl. Rust. t.* 89. *Dicks.*
H. Sicc. fasc. 14. 11.

C. minor vulgaris. *Raii Syn.* 275.

Smilax lenis minor. *Ger. em.* 861.

β. *Convolvulus arvensis minimus.* *Raii Syn.* 276.
Pluk. Phyt. t. 24. *f.* 3.

γ. *C. flore minimo ad ungucem ferè secto.* *Dill. in*
Raii Syn. 276.

Angl. Small Bindweed.

In arvis et ad scpes vulgaris, hortorum vitium summum, præcipuè solo glareoso.

Perennis. Fl. Junio, Julio.

Radix repens, profundè descendens, vivacissima, vix extricanda. Caules volubiles, angulati. Folia alterna, petiolata, sagittata, lobis acutis, glabriuscula. Pedunculi axillares, plerumque uniflori, supernè incrassati,
medium

medium versus bracteis binis, oppositis, parvis, muniti. Flores incarnati, albo varii, pulchri, suavcolentes, cælo fereno tantum expansi.

2. CONVULVULUS foliis sagittatis posticè truncatis, pedunculis tetragonis unifloris, bracteis cordatis calyci approximatis. *sepium.*

Convolvulus sepium. Sp. Pl. 218. Hudf. 88.

With. 240. Relb. 87. Sibth. 76. Curt. Lond.

fasc. 1. t. 13. Mart. Fl. Rust. t. 88. Engl.

Bot. t. 313. Fl. Dan. t. 458.

C. major. Raii Syn. 275.

Smilax lenis five lævis major. Ger. em. 861.

Angl. Great Bindweed.

In sepibus subhumidis, et falicetis.

Perennis. Fl. Julio, Augusto.

Radix repens, vix eradicanda. Caules volubiles, sexpedales, angulati. Folia alterna, petiolata, dependentia, sagittata, angulis posticis truncatis. Pedunculi tetragoni, uniflori. Bractææ cordatæ, calycem occultantes. Flores formosi, magni, plerumque albi, subinde incarnati, aliquando saturatè rosei.

3. CONVULVULUS foliis reniformibus, pedunculis unifloris, angulis alatis. *Soldanella.*

Convolvulus Soldanella. Sp. Pl. 226. Hudf. 89.

With. 240. Engl. Bot. t. 314.

C. maritimus, Soldanella dictus. Raii Syn. 276.

Soldanella marina. Ger. em. 838.

Brassica marina. Cord. Hist. 205. 2. Matth.

Valgr. v. 1. 427.

Angl. Sea Bindweed.

In littoribus maritimis arenosis.

Perennis. Fl. Julio.

Radix repens, ut in prioribus. Caules procumbentes, foliosi, breves, pauciflori. Folia alterna, petiolata, reniformia,

formia, subinde cordata et angulata, lævia, subcarnosa. Pedunculi axillares, solitarii, erectiusculi, supernè incrassati, quadranguli, angulis dilatatis, membranaceis, uniflori. Flores magni, erecti, incarnati, plicis longitudinalibus flavidis, foli meridiano expansi, fugaces. Capsula subrotunda. Semina angulata, magna.

90. POLEMONIUM. *G. Pl.* 87. *Juss.* 136.
Gærtn. t. 62.

Cor. quinquepartita. *Stam.* squamis basin corollæ claudentibus inserta. *Stigma* trifidum.
Capsula supera, trilocularis.

*cærule-
um.*

1. POLEMONIUM foliis pinnatis, floribus erectis, calycibus tubo corollæ longioribus.

Polemonium cæruleum. *Sp. Pl.* 230. *Hudsf.*

89. *With.* 241. *Engl. Bot. t.* 14. *Fl. Dan.*
t. 255.

P. vulgare cæruleum. *Raii Syn.* 288.

Valeriana græca. *Ger. em.* 1076.

Angl. Jacob's Ladder. *Greek* Valerian.

In dumetis Angliæ borealis rariùs.

Below Malham Cove, Yorkshire. *Ray.* At the Lover's Leap, Buxton, Mr. Wood; near Bake-well, Mr. Whately; in Gordale, Yorkshire, Mr. Gough. *Withering.*

Perennis. *Fl.* Junio.

Radix fibrosa. *Herba* glabra. *Caulis* erectus, bipedalis, foliosus, paniculatus. *Folia* alterna, impari-pinnata, multijuga, foliolis elliptico-lanceolatis, integerrimis. *Flores* campanulato-rotati, cærulei, sæpè albi. *Calyx* campanulatus, semiquinquesfidus.

91. CAMPANULA. *G. Pl.* 88. *Juss.* 164.
Gærtn. t. 31.

Cor. campanulata, fundo clauso valvis stamini-feris.

feris. *Stigma* trifidum. *Capf.* infera, poris lateralibus dehiscentes.

1. CAMPANULA foliis radicalibus reniformibus ; *rotundifolia.*
caulinis linearibus.

Campanula rotundifolia. *Sp. Pl.* 232. *Huds.* 95.

With. 241. *Relb.* 93. *Sibth.* 248. *Curt.*

Lond. fasc. 4. *t.* 21. *Fl. Dan.* *t.* 855. *Raii*

Syn. 277. *Ger. em.* 452.

β. *C. uniflora.* *Huds. ed.* 1. 81. nec *Linn.* *Fl.*
Dan. *t.* 189.

C. minor alpina, rotundioribus imis foliis. *Raii*

Syn. 277. *Chuf. Hist. lib.* 5. 173.

Angl. Round-leaved Bell-flower.

In ericetis, muris, et ad margines agrorum. β in
rupibus alpinis.

Perennis. *Fl.* Augusto, Septembri.

Radix sublignosa. *Caules* erecti, teretes, glabri, foliosi,
uni- vel multi-flori. *Folia* glabra ; radicalia petiolata,
cordata vel reniformia, ferrata ; caulina sessilia, alterna,
linearia, plerumque integerrima. *Flores* subpaniculati,
tremuli, cærulei, rarius albi. *Dentes calycini* subulati,
integerrimi, horizontaliter patentes ; in β erectiusculi.

2. CAMPANULA foliis strictis : radicalibus lan- *patula.*
ceolato-ovalibus, paniculâ patulâ, calyce den-
ticulato.

Campanula patula. *Sp. Pl.* 232. *Huds.* 95.

With. 242. *Engl. Bot.* *t.* 42. *Fl. Dan.*

t. 373.

C. esculentæ facie, ramis et floribus patulis.

Dill. Hort. Elth. 68. *t.* 58.

C. minor alba, five purpurea. *Ger. em.* 452 ?

Angl. Spreading Bell-flower.

In pascuis et ad sepes rarius.

On the outside of Buddon wood, Leicestershire.

Mr. Woodward. About Worcester, Malvern,
Litchfield,

Litchfield, and near Hagley. *Withering*. Near Holt in the road to Binham, Norfolk. *Rev. R. B. Francis, F.L.S.* At Cobham, Surrey.

Biennis. Fl. Julio, Augusto.

Radix attenuata, alba. Caulis erectus, angulatus, scaber, foliosus, paniculatus. Folia scabriuscula, crenata, elliptico-lanceolata; radicalia obtusa, latiora, petiolata; caulina acuminata, sessilia, alterna. Panicula patentiissima. Flores erecti, basi attenuati, dilute violacci. Dentes calycini lanceolati, utrinque basin versus denticulati.

Rapunculus.

3. CAMPANULA foliis undulatis: radicalibus lanceolato-ovalibus, paniculâ coarctatâ.

Campanula Rapunculus. *Sp. Pl.* 232. *Huds.* 95. *With.* 242. *Engl. Bot. t.* 283.

Rapunculus esculentus. *Raii Syn.* 277.

Rapuntium parvum. *Ger. em.* 453.

Rapum sylvestre. *Fuchs. Hist.* 214. *Ic.* 122.

Raponeoli. *Brunf. Herb. v.* 2. 84.

Angl. Rampion Bell-flower.

In aggeribus fossarum, et arvis requietis.

Near Croydon and Esher in Surrey. *Huds.* At Hindlip, Worcestershire. *Stokes.* Enville, Staffordshire. *With.* Near the castle at Old Buckenham, Norfolk. *Mr. Pitchford.*

Biennis. Fl. Julio, Augusto.

Radix fusiformis, alba, dulcis, esculenta. Caulis erectus, virgatus, angulatus, foliosus, scaber, pilis deflexis. Folia alterna, lanceolata, sessilia, scabra, undulata, denticulata, infima subelliptica. Panicula erecta, multiflora, sub-cylindræca, compacta. Pedunculi conferti, longitudine varii, uni- vel multi-flori, erecti. Dentes calycini subulati, vel integerrimi, vel basi denticulati. Flores ventricosi, præcedente minores.

latifolia. 4. CAMPANULA foliis ovato-lanceolatis, caule simpli-

simplicissimo tereti, pedunculis unifloris, fructibus cernuis.

Campanula latifolia. *Sp. Pl.* 233. *Huds.* 96.
With. 243. *Relb. Suppl.* 2. 10. *Abbot.* 48.
Lightf. 141. *Fl. Dan.* t. 85. *Engl. Bot.*
 t. 302.

C. maxima foliis latissimis. *Raii Syn.* 276.

Trachelium majus Belgarum, five giganteum.
Ger. em. 448.

Angl. Giant Bell-flower.

In dumetis et sepibus umbrosis.

Not unfrequent in Scotland, and the north of England; more rare in the south. In Suffolk on a clay soil. *Mr. Woodward.* Forehoe wood near Kimberley, Norfolk. *Mr. Rose.* Near Dunstable. *Abbot.*

Perennis. Fl. Augusto.

Radix ramosa, lactescens. *Caulis* erectus, 3- vel 4-pedalis, teretiusculus, vel polygonus, angulis obsolete, hispidulus, foliosus, multiflorus. *Folia* ovata, subpetiolata, duplicato-serrata, scabra. *Pedunculi* axillares, uniflori, vix ac ne vix biflori, erecti; fructiferi deflexi. *Flores* maximi, speciosi, cæruleo-violacei. *Calyx* ferulatus, subciliatus.

5. *CAMPANULA* foliis cordato-lanceolatis, caule *rapuncu-*
 ramofo, floribus sparsis secundis nutantibus, *loides.*
 calycibus reflexis.

Campanula rapunculoides. *Sp. Pl.* 234.

C. n. 692. *Hall. Hist.* v. 1. 307.

C. repens, flore minore cæruleo. *Baub. Hist.*
 v. 2. 806.

C. hortensis, *Rapunculi radice repente.* *Morif.*
Hist. v. 2. 460. *sect.* 5. t. 3. f. 32.

Angl. Creeping Bell-flower.

In dumetis rarissime.

At Blair in Scotland. *Fenwick Skrimshire, M. D.*
In some woods in Oxfordshire, among yew trees.
Herb. Buddl.

Perennis. Fl. Augusto.

Radix repens. Caulis erectus, supernè ramosus, foliosus, teres, hispidulus, pilis arctè deflexis. Folia cordato-lanceolata, inæqualiter ferrata, scabra; infima petiolata; summa sessilia. Flores racemosi, foliis sensim diminutis bracteolati, pedunculati, secundi, nutantes, magnitudine C. Rapunculi, cærulei. Calyx scaber, laciniis lanceolatis, integerrimis, reflexis.

Trachelium.

6. CAMPANULA caule angulato, foliis cordato-lanceolatis acutè ferratis, calycibusque hispidis, pedunculis axillaribus paucifloris.

Campanula Trachelium. Sp. Pl. 235. Hudf. 96. Wurb. 243. Relb. 94. Sibth. 80. Engl. Bot. t. 12. Fl. Dan. t. 1026.

C. vulgatioꝝ foliis Urticæ, vel major et asperior.
Raii Syn. 276.

Trachelium majus. Ger. em. 448.
Angl. Nettle-leaved Bell-flower.

In nemorosis, dumetis et sepibus.

Perennis. Fl. Julio.

Caulis angulato-teres, angulis subulatis, strictus, hispidus, foliosus. Folia petiolata, ovato-lanceolata; infima cordata; omnia profundè et inæqualiter ferrata, scabra, plerumque fetis pungentibus hispida. Pedunculi axillares, apud nos rariùs triflori, plerumque uniflori. Flores violacei, C. latifoliæ parùm minores. Calyx basi margineque hispidus, quandoque glaber.

glomerata.

7. CAMPANULA caule angulato simplici, floribus sessilibus, capitulo terminali, foliis ovatis crenatis.

Campanula glomerata. Sp. Pl. 235. Hudf. 96. Wurb.

With. 244. *Relb.* 94. *Sibth.* 80. *Engl. Bot.* t. 90.

C. pratensis, flore conglomerato. *Raii Syn.* 277.
Trachelium minus. *Ger. em.* 449.

Angl. Clustered Bell-flower.

In campis elatis calcareis.

Perennis. Fl. Julio, Augusto.

Radix lignosa. Caulis pedalis, erectus, simplex, teretiusculus, foliosus, scabriusculus. Folia ovata, crenata, pilosa, subtus pallida; radicalia longius petiolata, atque subcordata; caulina amplexicaulia. Flores omnes ferè terminales, capitati, bracteis ovatis obvallati, violacei, speciosi.

Huc pertinet *Gentiana collina*, *With.* 282. t. 11. f. 8.

8. **CAMPANULA** caule basi subramoso stricto, *hybrida*, foliis oblongis crenatis undulatis, corollâ calyce brevior, capsulâ prismaticâ.

Campanula hybrida. *Sp. Pl.* 239. *Huds.* 97.

With. 244. *Relb.* 95. *Sibth.* 81. *Engl. Bot.* t. 375. *Abbot.* 49.

C. arvensis erecta. *Raii Syn.* 278.

Speculum Veneris minus. *Ger. em.* 439.

Angl. Corn Bell-flower.

In arvis cretaceis.

Annua. Fl. Augusto.

Radix fibrosa, parva. Caulis basi plerumque ramosus, erectiusculus, flexuosus, angulatus, foliosus. Folia alterna, remota, sessilia, oblonga, undulata, subcrenata, margine scabra. Flores terminales, sessiles. Germen elongatum, prismaticum. Corolla violacea, calyce duplò brevior.

C. Speculum vix differt, nisi corollâ majore, cauleque ramosiori.

9. **CAMPANULA** foliis cordatis quinquelobis *hederacea*, petiolatis glabris, caule laxo.

Campa-

Campanula hederacea. *Sp. Pl.* 240. *Huds.* 97.
With. 244. *Dicks. Dr. Plants*, 56. *H. Sicc.*
fasc. 10. 9. *Engl. Bot. t.* 73. *Fl. Dan.*
t. 330.

C. cymbalariae foliis. *Raii Syn.* 277. *Ger.*
em. 452.

C. folio hederacco. *Bauh. Hist. v.* 2. 797.
Angl. Ivy-leaved Bell-flower.

In umbrosis humidis.

In Cornwall, Devonshire, Suffex and Wales not uncommon, also in some parts of Yorkshire and Oxfordshire. *Huds. With.* On a bog near High-beech, Epping Forest. *Mr. E. Forster.*

Perennis. *Fl.* Junio—Augusto.

Radices fibrosæ, parvæ. *Caules* prostrati, repentes, latè diffusi, graciles, ramosi, glabri, foliosi. *Folia* opposita vel alterna, petiolata, cordata, angulata, tenera, glabriuscula. *Pedunculi* sæpiùs terminales, longissimi, solitarii, uniflori. *Flores* nutantes, pallidè cyanei, fugaces. *Capfula* parva, hemisphærica.

92. PHYTEUMA. *G. Pl.* 89. *Juss.* 165.
Gærtn. t. 30.

Cor. rotata, quinquepartita; laciniis linearibus.
Stigma bi- five tri-fidum. *Capf.* bi- five tri-
 locularis, infera, lateraliter dehiscentes.

orbicula- I. PHYTEUMA capitulo subrotundo, foliis cre-
 re. natis; radicalibus cordatis.

Phyteuma orbicularis. *Sp. Pl.* 242. *Huds.* 97.
With. 245. *Engl. Bot. t.* 142. *Dicks. H.*
Sicc. fasc. 9. 1. *Jacq. Austr. t.* 437.

Rapunculus corniculatus montanus. *Raii Syn.*
 278. *Ger. em.* 455.

Angl. Round-headed Rampion.

In pascuis et ad vias, solo cretaceo, rariùs.

On the downs of Suffex, Hampshire, and Surrey.
Plentiful near Leatherhead.

Perennis. Fl. Augusto.

Radix lignosa. Caulis pedalis, erectus, simplex. Folia crenata, ciliata; radicalia cordato-ovata, petiolata; caulina sessilia, lanceolata, sensim minora. Capitulum terminale, erectum, multiflorum, bracteatum, post florescentiam elongatum. Flores violacei, sessiles, coarctati.

93. JASIONE. *G. Pl.* 455. *Juss.* 166.
Gærtn. t. 30.

Cor. rotata, quinquepartita; laciniis linearibus.
Stigma clavatum, bifidum. *Antheræ* basi connexæ. *Capf.* infera, semibilocularis, apice dehiscens.

I. JASIONE.

montana.

Jasione montana. *Sp. Pl.* 1317. *Huds.* 377.
With. 247. *Relh.* 328. *Sibth.* 85. *Curt. Lond.*
fasc. 4. *t.* 58. *Fl. Dan. t.* 319.

Rapunculus Scabiosæ capitulo cærulco. *Raii*
Syn. 278.

Scabiosa minima hirsuta. *Ger. em.* 723.

Angl. Sheep's-bit. Sheep's Scabious.

In pascuis et ericetis ficcioribus, arenosis, frequens.

Annua. Fl. Junio, Julio.

Radix sublignosa. Caules plurimi, erectiusculi, spithamæi, subinde pedales, terctiusculi, scabri, foliosi, apice subramosi. Folia lineari-oblonga, undulata, scabra; radicalia maxima; caulina alterna. Capitula terminalia, solitaria, erecta, hemisphærica, multiflora, involu-crata, involucri polyphylo, persistente. Flores pedicellati, cærulei. Corolla profundè quinquepartita. Antheræ basi connatæ, apice divergentes. Germen obovatum, calyce quinquesido persistente coronatum. Stylus staminibus longior. Stigma clavatum, emarginatum. Capsula, monente Gærtnero, non perfectè bilocularis, e dissepimentis ad axem fissis, neque receptaculo cen-

trali adnatis; quâ notâ ab omnibus reliquis *Campanulaceis* differt. Genus certè *Phyteumati* proximum, nec nisi fructu vix biloculari, vel antheris basi connexis, levi quidem discrimine, dignoscendum.

94. LOBELIA. *G. Pl.* 456. *Juss.* 165.

Rapuntium. *Gertn. l.* 30.

Cor. irregularis, suprâ longitudinaliter fissâ.

Stigma simplex. *Antheræ* connatæ in tubum.

Caps. infera, bi- five tri-locularis.

*Dort-
manna.*

1. LOBELIA foliis linearibus bilocularibus integerrimis, caule subnudo.

Lobelia Dortmanna. Sp. Pl. 1318. *Huds.* 377.

With. 245. *Engl. Bot. t.* 140. *Dicks. H. Sicc.*

fasc. 1. 14. *Lightf.* 505, *t.* 21. *Fl. Dan.*

t. 39.

Gladiolus lacustris. Ger. em. 105.

G. lacustris Dortmanni. Raii Syn. *287.

Angl. Water Lobelia.

In lacubus Angliæ borealis, Cambriæ, et Scotiæ.

In most of the lakes with clear gravelly bottoms in Westmoreland and Cumberland. Near the inn at Bowness on the banks of Wynandermere plentifully.

Perennis. Fl. Julio.

Radix e fibris albis, simplicibus. Herba glabra, lactescens. Caulis plerumque solitarius, erectus, simplex, teres, nudiusculus, cavus, aquâ ad medium usque immersus, apice racemosus. Folia omnino innumera, omnia ferè radicalia, linearia, recurva, depressò-teretiuscula, intùs bilocularia, dissépimento longitudinali. Flores remotiusculi, pedicellati, nutantes, cyanei. Bractæ solitariae ad basin pedicellorum, ovatae, breves. Calyx pentaphyllus, apicem germinis cingens, tubo brevior. Stamina intra fissuram corollæ fursùm exserta, antheris connatis stylum arctè amplectentibus. Stigma staminibus paulò longius, emarginatum. Capsula elliptica, bilocularis.

2. LOBELIA caule erectiusculo, foliis inferioribus *urens* obovatis denticulatis; superioribus lanceolatis ferratis, floribus racemosis.

Lobelia urens. *Sp. Pl.* 1321. *Huds.* 378.

With. 246. *Curt. Lond. fasc.* 6. *t.* 63.

Dickf. H. Sicc. fasc. 16. 9.

Rapunculus galeatus Blæsenfis, seu Soloniensis, flore violacco minore. *Morif. Hist. v.* 2. 467. *sect.* 5. *t.* 5. *f.* 56.

Rapuntium urens Soloniense. *Bocc. Sic.* 20. *t.* 11.

Angl. *Acrid Lobelia.*

In ericetis glareosis dumosis Angliæ australioris.

On Shute common between Axminster and Honiton, Devonshire. *Mr. Newberry.* On the slope of a heath called Kilmington hill, two miles from Axminster, close to the road; Lord Webb Seymour. *Curtis.*

Perennis. Fl. Augusto, Septembri.

Radix fibrosa. Herba glabra, lactescens, acris. Caulis erectus, ramosus, foliosus, scabriusculus, obsolete angulatus. Folia sparsa, lanceolata, denticulato-ferrata; inferiora obovata. Racemi terminales, multiflori. Flores brevius pedicellati, secundi, bracteis fetaceis, scabris, solitariis, ad basin pedunculorum. Corolla dilute violacea. Antheræ pubescentes. Calycis foliola fetacea, scabra, longitudine tubi.

95. IMPATIENS. *G. Pl.* 458.

Balsamina. *Juss.* 270. *Gærtn. t.* 113.

Cor. pentapetala, irregularis, nectario cucullato.

Antheræ apice connatæ. *Caps.* supera, quinquevalvis, elastica. *Cal.* diphyllus.

1. IMPATIENS pedunculis multifloris solitariis, *Noli mē tangere.* foliis ovatis, geniculis caulinis tumentibus.

Impatiens Noli me tangere. *Sp. Pl.* 1329.

Huds. 380. *With.* 263. *Fl. Dan.* t. 582.

Balsamine lutea, five Noli me tangere. *Raii*

Syn. 316.

Balsamita altera Noli me tangere. *Col. Ecphr.*

t. 150.

Perficaria filiquosa. *Ger. em.* 446.

Angl. Yellow Balsam. Touch me not.

In umbrosis humidis irriguis rariùs.

In several parts of Westmoreland; also in Lancashire, Yorkshire and Wales. *Ray.* On the banks of Wynandermerc, in little brooks and watery places near Rydall-hall, plentifully.

Annua. Fl. Augusto.

Radix fibrosa, parva. Herba succosa, tenerrima, pallida et subpellucida, glabra, ramosissima, ramis alternis. Folia alterna, petiolata, ovata, obtusa, ferrata. Pedunculi axillares, solitarii, racemosi, 3- vel 4-flori. Flores citrini, rubro punctati, elegantes. Calycis foliola æqualia, opposita, concava. Petala maximè inæqualia. Nectarium infundibuliforme, apice recurvo. Capsula oblonga gibbosa, pendula; valvulis elasticè diffilientibus ad semina, vix licet matura, spargenda.

Folia omninò noctu flaccida dependent. *Villars Dauph.* v. 2. 670.

96. VIOIA. *G. Pl.* 457. *Juss.* 294.

Gertn. t. 112.

Cor. pentapetala, irregularis, posticè cornuta.

Antheræ subcohærentes. *Caps.* supera, trivalvis, unilocularis. *Cal.* pentaphyllus, basi productus.

birta. 1. VIOIA acaulis, foliis cordatis petiolisque pilosohispidis, calycibus obtusis.

Viola birta. *Sp. Pl.* 1324. *Huds.* 379. *With.*

260. *Relb.* 329. *Sibth.* 83. *Curt. Lond.* fasc.

fasc. 1. t. 64. Fl. Dan. t. 618. Sym. Syn. 60.

V. martia major, hirsuta, inodora. Morif. Hist.

v. 2. 475. sect. 5. t. 35. f. 4. Raii Syn. 365. Angl. Hairy Violet.

In nemorosis et sepibus, solo calcareo, rariùs.

At Marham, Norfolk. *Mr. Crowe.* Near Bury.

Mr. Woodward. On St. Vincent's rocks, Bristol. *Mr. Dyer.*

Perennis. Fl. Aprili.

Radix sublignosa, fibrosa. Caulis nullus. Stolones breves, horizontales, simplices, cæspitosi, non radican-tes. Folia erectiuscula, longiùs petiolata, cordata, ferrata, venosa, utrinque piloso-hispida. Petioli hispidi, pilis patenti-deflexis, brevibus. Stipulæ lanceolatae, radicales, dentatae. Scapi foliis longiores, glabri, nec pilosi, erecti, uniflori, bracteis geminis, subulatis, infra medium positis. Flores nutantes, cærulei, inodori. Calyx glaber. Petala lateralìa lineâ pilosâ supra basin notata. Antheræ membranaceo-dilatatae, approximatae, vix coherentes. Capsula pilosa.

2. VIOLA acaulis, stolonibus reptantibus, foliis *odorata*, cordatis petiolisque glabriusculis, calycibus obtusis.

Viola odorata. Sp. Pl. 1324. Hudf. 378. With. 260.

Relb. 329. Sibth. 83. Curt. Lond. fasc. 1. t. 63.

Fl. Dan. t. 309. Sym. Syn. 60. Engl. Bot. t. 619.

V. martia purpurea. Raii Syn. 364.

V. nigra sive purpurea. Ger. em. 850.

β. *V. martia alba. Raii Syn. 364.*

V. flore albo. Ger. em. 850.

Viola, flos recens. Pharmac. Lond.

Angl. Sweet Violet.

In nemorosis, sepibus, et aggeribus fossarum.

Perennis. Fl. Martio, Aprili.

Radix præcedentis. Caulis nullus. Stolones longissimi, prostrati, radicanes, foliosi, etiam primo anno sæpè floriferi. Folia ut in priore, sed glabra, subtùs levissimè pubescentia. Petioli vel omninò glabri, vel pilis raris hirsutiusculi. Stipulæ binæ, lanceolatæ, dentatæ. Scapi supra medium bracteati, glabri. Flores saturatè purpurei, fragrantès, odore omnibus grato; subinde albi, vel carnei. Cætera ut in præcedente. Has optimè distinxit Curtisius, qui in utrisque flores seriores apetalos invenit.

palustris. 3. VIOLA acaulis, foliis reniformibus glabris, radice repente.

Viola palustris. *Sp. Pl.* 1324. *Huds.* 379. *With.* 261. *Lightf.* 506. *Curt. Lond. fasc.* 3. t. 58. *Engl. Bot.* t. 444. *Abbot.* 190. t. 3.

V. palustris rotundifolia glabra. *Morif. Hist.* v. 2. 475. *sect.* 5. t. 35. f. 5. *Raii Syn.* 364.

β. V. rubra striata Eboracensis. *Raii Syn.* 365. *Fl. Dan.* t. 83.

Angl. Marsh Violet.

In paludibus muscosis.

Perennis. Fl. Aprili.

Radix fibrosa, repens. Caulis nullus. Folia cordato-reniformia, crenata, glabra. Petioli semicylindrici, glaberrimi. Stipulæ lato-lanceolatæ, ferrulatæ. Scapi tetragoni, glabri, medium versus bracteati. Flores præcedentibus minores, inodori, pallidè cærulescentes vel carnei, venis saturationibus picti. Stigma urceolatum, suprà gibbosum.

canina. 4. VIOLA caule adultiore adscendente canaliculato, foliis oblongo-cordatis, calycibus acutis.

Viola canina. *Sp. Pl.* 1324. *Huds.* 379. *With.* 261. *Relh.* 329. *Sibth.* 83. *Curt. Lond. fasc.* 2. t. 61. *Engl. Bot.* t. 620.

V. canina sylvestris. *Ger. em.* 851.

V. martia

- V. *martia* inodora sylvestris. *Raii Syn.* 364.
 β. V. *canina* flore albo. *Merr. Pin.* 125. *Dill. in Raii Syn.* 364.
 γ. V. *caninæ* varietas minor. *Dill. in Raii Syn.* 364.
t. 24. *f.* 1.
 δ. V. *alpina*. *Hudf. ed.* 1. 379.
 V. *martia alpina*, folio tenello circinato. *Raii Syn.* 366.
Angl. Dog's Violet.

In nemorosis, dumetis, sepibus et ericetis vulgaris.

Perennis. Fl. Aprili, Junio.

Radix sublignosa. Herba primò acaulis est; at caules citò enascuntur, adscendentes, ramosi, foliosi, angulati, hinc canaliculati, glabriusculi. Folia cordato-oblonga, crenata, glabra, petiolis glabris apice subulatis. Stipulæ lanceolatæ, ferratæ, ciliatæ. Pedunculi axillares, solitarii, erecti, tetragoni, apicem versus bracteati, bracteis subulatis. Flores cærulei, inodori. Antheræ vix coherentes. Folia calycis, monente Curtisio, acuta. Stigma gibbosum. Capsula oblonga.

Varietatem alpinam, δ, ex Hudsono mutuatus sum, nec ipse inveni.

5. VIOLA caule adscendente tereti, foliis ovato-lanceolatis, stipulis inciso-ferratis. *lactea.*

Viola lactea. *Engl. Bot. t.* 445. *Syn. Syn.* 61.

V. *canina*, var. 3. *With.* 262.

Angl. Cream-coloured Violet.

In ericetis humidis montosis.

On the wolds at Tunbridge. *Mr. T. F. Forster jun.*

At Pendarvis, Cornwall; Mr. Stackhouse. *With.*

Perennis. Fl. Maio.

Radix fibrosa. Caules adscendentes, ramosi, teretes, foliosi, glabri, geniculis radicales. Folia longius petiolata, ovato-oblonga, basi in petiolum decurrentia, crenata, glabra. Stipulæ profundè ferratæ, sæpè inciso-pinnatifidæ, ferraturis apice glandulosis, nec setiferis.

Pedunculi elongati, supra medium bracteati, glabri. Bracteae lanceolatae. Calyx acutus. Corolla caeruleo-albida, vel dilute caerulea, venis purpureis. Antherae discretae.

Folia in hortis cordata evadunt, floresque caerulei. Forsitán præcedentis varietas. Videtur distincta a *V. pumila* *Villars Dauph. v. 2. 666.*

tricolor. 6. *VIOLA* caule angulato diffuso, foliis oblongis dentato-crenatis, stipulis lyrato-pinnatifidis.

Viola tricolor. *Sp. Pl.* 1326. *Huds.* 380. *With.* 262. *Relb.* 330. *Curt. Lond. fasc. 1. t. 65.* *Fl. Dan. t. 623.* *Woodv. Suppl. t. 252.* *Sym. Syn.* 61. *Raii Syn.* 365. *Ger. em.* 854.

Garden Pansie. *Pet. Herb. t. 37. f. 8.*

β. *Viola arvensis.* *Sibth.* 84. *Sym. Syn.* 61.

V. bicolor arvensis. *Raii Syn.* 366.

V. tricolor petræa. *Ger. em.* 854.

Corn Pansie. *Pet. H. Brit. t. 37. f. 9.*

Angl. Pansy Violet. Heart's-case.

In arvis.

Annua. Fl. Maio—Septembri.

Radix fibrosa, parva. Caulis ramosus, diffusus, foliosus, angulatus, pubescens; rami erecti. Folia alterna, petiolata, ovato-oblonga, magis vel minus profundè crenata vel ferrata, margine præcipuè pubescentia, subinde glabra; inferiora plerumque cordato-subrotunda. Stipulae pinnatifidæ, laciniâ terminali maximâ, ovatâ vel oblongâ, sæpius crenatâ. Flores axillares, solitarii, longissimè pedunculati, pedunculis striatis, nudis. Petala obcordata, flavo purpureoque varia, calyce plerumque longiora; in β breviora, albida.

lutea. 7. *VIOLA* caule triquetro simplici, foliis ovato-oblongis crenatis ciliatis, stipulis palmato-incisis,

Viola

Viola lutea. *With.* 263. *Sym. Syn.* 61. *Hudf.*
ed. 1. 331.

V. grandiflora. *Hudf.* 380. *Lightf.* 508.

V. montana lutea grandiflora nfrs. *Raii Syn.*
365.

Great Yellow Pansie. *Pet. H. Brit. t.* 37. *f.* 10.
Angl. Yellow Mountain Pansy.

In pascuis montosis.

Very common in the mountainous parts of Wales
and the north of England, as well as in Scotland.

Perennis. Fl. Majo—Septembri.

Radix fibrosa, tenuis. *Caulis simplex, adscendens, tri-*
queter, foliosus, vix pilosus, basi attenuatus et infirmus.
Folia præcedenti fimillima, at manifestius ciliata. Sti-
pulæ profundè palmato-lobatæ, laciniis oblongis, diva-
ricatis, ciliatis, exterioribus minoribus. Pedunculi
longissimi. Flores magnitudine V. tricoloris majoris,
lutei, basi nigro radiati, quandoque cærulei. Calyx
posticè dentatus.

97. VERBASCUM. *G. Pl.* 97. *Juss.* 124.

Gærtn. t. 55.

Cor. rotata, irregularis. Stam. declinata, bar-
bata. Caps. supera, bilocularis, valvulis in-
flexis, polysperma. Stigma simplex.

1. VERBASCUM foliis decurrentibus utrinque *Thapsus,*
tomentosis, caule simplici.

Verbascum Thapsus. *Sp. Pl.* 252. *Hudf.* 89.

With. 248. *Relb.* 88. *Sibth.* 77. *Engl.*

Bot. t. 549. *Fl. Dan. t.* 631.

V. mas latifolium luteum. *Raii Syn.* 287.

Thapsus barbatus. *Ger. em.* 773.

β. Var. 2, *Thapso-nigrum.* *With.* 248.

Angl. Great Mullein. High Taper.

In aggeribus et ruderatis, solo calcareo vel glareoso.

Biennis.

Biennis. Fl. Julio, Augusto.

Radix fusiformis. Caulis erectus, simplex, strictus, 3—5 pedalis, foliosus, tomentosus, angulatus, alatus. Folia alterna, decurrentia, oblonga, ferè integerrima, utrinque densissimè lanata, villis albis, ramosissimis, implexis. Spica terminalis, erecta, cylindracea, multiflora. Flores sessiles, coarctati, flavi, rarius albi. Stamina lutea. Antheræ miniatæ. Stigma clavatum.

β varietas hybrida, sterilis, patre V. nigro, sub oculis D. Edvardi Robson in horto enata, gaudet caule ramoso, foliis superioribus tantum decurrentibus, staminum villis violaceis. *With.*

Herba mucilaginis, emolliens, subnarcotica.

Lychnitis. 2. VERBASCUM foliis cuneiformi-oblongis supra denudatis, caule angulato paniculato.

Verbascum Lychnitis. *Sp. Pl.* 253. *Huds.* 90. *With.* 249. *Fl. Dan.* t. 536. *Engl. Bot.* t. 58.

V. flore albo parvo. *Raii Syn.* 287.

V. Lychnite Matthioli. *Ger. em.* 775.

β. V. Thapsi. *Sp. Pl.* 1669.

V. Thapsoides. *Huds.* 90. *With.* 249. *Syn. Syn.* 56.

V. angustifolium ramosum, flore aurco, folio crassiore. *Bauh. Hist.* v. 3. 860.

Angl. White Mullein.

In pascuis et ad vias, solo cretaceo.

Found chiefly in Kent, and there plentifully. In lanes about Dartford very common. At Kinver, Staffordshire. *Dr. Stokes.* β in Kent. *Huds.*

Biennis. Fl. Julio, Augusto,

Caulis erectus, tripedalis, strictus, angulatus, lanatus, foliosus, supernè paniculato-ramosissimus. Folia elliptico-oblonga, subcuneiformia, crenata, subtus densè lanata, supra denudata, reticulato-venosa; radicalia basi attenuata; caulina ovata, sessilia, nec decurrentia. Paniculæ rami racemosi, multiflori. Flores pedicellati, fasciculati,

fasciculati, lactei, staminibus flavescentibus, antheris croceis.

Varietatem β ex fide Linnæi et Hudsoni assumpsi, nec ipse adhuc vidi.

3. VERBASCUM foliis ovato-oblongis subser-^{pulveru-}
ratis utrinque pulverulentis, caule tereti pa-^{lentum,}
niculato.

Verbascum pulverulentum. *Curt. Cat. Salisb.*
Prod. 104. *Engl. Bot. t.* 487.

V. pulverulentum, flore luteo parvo. *Raii Syn.*
287.

V. *Lychnitis* β . *Huds.* 90. var. 1. *With.* 249.

β . Var. Nigro-pulverulentum.

Angl. Yellow Hoary Mullein. Norfolk Mullein.

In ruderatis, sepibus, et marginibus agrorum, solo
glareoso vel calcareo.

About Norwich every where, and in most parts of
Norfolk; also at Bury in Suffolk. At Woller-
ton near Nottingham. *Ray.*

β . At Helleston near Norwich frequent.

Biennis. Fl. Julio.

Herba tota tomento pulverulento, facile detergendo, tecta.
Caulis erectus, 3—4-pedalis, teres, foliolus, supernè
paniculatus, pyramidatus, ramosissimus, multiflorus.
Folia omnia sessilia, utrinque pulverulenta, obsolete
ferrata vel erenata; radicalia elliptico-oblonga, pedalia;
caulina ovata, acuminata, ferè amplexicaulia. Inflo-
rescentia præcedentis, sed ramosior, et flores majores,
aurei, staminum villo albo, antheris miniatis.

Var. β , e polline V. nigri ut videtur enata, gaudet habitu
V. pulverulenti, at foliis magis conceinnè erenatis, radi-
calibus petiolatis, caule petiolisque purpureis, staminum
villo violaceo.

4. VERBASCUM foliis oblongo-cordatis petio-^{nigrum,}
latis undulato-crenatis subpubescentibus.

Verbascum nigrum. *Sp. Pl.* 253. *Huds.* 90.
With.

With. 250. *Relb.* 88. *Sibth.* 77. *Engl. Bot.* t. 59. *Ger. em.* 775.

V. nigrum flore parvo, apicibus purpureis. *Raii* *Syn.* 288.

Angl. Dark or Black Mullein.

In glaucosis vel calcareis.

Plentiful in Kent, also in the retired grassy lanes so frequent in Norfolk and Suffolk. At Helleston and Costesby near Norwich.

Perennis. Fl. Julio, Augusto.

Caulis strictus, foliosus, angulatus, striatus, purpurascens, subinde pubescens, parum ramosus. Folia radicalia cordata, longè petiolata; caulina cordato-lanceolata, petiolata; summa ferè sessilia; omnia undulato-crenata, vel subserrata, reticulato-venosa, saturatè viridia, vel nuda, vel pubescentia. Spica terminalis, plerumque simplex et solitaria, multiflora, laxiuscula. Flores fasciculati, pedicellati, aurei, staminum villo violaceo, antheris croceis.

virga-
tum.

5. *VERBASCUM* foliis oblongo-lanceolatis dentatis sessilibus; radicalibus sublyratis pubescentibus, caule ramoso, floribus aggregatis subsessilibus.

Verbascum virgatum. *With.* 250. *Syn. Syn.* 56. *Engl. Bot. t.* 550.

Blattaria flore amplo. *Ger. em.* 778.

B. magno flore. *Bauh. Hist. v.* 3. 859. *Lob. Ic.* 564.

Angl. Large-flowered Mullein.

In arvis et ad vias.

First found by Mr. Waldron Hill of Worcester, in a field on the south side of a lane leading from Gregory's mill to the turnpike road, near that town. By the side of the turnpike road from Worcester to Ombersley, opposite to the lane leading to Bever. Cultivated for three years

without any perceptible alteration ; Dr. Stokes.
With.

Biennis. Fl. Augusto.

Caulis erectus, 5- vel 6-pedalis, a basi ramosus, lignosus, foliosus, teres, sed præ foliis decurrentibus ferè angulatus vel alatus. Rami alterni, simplices, elongati, virgati, foliosi, apice racemosi. Folia radicalia facie Primulæ vulgaris, lanceolato-oblonga, rugosa, inæqualiter dentato-crenata, basi pinnatifido-incisa ; suprà nitidiuscula, pilis sparsis ; subtùs venosa, lanata : caulina oblongo-lanceolata, dentata et incisa, subfessilia ; superiora cordata, dentata, fessilia. Flores fasciculati, subfessiles, flavi, ore purpurascente, maximi. *Stokes.*

6. VERBASCUM foliis amplexicaulibus oblongis *Blattaria.*
glabris ferratis, pedunculis unifloris solitariis. *ria.*

Verbascum Blattaria. *Sp. Pl.* 254. *Huds.* 91.

With. 251. *Engl. Bot. t.* 393.

Blattaria lutea. *Raii Syn.* 288.

B. Plinii. *Ger. em.* 776. *Lob. Ic.* 564.

Angl. Moth Mullein.

In glareosis rariùs.

In Devonshire and Cornwall not uncommon. On a bank about three miles from Rochester, near the river Medway. *Mr. Jacob Rayer.*

Annua. Fl. Julio.

Radix fusiformis. Caulis tripedalis, erectus, ramosus, foliosus, angulatus, glaber. Folia obovato-oblonga, duplicato-ferrata, glabra, amplexicaulia ; radicalia sublyrata. Racemi terminales, piloso-glandulosi, stricti, multiflori. Flores pedunculati, solitarii, aurei, purpureo lineati, bracteis solitariis ovatis ad basin pedunculorum. Stamina villis purpureis tecta.

98. DATURA. *G. Pl.* 98. *Juss.* 125.

Cor. infundibuliformis, plicata. *Cal.* tubulosus, angulatus, deciduus. *Capf.* supera, bilocularis, quadrivalvis.

Stramonium.

- * 1. *DATURA* pericarpis spinosis erectis ovatis; foliis ovatis glabris.

Datura Stramonium. *Sp. Pl.* 255. *Huds.* 92. *With.* 251. *Curt. Lond. fasc.* 6. *t.* 17. *Fl. Dan.* *t.* 436.

Solanum, pomo spinoso oblongo, flore calathoides, *Stramonium* vulgò dictum. *Raii Syn.* 266.

Stramonium spinosum. *Ger. em.* 348. *Angl.* Thorn-apple.

In finetis, ruderatis et ad vias, e rejectamentis hortorum.

By the road side beyond Brook, Norfolk, in the way to Bungay. About London not uncommon.

Annua. Fl. Julio.

Herba lurida, narcotica, fœtens, glabra. Caulis ramissimus, dichotomus, patens, tripedalis, teres, foliosus. Folia e dichotomiâ caulis, petiolata, ovata, acuta, sinuata, basi inæqualia. Flores axillares, solitarii, brevius pedunculati, erecti, albi. Calyx supra basin deciduus. Fructus quadrifurcatus, spinis horridus. Semina reniformia.

99. *HYOSCYAMUS.* *G. Fl.* 98. *Juss.* 124. *Gærtn. t.* 76.

Cor. infundibuliformis, obtusa, irregularis. *Stam.* inclinata. *Caps.* operculata, bilocularis.

niger.

1. *HYOSCYAMUS* foliis amplexicaulibus sinuatis, floribus sessilibus.

Hyoscyamus niger. *Sp. Pl.* 257. *Huds.* 92. *With.* 252. *Relb.* 90. *Sibth.* 77. *Ger. em.* 353. *Woodv. Med. Bot. t.* 52. *Engl. Bot. t.* 591. *Dreves. Bilderb. t.* 47.

H. vulgaris. *Raii Syn.* 274.

Hyoscyamus.

Hyoicyamus. *Branf. Herb.* 229. *Riv. Monop.*

Irr. t. 102.

Angl. Common Henbane.

In rudcratis, aggeribus et ad vias.

Anna. Fl. Julio.

Radix fusiformis. Herba viscosa, pilosa, foetidissima.

Caulis ramosus, teres. Folia alterna, sessilia, oblonga, pinnatifido-sinuata. Spicæ terminales, simplices, recurvæ, foliosæ. Flores sessiles, erecti. Calyx urceolatus, margine quinquesidus, reticulato-venosus, persistens. Corolla flava, venis violaceis pulcherrimè picta.

Planta tota venenata, narcotica; semina tantum, ex oleo cotyledonum, innoxia.

100. ATROPA. *G. Pl.* 99. *Juss.* 125.

Gertr. t. 131.

Cor. campanulata. *Stam.* distantia. *Bacca* super, bilocularis.

1. ATROPA caule herbaceo, foliis ovatis integris. *Bella-*
Atropa Belladonna. *Sp. Pl.* 260. *Huds.* 93. *donna.*

With. 252. *Relb.* 90. *Sibth.* 78. *Curt. Lond.*

fasc. 5. *t.* 16. *Woodw. Med. Bot. t.* 1. *Jacq.*

Austr. t. 309. *Engl. Bot. t.* 592.

Belladonna. *Raii Syn.* 265.

Solanum lethale. *Ger. em.* 340.

Angl. Dwale, or Deadly Nightshade.

In sepibus et rudcratis, præcipuè solo calcareo.

Perennis. Fl. Junio.

Radix carnosa, crassa, repens. Caules herbacei, annui, erecti, tripedales, ramosi, foliosi, teretes, subpubescentes. Folia lateralialia, gemina, magnitudine inæqualia, petiolata, ovata, acuta, integerrima, sordidè viridia, glabriuscula. Pedunculi laterales, subaxillares, solitarii, uniflori, teretes, nutantes. Flores sordidè violacei. Calyx subpubescens, viscosus. Antheræ magnæ, albidæ. Stigma capitatum. Bacea depressa, sulcata,

nitida, atro-purpurea, polysperma, succo violaceo, subdulci, maximè narcotica et lethalis.

101. SOLANUM. *G. Pl.* 100. *Juss.* 126.

Gærtn. t. 131.

Cor. rotata. *Antheræ* subcoalitæ, apice poro gemino dehiscentes. *Bacca* supera, bilocularis.

Dulcamara.

1. SOLANUM caule inermi frutescente flexuoso, foliis superioribus hastatis, racemis cymosis.

Solanum Dulcamara. Sp. Pl. 264. *Huds.* 93.

With. 253. *Relb.* 91. *Sibth.* 78. *Curt.*

Lond. fasc. 1. t. 14. *Fl. Dan. t.* 607. *Woodv.*

Med. Bot. t. 33. *Engl. Bot. t.* 565.

S. lignosum, seu *Dulcamara. Raii Syn.* 265.

Amara dulcis. Ger. em. 350.

β. *Solanum lignosum*, seu *Dulcamara marina. Raii Syn.* 265.

Angl. Woody Nightshade. Bitter-sweet.

In sepibus humidis, et umbrosis.

Frutex. Fl. Junio, Julio.

Radix lignosa. Caulis fruticosus, scandens, teretiusculus, ramulosus. Folia alterna, petiolata; inferiora cordata; superiora hastata; omnia integerrima, glabra; “in β hirsuta,” *Huds.* Flores in racemis vel paniculis cymæformibus, non verè cymosi, oppositifolii et terminales, nutantes, formosi, corollæ laciniis reflexis, violaceis, basi punctis duobus viridibus notatæ. *Antheræ* citrinæ, connatæ. *Bacca* elliptica, rubra, succosa, amara atque venenosa.

nigrum. 2. SOLANUM caule inermi herbáceo, foliis ovatis dentato-angulatis, umbellis lateralibus nutantibus.

Solanum nigrum. Sp. Pl. 266. *Huds.* 92.

With. 254. *Relb.* 91. *Sibth.* 78. *Curt.*

Lond. fasc. 2. t. 14. *Fl. Dan. t.* 460. *Engl.*

Bot. t. 566.

S. vul-

S. vulgare. *Raii Syn.* 265.

S. hortense. *Ger. em.* 339.

Angl. Common or Garden Nightshade.

In ruderatis, cultis, et fimetis.

Annua. Fl. Junio—Septembri.

Radix ramosa. *Caulis* ramosus, patens, foliosus, aliquando sublignosus, angulatus. *Folia* petiolata, ovata, basi producta, repanda, subpubescentia. *Umbellæ* laterales, ex internodiis caulinis, nutantes, paucifloræ, pubescentes. *Flores* albi, inoschati. *Baccæ* nigrae; subinde, monente Hudsono, lutci. *Planta* narcotica, dia-phoretica et diuretica, sed cautè adhibenda.

102. *CHIRONIA.* *G. Pl.* 101. *Juss.* 142.

Gærtn. t. 114.

Cor. hypocrateriformis. *Stam.* tubo inserta.

Antherae demum spirales. *Stylus* declinatus.

Pericarpium superum, valvulis inflexis biloculare.

1. *CHIRONIA* caule herbaceo dichotomè paniculato, foliis ovato-lanceolatis, calyce tubo brevioris. *Centaureum.*

Chironia Centaurium. *Curt. Lond. fasc.* 4. *t.* 22.

With. 255. *Sibth.* 75. *Engl. Bot. t.* 417.

Woodv. Med. Bot. t. 157. *Wilden. Sp. v.* 1. 1068.

Gentiana Centaurium. *Sp. Pl.* 332. *Hudsf.* 102.

Relb. 100. *Dicks. H. Sicc. fasc.* 7. 3.

Centaurium minus. *Raii Syn.* 286. *Tillandsf.*

Is. 27.

C. parvum. *Ger. em.* 547.

Centaureum minus, cacumen. *Pharmac. Lond.*

Angl. Common Centaury.

In pascuis glareosis siccioribus.

Annua. Fl. Julio, Augusto.

VOL. I.

S

Radix

Radix parva, ramosa. Caulis erectus, pedalis, foliosus, quadrangulus, supernè paniculato-dichotomus; rariùs e basi ramosus. Folia opposita, sessilia, elliptico-oblonga, integerrima, trinervia, lævia. Flores e dichotomiâ caulis, sessiles, erecti, soli clariori tantum expansi. Calyx semi-quinquesidus, laciniis subulatis, erectis, tubo corollæ adnatis, duplòque brevioribus. Corollæ limbus æqualis, patens, roseus, nitidus, laciniis ellipticis, concaviusculis. Stamina declinata. Antheræ ter tortæ. Stigma capitatum, bipartitum. Herba glabra, valdè amara.

Varietatem floribus candidissimis invenit Nobiliss. Marchio de Blandford, Soc. Linn. Soc. Honor.

pulchella. 2. **CHIRONIA** caule herbaceo undique ramosissimo, foliis ovatis, laciniis calycinis subulatis tubo parùm brevioribus.

Chironia pulchella. Swartz. *Act. Holm.* 1783, p. 85. t. 3. f. 8, 9. *With.* 255. *Engl. Bot.* t. 458.

Gentiana Centaurium β. *Sp. Pl.* 333. *With.* 255.

Centaurium minus palustre ramosissimum, flore purpureo. Vaill. *Paris.* 32. t. 6. f. 1. *Engl.* Dwarf branched Centaury.

In arenosis maritimis.

At Gorleston, Suffolk. *R. Stone Esq.* On the downs at Port Owen, on the north coast of Cornwall, J. Watt, Esq. *With.* Braunsion Burrows. *Rev. Dr. Goodenough.*

Annua. Fl. Augusto, Septembri.

Radix parva, ramosa. Caulis ramosissimus, vix biuncialis, dichotomus, quadrangulus, glaber. Folia ovato-elliptica, lævia, tri- vel quinque-nervia. Flores e dichotomiâ caulis, plerumque pedicellati, erecti. Calyx ad basin usque quinquesidus, laciniis subulatis, attenuatis, liberis, longitudine ferè tubi. Corolla gracillima, limbi laciniis elliptico-lanceolatis, roseis. Antheræ semitortæ.

Variat

Variat caule simplici, attenuato, paucifloro, calyce tamen semper dignoscenda.

103. SAMOLUS. *G. Pl.* 89. *Juss.* 97.

Gertn. t. 30.

Cor. hypocrateriformis, quinquefida, squamis interstinctis. *Stamina* tubo inserta, laciniis opposita. *Caps.* femiinfera, unilocularis, quinquedentatus.

1. SAMOLUS foliis obtusis, racemo multifloro, *Valerandi.*
pedicellis bracteolatis.

Samolus Valerandi. Sp. Pl. 243. *Huds.* 94.

With. 246. *Relb.* 92. *Sibth.* 79. *Curt.*

Lond. fasc. 4. *t.* 20. *Fl. Dan. t.* 198. *Raii*

Syn. 283.

Anagallis aquatica rotundifolia. Ger. em. 620.

Angl. Brookweed. *Water Pimpernel.*

In palustribus.

Perennis. Fl. Julio.

Radix e fibris longis, subsimplicibus, albis. Herba glabra. Caulis pedalis, erectus, foliosus, teres, apice subramosus. Folia alterna, subsessilia, obovata, obtusa, integerrima; radicalia petiolata, numerosa. Racemi terminales, erecti, multiflori. Pedicelli sæpius gemini, uniflori, erecti, geniculati, bractea ad geniculum solitaria, lanceolata. Flores nivei, parvi. Calyx campanulatus, vix femiquinquesidus. Corollæ tubus cylindricus; limbus patens, laciniis obtusis, æqualibus, squamis lanceolatis. Stamina tubo breviora, inclusa. Stylus brevissimus. Stigma capitatum. Capsula femiinfera, apice dehiscens, quinquevalvis. Semina numerosa, angulata, receptaculo centrali, globoso, inserta.

Genus Montiæ, itaque *Portulaccis* Jussieu, affine. Alia species est *Sheffieldia repens* Forsteri, monente amicissimo Dryandro.

104. LONICERA. *G. Pl.* 93.

Caprifolium. *Juss.* 212. *Gertn. t.* 27.

Cor. monopetala, irregularis. *Bacca* polysperma, infera.

Caprifolium. 1. LONICERA floribus ringentibus verticillatis terminalibus, foliis deciduis: summis connatoperfoliatis.

Lonicera Caprifolium. *Sp. Pl.* 246. *Jacq. Austr. t.* 357. *Ait. H. Kew. v.* 1. 230 α .

Periclymenum. *Matth. Valgr. v.* 2. 321. *Camer. Epit.* 713.

P. perfoliatum. *Ger. em.* 891.

Angl. Pale perfoliate Honeysuckle.

In nemorosis et dumetis.

In a wood near Elsfield, Oxfordshire, plentifully.

Rev. Mr. Butt. In Chalk-pit close, Hinton, Cambridgeshire, certainly wild; also in another coppice in the same parish. *Rev. Mr. Relban.*

Frutex. Fl. Maio, Junio.

Caulis lignosus, volubilis. *Rami* suboppositi, teretes, glabri. *Folia* omnia ferè connata, elliptica, obtusa, integerrima, glabra, subtùs glaucescentia; summa connatoperfoliata, suborbiculata, florifera. *Flores* verticillati, patentes, flavescentes, basi rubicundi, suaveolentes. *Baccæ* miniatæ, calyce ferè integerrimo coronatæ.

Periclymenum. 2. LONICERA capitulis ovatis imbricatis terminalibus, foliis omnibus distinctis deciduis, floribus ringentibus.

Lonicera Periclymenum. *Sp. Pl.* 247. *Huds.* 94. *Witb.* 246. *Relb.* 93. *Sibth.* 81. *Curt. Lond. fasc.* 1. t. 15. *Fl. Dan. t.* 908. *Ait. Hort. Kew. v.* 1. 231.

Caprifolium Germanicum. *Raii Syn.* 458.

Periclymenum. *Ger. em.* 891.

β . Capri-

β. Caprifolium non persoliatum, foliis finuosis.
Dill. in Raii Syn. 458.

Periclymenum foliis quercinis. *Merr. Pin.* 92.
Angl. Common Honeyfuckle or Woodbine.

In sepibus et nemorosis frequens.

β. In a wood near Kimberley, Norfolk. *Mr. Woodward.*

Frutex. Fl. Junio, Julio.

Caulis lignosus, volubilis. Rami oppositi, teretes. Folia opposita, subpetiolata, elliptica, integerrima, sæpius pubescentia, subtus glauca. Flores in capitulo terminali, radiato-patentes, ochroleuci, rubicundi, vespere imprimis suaveolentes. Baccæ rubræ, calyce quinque-dentato persistente coronatæ, amaræ, subdulces.

105. RHAMNUS. *G. Pl.* 105. *Juss.* 380.

Gærtn. t. 106.

Cal. urceolatus. *Petala* 5, staminibus opposita.
Bacca supera.

1. RHAMNUS spinis terminalibus, floribus quadrifidis dioicis, foliis ovatis, caule erecto, baccæ tetraspermæ. *catbarticus.*

Rhamnus catharticus. *Sp. Pl.* 279. *Huds.* 98.

With. 256. *Relb.* 95. *Sibth.* 82. *Woodv.*

Med. Bot. t. 114. *Fl. Dan. t.* 850. *Raii Syn.* 466.

R. solutivus. *Ger. em.* 1337.

Spina cervina, bacca. *Pharmac. Lond.*

Angl. Buckthorn.

In sepibus et nemoribus.

Frutex. Fl. Maio, Junio.

Rami rigidi, alterni, teretes, apice spinosi. Folia fasciculata, petiolata, ovata, serrata, nervosa; juniora pubescentia. Flores ex eâdem gemmâ cum foliis, pedunculati, luteo-virides, quadrifidi, plerumque, nec semper,

dioici. Stigma quadrifidum. Baccæ nigræ, tetraspermæ, catharticæ, succo in immaturis luteo tinctorio.

Frangula. 2. RHAMNUS inermis, floribus monogynis hermaphroditis, foliis integerrimis glabris, baccâ dispermâ.

Rhamnus Frangula. *Sp. Pl.* 280. *Huds.* 98. *With.* 259. *Kelb.* 96. *Fl. Dan. t.* 278. *Engl. Bot. t.* 250.

Frangula, seu *Alnus nigra baccifera.* *Raii Syn.* 465.

Alnus nigra five *Frangula.* *Ger. em.* 1470. *Angl.* Berry-bearing Alder. Alder Buckthorn.

In nemoribus et dumosis.

Frutex. *Fl. Maio.*

Frutex quadripedalis et ultra, ramosissimus. Folia opposita, petiolata, obovata, vel elliptico-lanceolata, integerrima, glabra, venosa. Pedunculi axillares, binati vel ternati, uniflori. Flores albidî, antheris atro-purpureis. Stigma bifidum. Bacca atro-violacea, disperma.

106. EUONYMUS. *G. Pl.* 107. *Juss.* 377. *Gærtn. t.* 113.

Cal. planus. *Petala* 5. *Caps.* pentagona, quinquelocularis, quinquevalvis, colorata. *Sem.* calyptrata.

europæus. 1. EUONYMUS floribus plerisque quadrifidis, petalis acutis, ramis lævibus.

Euonymus europæus. *Sp. Pl.* 286. *Huds.* 98. *With.* 259. *Relb.* 96. *Sibth.* 82. *Engl. Bot. t.* 362.

E. vulgaris. *Raii Syn.* 468.

E. Theophrasti. *Ger. em.* 1468.

Angl. Spindle-tree. Prickwood.

In sepibus et dumetis.

Frutex.

Frutex. Fl. Maio.

Rami glabri, virides, teretes; juniores quadranguli. Folia opposita, petiolata (nec sessilia) lanceolata, ferrata, glabra. Pedunculi axillares, dichotomi, divaricati. Flores albid, fœtidi, dempto primario quadrifidi. Capsulæ depressæ, angulatæ, purpureæ. Semina alba, arillo croceo.

107. RIBES. *G. Pl.* 111. *Juss.* 310.

Cal. superus, campanulatus, quinquefidus. *Petal.* et *Stamina* calyci inserta. *Stylus* bifidus. *Bacca* polysperma.

* *Inermia.*

1. RIBES inerme, racemis glabris pendulis, flori-*rubrum.* bus planiusculis, petalis obcordatis.

Ribes rubrum. *Sp. Pl.* 290. *Huds.* 99. *With.* 264. *Relb.* 97. *Sibth.* 84. *Lightf.* 146. *Woodv. Med. Bot. t.* 74. *Fl. Dan. t.* 967.

R. vulgaris fructu rubro. *Raii Syn.* 456. *Ger. em.* 1593.

β. *R. vulgaris* fructu dulci. *Raii Syn.* 456.

γ. *R. fructu parvo.* *Merr. Pin.* 104. *Dill. in Raii Syn.* 456.

Ribes rubrum, fructus. *Pharmac. Lond.* *Angl.* Common Currants.

In sylvis et ad ripas fluviorum Angliæ borealis.

Undoubtedly wild about the banks of the Tees.

Mr. Robson. In the isle of Iffa, amongst the brush-wood on the bank of the Sound. *Lightf.*

In hedges by accident frequent.

Frutex. Fl. Maio.

Rami glabri. Folia alterna, longius petiolata, quinqueloba, duplicato-ferrata, subpubescentia. Racemi simplices, nutantes, multiflori; fructiferi penduli. Bractæ solitariae, ovatae, minutæ, pedicellis multò breviores. Flores luteo-virides, cyathiformes. Petala ob-

cordata. Baccæ globosæ, acidæ, rubræ, in hortis sæpè albæ vel incarnatæ.

alpinum, 2. RIBES inerme, racemis erectis, bracteis flore longioribus, foliis subtus lucidis.

Ribes alpinum. *Sp. Pl.* 291. *Huds.* 99. *With.* 264. *Lightf.* 146. *Jacq. Austr.* t. 47. *Fl. Dan.* t. 968. *Dicks. H. Sicc. fasc.* 15. 13.

R. alpinus dulcis. *Raii Syn.* 456.

Angl. Tasteless Mountain Currants.

In sylvis, sepibus et dumetis, Angliæ borealis.

In Yorkshire, especially about Bradford. In Durham. *Mr. Robson*.

Frutex. Fl. Aprili, Maio.

Folia quam in præcedente minora, triloba, ferrata, glabriuscula, subtus præcipuè lucida. Racemi semper erecti, floribus brevius pedicellatis, luteo-viridibus. Bracteæ lanceolatæ, florem superantes. Baccæ ellipticæ, rubræ, mucilaginosæ et insipidæ.

Variat floribus dioicis, ex observationibus D. D. Sneyd & Robson. *With.*

spicatum, 3. RIBES inerme, spicis erectis, petalis oblongis, bracteis flore brevioribus.

Ribes spicatum. *Robson in Trans. of Linn. Soc.* v. 3. 240. t. 21. *With.* 265. *Sym. Syn.* 62.

Angl. Acid Mountain Currants.

In sylvis Angliæ borealis.

Near Richmond in Yorkshire, and between Piersbridge and Gainford in the county of Durham.

Mr. Robson.

Frutex. Fl. Maio.

Folia *R. rubri* habet, at lobis ferraturisque acutioribus, subtusque magis tomentosa. Spicæ semper erectæ, floribus brevissimè pedicellatis, rubido-fuscis. Bracteæ ovatæ, exiguæ, vix pedicellum, minimè florem, superantes, apice reflexæ. Petala cuneato-oblonga. Baccæ globosæ,

globosæ, colore et gustu R. rubro similes. Semina fertilia. *Robson.*

4. RIBES inerme, racemis erectis: fructiferis pen- *petræum.*
 dulis, petalis obtusis; bracteis flore brevioribus.
Ribes petræum. Wulf. in Jacq. Misc. v. 2. 36.
Jacq. Ic. Rar. v. 1. t. 49. Wilden. Sp. v. 1.
1153.
Angl. Rock Currants.

In montosis.

Near Egleston in the county of Durham. *Rev.*
Mr. Harriman. Near Consccliffe in that county.
Mr. Robson.

Frutex. Fl. Maio.

Flores racemosi, nec spicati. Racemi fructiferi penduli.
 Omnino diversum a R. alpino. *D. Harriman.*
 Facies prorsus antecedentis. *Anne species distincta?*

5. RIBES inerme, racemis pilosis pendulis pedun- *nigrum.*
 culo simplici ad basin, floribus oblongis.
Ribes nigrum. Sp. Pl. 291. Hudf. 99. With.
265. Relb. 97. Woodv. Med. Bot. t. 75. Fl.
Dan. t. 556.
R. nigrum vulgò dictum folio olente. Raii Syn.
456.
Ribes nigrum, fructus. Pharmac. Lond.
Angl. Black Currants.

In dumetis et sepibus humidis, et ad ripas fluviorum.

On Costesý island near Norwich. *Mr. Rose.* In
 alder swamps in Norfolk. *Mr. Woodward.* Not
 unfrequent in various parts of England, un-
 doubtedly wild. In the wood on the bank of
 the Sound of Isla. *Lightf.*

Frutex. Fl. Maio.

Dignoscitur habitu humiliore, foliis olentibus, subtùs
 glandulosis;

glandulosis, racemis pilosis, calyce tubuloso, fructu nigro, at præcipue pedunculo unifloro, solitario, ad basin racemi, ab eoque distincto.

Petala quandoque in stamina mutantur: metamorphosis hæcenus inaudita. *With.*

* * *Aculeata.*

Grossularia.

* 6. RIBES ramis aculeatis, petiolis pilosis, pedunculis unifloris, bracteis binis, fructu hirsuto.

Ribes Grossularia. *Sp. Pl.* 291. *With.* 266. *Sym. Syn.* 62.

R. Uva crispa. *Fl. Dan. t.* 546.

Angl. Rough Gooseberry.

In sepibus et rudcratis, vix indigena.

In hedges, Walcot; on old buildings and church towers, Stokes; in woods and hedges about Darlington, common; Mr. Robson. *With.*

Frutex. Fl. Aprili.

Fruticulus ramosus; aculei sub gemmis solitarii, binati, vel ternati. Folia triloba incisa, subpubescentia. Petioli pilosi, foliis sæpius longiores. Pedunculi uniflori, nutantes, medio bracteis binis vel ternis, oppositis, ovatis, ciliatis. Germen villosum. Baccæ pendulæ, hispidae.

Variat fructibus glabris. *D. Robson.* Anne species a frequente distincta?

Uva-crispa.

* 7. RIBES ramis aculeatis, pedunculis unifloris, bracteis connato-tubulosis, fructu glabro.

Ribes Uva-crispa. *Sp. Pl.* 292. *With.* 266. *Sym. Syn.* 62. *Relb.* 98. *Sibth.* 84. *Culhum.* 88.

Uva crispa. *Fuchf. Hist.* 187. *Ger. em.* 1324.

Angl. Smooth Gooseberry.

In sepibus et nemorosis.

In several places in Cambridgeshire. *Relb.* In Oxfordshire. *Sibth.* In Norfolk not unfrequent,

quent, but apparently not a truly indigenous plant.

Frutex. Fl. Aprili.

A præcedente vix differt, nisi baccis glabris, bracteis basi in tubum connatis, pilisque petiolorum glandulosis: notis lubricis reverà et incertis.

108. HEDERA. *G. Pl.* 111. *Juss.* 214.

Gærtn. t. 26.

Cal. quinquedentatus. *Petala* 5, basi dilatata.

Bacca pentasperma, calyce cincta.

1. HEDERA foliis ovatis lobatisque.

Helix.

Hedera Helix. *Sp. Pl.* 292. *Huds.* 100. *With.*

267. *Relb.* 98. *Sibth.* 85. *Curt. Lond.*

fasc. 1. *t.* 16. *Fl. Dan. t.* 1027.

H. communis major et minor. *Raii Syn.* 459.

H. corymbosa et *H. Helix.* *Ger. em.* 857.

Angl. Common Ivy.

In sylvis, sepibus, et ad muros.

Frutex. Fl. Octobri.

Caulis scandens, radicans, sub-complanatus, arborum truncis murisque arctè fulcris adhærens. Rami superne liberi, teretes, ramosissimi. Folia sparsa, petiolata, sempervirentia, integerrima, obscure viridia, lucida; caulina quinqueloba; summorum ramorum ovata, indivisa. Pedunculi communes corymbosi; partiales umbellati, tomento stellato tecti. Bractæ parvæ, ovatæ, acutæ. Flores virides. Petala ovata, basi lata. Stamina patentia, petalis longiora. Stylus brevis, fulcatus, stigmate simplici. Baccæ læves, nigræ, farinosæ, atomatæ.

109. ILLECEBRUM. *G. Pl.* 114. *Juss.* 89.

Cal. pentaphyllus, cartilagineus, inferus. *Cor.*

nulla. *Stigma* simplex. *Caps.* quinquevalvis, monosperma.

1. ILLE-

*verticil-
latum.*

1. ILLECEBRUM floribus verticillatis nudis,
caulibus procumbentibus.

Illecebrum verticillatum. *Sp. Pl.* 298. *Huds.*

100. *With.* 267. *Dicks. H. Sicc. fasc.* 12. 13.

Fl. Dan. t. 335.

Corrigiola. *Raii Syn.* 160.

Polygala repens. *Ger. em.* 563.

Angl. Whorled Knotgrass.

In palustribus, udis et humidioribus pascuis, Cornu-
biæ et Devonix.

About Pensans. *F. Borone.* In Devonshire not
uncommon. *With.*

Perennis. Fl. Julio.

Radix fibrosa, subrepens. Caules herbacei, procumben-
tes, ramosi, teretes, foliosi, glabri. Folia opposita, vix
petiolata, ovato-spatulata, integerrima, carnosa, glabra.
Stipulæ intrafoliaceæ, scariosæ, lacerae. Flores nivei,
verticillati, numerosi. Calyx aristatus, persistens.

110. GLAUX. *G. Pl.* 114. *Juss.* 333.

Cal. monophyllus, campanulatus, coloratus, in-
ferus. *Cor.* nulla. *Caps.* unilocularis, quin-
quevalvis, pentaîperma.

*mariti-
ma.*

1. GLAUX.

Glaux maritima. *Sp. Pl.* 301. *Huds.* 101.

With. 268. *Engl. Bot.* t. 13. *Fl. Dan.*

t. 548. *Raii Syn.* 285.

G. exigua maritima. *Ger. em.* 562.

Polygonum maritimum, longiùs radicum no-
stras. *Raii Syn.* 161? *Conf. Pluk.* t. 53. f. 3.

Pet. H. Brit. t. 10. f. 6.

Alfine bifolia, fructu Coriandri, radice genicu-
latâ. *Loef. Pruff.* 13. t. 3.

Angl. Black Salt-wort.

In maritimis cœnosis copiosè.

Perennis.

Perennis. Fl. Junio, Julio.

Radix e fibris longis, tortuosis, subcarnosis. Caulis erectus, palmaris, subramosus, teres, dense foliosus. Folia opposita, oblonga, integerrima, glabra, subtus pallidiora. Flores axillares, solitarii, subseffiles, incarnati.

III. THESIIUM. *G. Pl.* 114. *Juss.* 75.

Gærtn. t. 86.

Cal. monophyllus, superus, cui stamina inserta.

Cor. nulla. *Semen* unicum, corticatum.

I. THESIIUM spicâ ramosâ, bracteis ternis, foliis *linophyllum*.
linearis-lanceolatis, tubo calycino brevissimo. *lum.*

Thesium linophyllum. *Sp. Pl.* 301. *Huds.* 101.

With. 268. *Relb.* 99. *t.* 1. *Sibth.* 414. *Engl.*

Bot. t. 247. *Dickf. H. Sicc. fasc.* 11. 5.

Linaria adulterina. *Raii Syn.* 202. *Ger. em.*

555.

Angl. Bastard Toad-flax.

In pascuis clatis cretaceis.

In Cambridgeshire frequent; also in Suffolk, Dorsetshire, and near Shouldham in Norfolk. *With.* and *Engl. Bot.*

Perennis. Fl. Julio.

Radix lignosa. Herba rigida, glabra. Caules adscendentes, angulati, foliosi, spithamei. Folia sparsa, secunda, linearia, margine scabra. Spicæ terminales, plerumque ramosæ, multifloræ. Pedicelli alterni, uniflori, erecti, bracteis ternis, lanceolatis, acutis, inæqualibus, flori approximatis. Calycis tubus brevissimus, margine patente, acutè quinquefido, denticulato, albo. Stamina brevissima, laciniis calycinis opposita. Stigma emarginatum. Semen ovale, pentagonum, osseum, basi calycis induratâ vestitum.

II 2. VINCA. *G. Pl.* 115. *Juss.* 144. *Gærtn. t.* 117.

Cor. contorta, hypocrateriformis, infera. *Folliculi* duo, erecti. *Semina* nuda.

I. VIN-

- minor.* 1. VINCA caulibus procumbentibus, foliis elliptico-lanceolatis margine glabris, floribus pedunculatis, dentibus calycinis lanceolatis.
Vinca minor. *Sp. Pl.* 304. *Huds.* 91. *With.* 268. *Relb.* 89. *Sibth.* 79. *Curt. Lond. fasc.* 3. *t.* 16.
V. pervinca minor. *Raii Syn.* 268. *Ger. em.* 894.
Angl. Lesser Periwinkle.

In dumetis et ad scpes rariùs.

By Honingham church, Norfolk, on a bank facing the south, plentifully, also in several lanes in that parish, undoubtedly wild.

Perennis. Fl. Maio.

Radix repens, fibris ramosis. Caules teretes, glabri, flagelliformes, foliosi; floriferi erecti, altitudine varii; demum prostrati, radicanes. Folia opposita, petiolata, elliptico-lanceolata, integerrima, nitida, glaberrima, sempervirentia. Flores axillares, alterni, solitarii, pedunculati, erecti, inodori. Calyx campanulatus, quinquefidus, glaber, laciniis lanceolatis, brevibus. Corolla violacea, ore albo.

- major.* 2. VINCA caulibus erectiusculis, foliis ovatis ciliatis, floribus pedunculatis, dentibus calycinis setaccis clongatis.
Vinca major. *Sp. Pl.* 304. *Huds.* 91. *With.* 269. *Relb.* 89. *Sibth.* 79. *Curt. Lond. fasc.* 4. *t.* 19. *Engl. Bot. t.* 514.
Clematis daphnoides major. *Raii Syn.* 268. *Ger. em.* 894.
Angl. Greater Periwinkle.

In dumetis et nemoribus, præcipuè humidis.

Ile of Wight. *With.* In a field at Beckenham in Kent. *Curtis.*

Perennis. Fl. Maio.

Præcedente omnibus partibus major. Caules erecti, demum apice radicales. Folia latè ovata, margine ciliis minutis scabra. Flores cærulæ, nec violacei. Dentes calycini elongati, setacei, sæpius ciliati, apice patentés. Folliculi glaberrimi, nitidi, apice subulati, divaricati. Semina plerumque bina, unum super alterum, oblonga, sulco longitudinali supernè notata.

II. DIGYNIA.

113. HERNIARIA. *G. Pl.* 121. *Juss.* 89.

Cal. quinquepartitus, inferus. *Cor.* nulla. *Stamina* squamis 5 filiformibus interjectis. *Caps.* monosperma, calyce tecta.

1. HERNIARIA glabra herbacea.

Herniaria glabra. *Sp. Pl.* 317. *Huds.* 108.

With. 269. *Engl. Bot. t.* 206. *Fl. Dan.* t. 529.

Herniaria. *Raii Syn.* 160. *Ger. em.* 569.

Angl. Smooth Rupture-wort.

In arenosis vel glareosis.

On the Lizard point, Cornwall. *Ray.* Near Newmarket. *Rev. Mr. Hemsted.*

Perennis. Fl. Julio, Augusto.

Radix fusiformis, sublignosa, ex observationibus D. Stackhouse perennis, nec annua. Caules prostrati, divaricati, alternatim ramosi, teretes, foliosi, glabri. Folia ovata, integerrima, glabra, margine subciliata; inferiora opposita; superiora alterna. Stipulæ membranaceæ. Racemi solitarii, alterni, oppositifolii, foliosi, multiflori. Flores sessiles, congesti, luteo-virides.

Varietas

Varietas β apud Hudsonum mihi videtur *Glaux maritima*,
ab *H. lenticulatâ* Linnæi longè quidem diversa.

hirsuta. 2. *HERNIARIA hirsuta* herbacea.

Herniaria hirsuta. *Sp. Pl.* 317. *Huds.* 109.
With. 270. *Raii Syn.* 161. *Pet. H. Brit.*
t. 10. *f.* 10. *Baub. Hist.* v. 3. 379. *f.* 1.
Angl. Hairy Rupture-wort.

In arenosis rariùs.

At Colney-Hatch near Barnet. *Huds.*

Perennis. Fl. Julio, Augusto.

Prioris varietas, ex sententiâ D. Staekhouse; nec differt
nisi caule pilis patentibus hirsuto, foliorumque paginis
inferioribus, calycibusque etiam densè fetosis.

114. *CHENOPODIUM.* *G. Pl.* 121. *Juss.* 85.
Gærtn. t. 75.

Cal. quinquepartitus, inferus. *Cor.* nulla. *Se-*
men unicum, lenticulare, calyce clauso, pen-
tagono, tectum.

* *Foliis angulosis.*

Bonus 1. *CHENOPODIUM* foliis triangulari-sagittatis
Henri- integerrimis, spicis compositis aphyllis.
cus.

Chenopodium Bonus Henricus. *Sp. Pl.* 318.
Huds. 104. *With.* 270. *Relb.* 102. *Sibth.* 87.
Curt. Lond. fasc. 3. t. 17. *Fl. Dan.* t. 579.

Blitum perenne, Bonus Henricus dictum. *Raii*
Syn. 156.

Bonus Henricus. *Ger. em.* 329.

Wild Spinach. *Pet. H. Brit.* t. 7. *f.* 12.

Angl. Perennial Goosefoot. Mercury Goose-
foot.

In ruderatis at ad vias frequens.

Perennis. Fl. Maio—Augusto.

Radix

Radix ramosa. Caules erectiusculi, teretes, pedales; basi sæpè ramosi; supernè simplices, foliosi, striati, unctuofo-pulverulenti. Folia alterna, petiolata, laxè patentia, hastato-deltaidea, vel sagittata, subrepanda, integerrima, venosa, subtus præcipuè subpulverulenta. Spicæ pedunculatæ, compositæ, multifloræ; inferiores axillares; superiores in spicam communem pyramidatam, aphyllam, digestæ. Flores congesti, virides, pulverulenti. Calyx apice dilatatus, erosus. Styli duo vel tres, subulati, patentes, pubescentes.

“Flores fœminei plurimi, inter hermaphroditos.” *Curtis*.

2. CHENOPODIUM foliis triangularibus denticulatis, racemis confertis strictissimis cauli approximatis longissimis subaphyllis.

Chenopodium urbicum. *Sp. Pl.* 318. *Huds.* 104. *With.* 270. *Lightf.* 148? *Sibth.* 87.

C. erectum foliis triangularibus dentatis, spicis e foliorum alis plurimis longis erectis tenuibus. *Dill. in Raii Syn.* 155.

C. latifolium, minùs ramosum, florum petiolis longissimis, ex foliorum alis confertim enascentibus. *Buxb. Fl. Hallens.* 69. t. i. opt.

Broad-pointed Blite. *Pet. H. Brit.* t. 8. f. 8.

Angl. Upright Goosefoot.

In fimetis et ad margines fossarum secundum vias publicas passim. *Dill.*

My specimen is from Oxfordshire.

Annua. Fl. Augusto, Septembri.

Caulis strictus, simplex vel ramosus, angulatus, rubro lineatus. Folia deltaideo-oblonga, subhastata, basi vix producta, medio præcipuè denticulata (nec profundè dentata), apicem versus integerrima. Racemi axillares, erecti, cauli sæpiùs adpressi, ramosi, ferè aphylli, superiores in paniculam communem, aphyllam, digesti.

“Semina magnitudine ferè *Brassicæ Napi*.” *Curtis sub C. rubro*.

rubrum. 3. **CHENOPODIUM** foliis rhomboideo-triangularibus finuato-dentatis, racemis erectis compositis foliolosis.

Chenopodium rubrum. *Sp. Pl.* 318. *Huds.* 105. *With.* 271. *Relb.* 103. *Sibth.* 88. *Curt. Lond. fasc.* 6. *t.* 21.

Blitum Pes anserinus dictum. *Raii Syn.* 154.

Atriplex sylvestris latifolia, five Pes anserinus: *Ger. em.* 328.

Pes anserinus. *Fuchf. Hist.* 653. *Dalechamp. Hist.* 542. *f.* 2.

Sharp-pointed Blite. *Pet. H. Brit. t.* 8. *f.* 6.

β. Blito Pes anserinus dicto similis. *Raii Syn.* 154. *Angl.* Red Goosefoot.

In finetis et ruderatis.

Annua. Fl. Augusto.

A præcedente differt foliis profundè dentatis, ferè finuatis basi productis, racemis subpatentibus, foliolosis. In apricis magis rubescit.

“Semina minutissima,” *Curtis*; quod etiam in var. β notavit *Dill.* in *Raii Syn.*

murale. 4. **CHENOPODIUM** foliis ovatis nitidis acutis dentatis, racemis ramosissimis cymosis aphyllis.

Chenopodium murale. *Sp. Pl.* 318: *Huds.* 105. *With.* 272. *Relb.* 103. *Sibth.* 88. *Curt. Lond. fasc.* 6. *t.* 20.

Blitum Pes anserinus dictum acutiore folio. *Raii Syn.* 154.

Atriplex sylvestris latifolia altera. *Ger. em.* 328.

Thick Shining Blite. *Pet. H. Brit. t.* 8. *f.* 5.

Angl. Nettle-leaved Goosefoot.

In ruderatis et ad vias.

Annua. Fl. Augusto, Septembri.

Caulis ramosissimus. Folia triangulari-ovata, basi producta, sinuato-dentata, dentibus acuminatis, subincurvis.

Racemi

Racemi axillares, subinde etiam terminales, ramosissimi, cymosi, subdivaricati, omnino aphylli. Herba tota plerumque saturatè virens, lucida, ingrati odoris. Semina minutissimè punctulata.

5. CHENOPODIUM foliis cordatis angulato-*hybridum*,
dentatis acuminatis, racemis ramosissimis sub-
cymosis divaricatis aphyllis.

Chenopodium hybridum. *Sp. Pl.* 319. *Huds.*

105. *With.* 272. *Relb.* 105. *Sibth.* 89.

Abbot. 55. *Curt. Lond. fasc.* 4. t. 23.

C. Stramonii folio. *Dill. in Raii Syn.* 154.

Vaill. Paris. 36. t. 7. f. 2.

Maple Blite. *Pet. H. Brit.* t. 8. f. 7.

Angl. Maple-leaved Goosefoot.

In ruderalis humidioribus, minùs frequens.

In Battersea fields. *Curtis.* Near Northfleet plentifully. *Huds.* On the banks of some watery pits beyond Ely. *Sherard.* About Colchester. *Dale.*

Annua. Fl. Augusto.

Caulis gracilis, ramosus. Folia patula, magna, lætè viridia, omnino cordata, (nec basi producta,) acuminata, angulato-dentata, angulis utrinque tribus. Racemi plerumque aphylli, ramosissimi, subcymosi, divaricatospatentes. Herba lævis, foetens. Semina punctis majoribus notata.

6. CHENOPODIUM foliis rhomboideo-ovatis *album*,
erosis posticè integris: superioribus oblongis
integerrimis, feminibus lævibus.

Chenopodium album. *Sp. Pl.* 319. *Huds.* 106.

With. 271. *Relb.* 104. *Sibth.* 88. *Curt.*

Lond. fasc. 2. t. 15.

Blitum Atriplex sylvestris dictum. *Raii Syn.* 154.

Atriplex sylvestris. *Fuchs. Hist.* t. 119.

Frost Blite. *Pet. H. Brit. t. 8. f. 2.*

β. Blitum folio subrotundo *Dill. in Raii Syn. 155.*

Buddle's Round Blite. *Pet. H. Brit. t. 8. f. 4.*

γ. Chenopodium viride. *Sp. Pl. 319.*

C. album β. *Huds. 106.*

Atriplex sylvestris altera. *Ger. em. 326.*

δ. Chenopodium foliis integris racemosum. *Dill. in Raii Syn. 155.*

ε. C. crasso et obtuso Oleæ folio. *Dill. in Raii Syn. 156.*—non vidi.

Angl. White Goosefoot.

In fimetis, cultis et ruderatis passim.

Annua. Fl. Julio, Augusto.

Caulis ramosus, subinde rubicundus. Folia oleaceo-pulverulenta; inferiora subovata, dentato-erosa, posticè integerrima; superiora angustiora, omnino edentula. Racemi ramosi, erecti, serè aphylli. Semina lævissima, nec punctata.

γ gaudet colore viridi, minùsque albo, foliis angustioribus racemis laxioribus; δ foliis omnibus integerrimis.

*ficifoli-
um.*

7. CHENOPODIUM foliis hastato-sinuatis erosis posticè integris: superioribus oblongis integerrimis, feminibus punctatis.

Chenopodium viride. *Curt. Lond. fasc. 2. t. 16.*

C. serotinum. *Huds. 106. Sibth. 88.*

Blitum Ficùs folio. *Dill. in Raii Syn. 155.*

Buddle's Fig Blite. *Pet. H. Brit. t. 8. f. 3.*

β. Chenopodium erectum Chrysanthemi segetum folio. *Dill. in Raii Syn. 155.*—non vidi.

Angl. Fig-leaved Goosefoot.

In fimetis et ruderatis.

About London. *Curtis.*

Annua. Fl. Augusto.

A præcedente differt colore magis viridi, ramorum axillis purpurascens, foliis inferioribus hastatis, at præcipuè

puè (optimè monente Curtisio) feminibus excavato-punctatis. *C. serotinum* Linnæi discrepat caule quintuplò altiore, foliis majoribus, deltoideis, vix hastatis, tempore florendi seriore; femina ejus non vidi.

8. **CHENOPODIUM** foliis omnibus oblongis *glaucum*.
sinuato-repandis subtùs glaucis, racemis aphyllis glomeratis.

Chenopodium glaucum. *Sp. Pl.* 320. *Huds.* 106. *With.* 272.

C. angustifolium laciniatum minus. *Dill. in Raii Syn.* 155.

Atriplex angustifolia laciniata minor. *Baub. Hist.* v. 2. 972.

A. sylvestris secunda. *Tabernæm. Krauter-buch.* 811.

Rand's Oak Blite. *Pet. H. Brit.* t. 8. f. 1.
Angl. Oak-leaved Goosefoot.

In ruderatis, præcipuè arenosis.

About London. *Mr. Pitchford*.

Annua. Fl. Augusto.

Caules diffusi, crassi. Folia oblonga, subindè ovato-oblonga, obtusiuscula, sinuato-repanda, omnia ferè uniformia, supra viridia, subtus glauca. Racemi breves, glomerati, lobati, aphylli. Semina excavato-punctata.

* * *Foliis integris*.

9. **CHENOPODIUM** foliis rhombeo-ovatis *intolidum*.
gerrimis, racemis conglomeratis.

Chenopodium olidum. *Curt. Lond. fasc.* 5. t. 20. *With.* 273.

C. Vulvaria. *Sp. Pl.* 321. *Huds.* 107. *Relb.* 105. *Sibth.* 89. *Lightf.* 149. *Sym. Syn.* 65. *Woodv. Med. Bot.* t. 145.

Blitum foetidum, *Vulvaria dictum*. *Raii Syn.* 156.

Atriplex olida. *Ger. em.* 327.

Stinking Orache. *Pet. H. Brit. t. 7. f. 11.*

Angl. Stinking Goosefoot.

In ruderalis, præcipuè maritimis.

Common about London. *Curtis.* About Yarmouth and Lowestoft towards the sea.

Annua. Fl. Augusto.

Caules diffusi, ramosi, divaricati. Folia pulverulento-canescens, rhombico-ovata, indivisa et integerrima. Racemi axillares et terminales, condensati, lobati. Herba tota imigritèr foet, odore piscium salorum putrescentium perdurabili.

polyspermum. 10. **CHENOPODIUM** foliis ovatis integerrimis, caule prostrato, racemis cymosis divaricatis apyillis.

Chenopodium polyspermum. *Sp. Pl.* 321. *Huds.* 107. *With.* 273. *Relb.* 105. *Sibth.* 89. *Curt. Lond. fasc. 2. t. 17.*

C. Betæ folio. *Raii Syn.* 157.

Atriplex sylvestris, f. *Polysspermon.* *Ger. em.* 325. Allseed Blite. *Pet. H. Brit. t. 7. f. 10.*

Angl. Round-leaved Goosefoot. Allseed Goosefoot.

In ruderalis et fimetis minùs frequens.

Annua. Fl. Julio, Augusto.

Caules undique diffusi, prostrati. Folia integerrima, obtusa, saturatè viridia. Racemi axillares et terminales, cymosi, laxi, multiflori. Semina minutissimè punctata.

maritimum. 11. **CHENOPODIUM** foliis subulatis semicylindricis, floribus axillaribus sessilibus.

Chenopodium maritimum. *Sp. Pl.* 321. *Huds.* 107. *With.* 273. *Relb.* 106. *Fl. Dan.* t. 489. *Dicks. H. Sicc. fasc. 4. 7. Engl. Bot. t. 633.*

Blitum

Blitum Kali minus album dictum. *Raii Syn.*
156.

Kali minus. *Ger. em.* 535.

Angl. Sea Goosetoot. *Sea Blite.*

In littoribus maritimis copiosè.

Annua. Fl. Julio, Augusto.

Caulis erectus, ramosissimus. Folia semicylindrica, acutiuscula, lævia, succosa, salsa. Flores sessiles, glomerati. Bractæ binæ. Semina minutè striata, nitida.

115. BETA. *G. Pl.* 122. *Juss.* 85. *Gærtn. t.* 75.

Cal. quinquepartitus, basi carnosus. *Cor.* nulla.

Semen unicum, reniforme, intra substantiam baseos calycis.

1. BETA caulibus decumbentibus, floribus ge-*mariti-*
minis, laciniis calycinis integerrimis. *ma.*

Beta maritima. *Sp. Pl.* 322. *Huds.* 108. *With*
277. *Engl. Bot. t.* 285.

B. sylvestris maritima. *Raii Syn.* 157.

Sea Beet. *Pet. H. Brit. t.* 8. f. 9.

Angl. Sea Beet.

In maritimis cœnosis.

Perennis. Fl. Augusto.

Radix fusiformis, carnosa, crassa; extus nigra; intus alba.

Caulis plures, decumbentes, sulcati, foliosi, apice ramosi. Folia ovata, repanda, lævia; radicalia maxima, petiolata; caulina alterna, subsessilia, fursum secunda. Flores in racemis terminalibus foliolosis, sessiles, sæpius gemini, virides. Stigmata sæpè terna.

116. SALSOLA. *G. Pl.* 122. *Juss.* 85.

Gærtn. t. 75.

Cal. quinquepartitus, basi capsularis. *Cor.* nulla.

Semen unicum, cochleatum.

Kali. 1. *SALSOLA* herbacea decumbens, foliis subulatis spinosis scabris, calycibus marginatis axillaribus.

Salsola Kali. *Sp. Pl.* 322. *Huds.* 107. *With.* 278. *Dicks. H. Sicc. fasc.* 12. 14. *Fl. Dan.* t. 818. *Woodv. Med. Bot.* t. 143. *Engl. Bot.* t. 634.

Kali spinosum cochleatum. *Raii Syn.* 107.

Tragos Matthioli. *Ger. em.* 1117.

Angl. Prickly Saltwort.

In littoribus maritimis arenosis passim.

Annua. Fl. Julio.

Caules rigidi, ramosissimi, decumbentes, foliosi, pubescentes. Folia alterna, subulata, canaliculata, patentissima, scabra, apice mucronato-spinosa. Flores axillares, solitarii, sessiles, bracteis ad basin ternis, foliaceis, spinosis. Calycis foliola marginata, dilatata, repanda, rubicunda, scariosa, conniventia. Semen turbinatum, cotyledonibus cochleatis.

fruticosa. 2. *SALSOLA* erecta fruticosa, foliis semicylindricis obtusiusculis muticis.

Salsola fruticosa. *Sp. Pl.* 324. *Huds.* 108.

With. 278. *Engl. Bot.* t. 635.

Blitum fruticosum maritimum, Vermicularis frutex dictum. *Raii Syn.* 156.

Cali species, sive Vermicularis marina arborescens. *Baub. Hist.* v. 3. 704.

Angl. Shrubby Saltwort.

In littoribus maritimis, haud frequens.

On the Norfolk coast, discovered by Sir Thomas Brown, M. D. *Ray.* At Wells and Cley. *Mr. Crowe.* Thornham, Norfolk. *Rev. Mr. Sutton.* Near Southwold, Suffolk. *Mr. Woodward.* In Devonshire and Cornwall. *Huds.* At Weymouth. *A. B. Lambert, Esq.*

Frutex.

Frutex. Fl. Julio, Augusto.

Caulis lignosus, erectus, teres, ramosissimus; rami erecti, foliosi. Folia alterna, erectiuscula, semiteretia, carnosa, obtusiuscula, lævia, glaucescentia. Flores axillares, sessiles, solitarii, virides. Bractæ ternæ, concavæ, scariosæ, parvæ.

Frutex elegans, sempervirens, hortorum ornamentum haud ingratum.

117. ULMUS. *G. Pl.* 123. *Juss.* 408.
Gærtn. t. 49.

Cal. quinquefidus, inferus, persistens. *Cor.* nulla. *Capsula* membranacea, compresso-plana, monosperma.

1. ULMUS foliis duplicato-ferratis scabris basi *campestris*, inæqualibus, floribus subsessilibus congestis. *tris*,

Ulmus campestris. *Sp. Pl.* 327. *Huds.* 109.

With. 278. *Relb.* 106. *Sibth.* 86. *Woodv.*

Med. Bot. t. 197.

U. vulgatissima folio lato scabro. *Ger. em.* 1480. *Raii Syn.* 468.

Common Elm *Trec. Hunt. Evel. Sylv. tab. ad p.* 119.

*β. *Ulmus minor*, folio angusto scabro. *Ger. em.* 1480. *Raii Syn.* 469. Hæc est omnium cultarum varietatum mater, ex auctoritate D. Crowe; at e sententiâ *Milleri* vix indigena.

Ulmus, cortex interior. *Pharmac. Lond.*
Angl. Common Elm.

In sepibus, præcipuè *Angliæ australioris*,

β Chiefly about London. *Miller.*

Arbor. Fl. Aprili.

Cortex rimosus, in ramulis suberosus. Folia utrinque scabra, subtus ad venulas villosa. Flores e propriis gemmis, congesti, vix pedunculati, numerosi, fusco-incarnati. *Capsulæ* oblongæ.

Lignum durum, tenax.

2. UL-

- montana*. 2. ULMUS foliis duplicato-ferratis acuminatis basi inæqualibus, floribus pedunculatis ciliatis.
 Ulmus montana. *Willd.* 279. *Bauh. Pin.* 427.
 U. glabra. *Huds. ed.* 1. 95.
 U. campestris. *Fl. Dan. t.* 632. *Relb.* 107 β.
Huds. 129 γ.
 U. effusa. *Willden. Sp. v.* 1. 1325. *Sibth.* 87.
Sym. Syn. 67. *Abbot.* 55.
 U. latifolia. *Cullum.* 96.
 U. folio latissimo scabro. *Ger. em.* 1481. *Raii Syn.* 469.
 β. U. folio glabro. *Ger. em.* 1481. *Raii Syn.* 469.
 ex sententiâ D. Crowe.
 U. glabra. *Cullum.* 97.
Angl. Broad-leaved Elm. Wych Hasel. β Wych Elm.

In sylvis et sepibus, per totam Britanniam.

Arbor. M. Aprili.

Cortex ramulorum æqualis. Folia latiora quam in præcedente, minusque aspera, acuminata. Flores longius pedunculati, laxè patentes. Fructus subrotundus.

Lignum minus solidum.

Arbores hæc optimè illustravit botanicus indefessus et valdè acutus Johannes Goodyer, Johnsoni amicus. Confer *Ger. em.*

118. CUSCUTA. *G. Pl.* 66. *Juss.* 135.
Gærtn. t. 62.

Cal. 4- vel 5-fidus, inferus. *Cor.* monopetala, 4- vel 5-fida. *Capsula* bilocularis, basi circumscissâ. *Sem.* bina.

europæa. 1. CUSCUTA floribus subsessilibus, corollâ fauce nudâ, stigmatibus acutis.

Cuscuta europæa. *Sp. Pl.* 180. *Engl. Bot. t.* 378. *Fl. Dan. t.* 199. *Relb.* 101. *Abbot.* 36.
Dicks. H. Sicc. fasc. 15. 6.

C. major.

C. major. *Bauh. Pin.* 219.
Cassutha. *Fuchsf. Hist. t.* 348.
Angl. Greater Dodder.

In Carduis, Urticis, aliisque parasitica, rariùs.
 Near Aberdeen. *Mr. Alexander Smith.* Cambridge-
 shire. *Rev. Mr. Hemsted.*

Annua. Fl. Augusto, Septembri.

Radix parva, statim marcescens. Caules filiformes, volubiles, parasitici, e cortice alieno pabulum, tuberculis propriis radicauntibus, trahentes, aphylli, glabri. Flores fasciculati, laterales, ferè sessiles, hyalini, rubro tincti, plerumque quinquesidi. Corolla campanulata, fauce omnino nudâ. Stamina fauce inserta, laciniis alterna. Styli divaricati, stigmatibus acutis, nec, ut in *C. americanâ*, capitatis.

2. CUSCUTA floribus sessilibus, flaminibus basi *Epithy-*
 squamâ crenatâ munitis, stigmatibus acutis. *mum.*

Cuscuta Epithymum. *Syst. Veg. ed.* 13. 140.
Engl. Bot. sub p. 378. *Fl. Dan. t.* 427.
Abbot. 36. *Dicksf. H. Sicc. fasc.* 15. 5.

C. europæa. *Engl. Bot. t.* 55. *Hudsf.* 104.
Sp. Pl. 180 β. *Sibth.* 63.

C. major. *Raii Syn.* 281.

C. five Cassuthe. *Ger. em.* 577.

Epithymum, five Cuscuta minor. *Bauh. Pin.*
 219.

Angl. Lesser Dodder.

In Ericis, Thymis, aliisque suffruticis, parasitica frequens.

Perennis? Fl. Augusto.

Priore minor. Caules tortuosi. Flores sessiles, congesti, albi, calyce rubro, sæpiùs quadrisidi. Corolla infundibuliformis, tubo brevi. Stamina fauce inserta, laciniis alterna, squamâ lunatâ, crenatâ, ad singulorum basin, germiui incumbente. Styli elongati, acuti.

119. SWERTIA. *G. Pl.* 125. *Juss.* 142.
Gærtn. l. 114.

Cor. rotata; poris geminis nectariferis ad basin
 singulæ laciniae. *Caps.* supera, unilocularis,
 bivalvis.

perennis. † 1. SWERTIA corollis quinquefidis, foliis radi-
 calibus ovatis.

Swertia perennis. *Sp. Pl.* 328. *Huds.* 102.
With. 280. *Jacq. Fl. Austr. t.* 243.

Gentiana duodecima, punctato flore. *Clus. Hist.*
v. 1. 316.

G. major, flore cærulco punctato. *Morif.*
sect. 12. *t.* 5. *f.* 11.

G. Pennici minor. *Ger. em.* 433.

Angl. Marsh Swertia. Marsh Felwort.

In palustribus alpinis.

Found in Wales by Dr. Richardson. *Huds.*

Perennis. Fl. Augusto.

Formosa planta. Radix e fibris longis, albidis. Caulis
 erectus, simplex, pedalis, subtetragonus, glaber, parùm
 foliosus. Folia omnia ferè radicalia, petiolata, ovata
 vel elliptica, integerrima, obsolete nervosa, glabra. Spica
 terminalis, erecta, pedunculis oppositis, erectis, angu-
 latis, unifloris, basi bracteatis. Bracteæ sessiles, ellip-
 tico-oblongæ, integerrimæ. Flores cinereo-purpurei,
 tristes, inodori. Calycis laciniae subulatæ, patulæ.
 Corolla patens, laciniis ellipticis, acutis, nigro puncta-
 tis, poris nectariferis margine setosis. Stamina subulata.
 Antheræ versatiles. Germen ovatum, compressum.
 Stylus brevissimus, bipartitus. Herba valde amara.
 E speciminibus in Monte Cenisio lectis descriptionem
 confeci; nec indigenam adhuc vidi.

120. GENTIANA. *G. Pl.* 126. *Juss.* 141.
Gærtn. l. 114.

Cor. basi tubulosa; poris nectariferis destituta.
Caps.

Caps. supera, unilocularis, bivalvis, polysperma.

1. GENTIANA corollâ campanulatâ quinquefidâ, floribus pedunculatis, foliis linearibus. *Pneumonanthe.*
Gentiana Pneumonanthe. *Sp. Pl.* 330. *Huds.*
 102. *With.* 280. *Engl. Bot. t.* 20. *Dicks.*
H. Sicc. fasc. 8. 9. *Fl. Dan. t.* 269.
G. palustris angustifolia. *Raii Syn.* 274.
Pneumonanthe. *Ger. em.* 438. *Cord. Hist.*
 162. 2.
Angl. Marsh Gentian. Calathian Violet.

In ericetis turfosis humidiusculis.

On Stratton Strawless heath, near Norwich. *Mr. Stillingfleet.* On Rollesby and Hopton heaths, Norfolk. *Mr. Wigg.* On Leziate heath, Norfolk, abundantly. *Mr. Crowe.* On Hounslow heath sparingly; but plentifully on Wareham common, Dorset. *Rev. Dr. Goodenough.* In the northern counties in several places.

Perennis. Fl. Augusto, Septembri.

Radix e fibris simplicibus, flagelliformibus, minutè annulatis. Caules spithamæi, erecti, simplices, foliosi, quadranguli. Folia opposita, linearia, obtusa, integerrima, glabra. Flores pauci, terminales et axillares, pedunculati, erecti, formosi, campanulati, plicati, saturatè cyanei, plicis virescentibus. Antheræ connatæ. Styli recurvi. Herba amara.

2. GENTIANA corollâ quinquefidâ hypocrateriformi crenatâ: laciniis basi appendiculatis, foliis ovatis confertis.

Gentiana verna. *Sp. Pl.* 331. *Engl. Bot.*
t. 493. *Sym. Syn.* 64.
G. n. 644. *Hall. Hist. v.* 1. 286. *Davall.*
Gentianella alpina verna. *Ger. em.* 436. *How.*
Phytolog.

Phytolog. 46. *Merr. Pin.* 45. *Dill. Indic. Pl. dub. in Raii Syn.*

G. alpina verna minor, subrotundo folio, flore cæruleo. *Barell. Ic. t.* 98.

Angl. Spring Gentian.

In montosis sterilibus.

On the mountains betwixt Gort and Galloway, Ireland. *Mr. Heaton.* In Teesdale forest, Durham. *Rev. Mr. Harriman* and *Mr. Oliver.*

Perennis. Fl. Aprili.

Radix repens, fibrosa. Caules breves; basi ramosi, filiformes, nudi; supernè simplices, quadranguli, densè foliosi, uniflori. Folia opposita, conferta, ovata, integerrima, acuta, subindè obtusa et ferè spatulata. Flos terminalis, erectus, cyaneus, tubo albido. Calyx angulis quinque, acutis, undulatis, tubo brevior. Corollæ laciniæ crenatæ, acutiusculæ, subindè acutæ, basi utrinque auriculatæ. Stylus simplex. Stigma bifidum, crenatum.

G. Bavarica Linn. hujus varietas videtur.

nivalis. 3. GENTIANA corollâ quinquefidâ infundibuliformi, angulis calycinis æqualibus acutis, caule multifloro.

Gentiana nivalis. *Sp. Pl.* 332. *Dicks. in Tr. of Linn. Soc. v.* 2. 290. *With.* 280. *Fl. Dan. t.* 17. *Sym. Syn.* 64.

G. n. 647. *Hall. Hist. v.* 1. 287.

G. n. 5. *Ejusd. Enum.* 475. *t.* 7.

G. undecima minima. *Clus. Hist. v.* 1. 316. *Parm.* 291.

G. minima. *Lob. Ic.* 310.

Gentianella fugax minor. *Ger. em.* 437, quoad iconem.

Angl. Small Alpine Gentian.

In alpinis Scotiæ.

On Ben Lawers. *Dickson.*

Annua.

Annua. Fl. Augusto.

Radix fibrosa, parva. Caulis erectus, quadrangulus, ramosus, foliosus; ramis elongatis, erectis, unifloris. Folia elliptica; radicalia acuta. Flores erecti, graciles, pulcherrimè cyanci, extùs pallidè virentes. Calyx tubulosus, quinquedentatus, angulis acutis, vix cariuatis.

4. GENTIANA corollâ quinquefidâ hypocrateriformi fauce barbatâ, laciniis calycinis æqualibus. *Amarella.*

Gentiana Amarella. *Sp. Pl.* 334. *Huds.* 103. *With.* 281. *Relb.* 100. *Sibth.* 85. *Abbot.* 56. *Engl. Bot. t.* 236. *Fl. Dan. t.* 328. *Dicks. Dr. Pl.* 5.

Gentianella fugax autumnalis elatior, Centaurii minoris folio. *Raii Syn.* 275 ?

Gentianellæ species quibusdam, an Cordo Pœumonanthe, aut Gentiana fugax altera Clusii. *Baub. Hist. v.* 3. 526. *f.* 1. *benè.*

β. *G. fugax verna seu præcox.* *Raii Syn.* 275. *Angl. Autumnal Gentian.*

In pascuis calcareis.

In the pastures above the rocks opposite Matlock bath, Derbyshire, plentifully; also on the downs above St. Vincent's rocks, Bristol, and in other parts of England on a limestone or chalky soil.

Annua. Fl. Augusto. β Aprili, Maio.

Radix parva, tortuosa. Caulis erectus, altitudine varius, foliosus, quadrangulus, supernè paniculato-ramosus. Folia ovata, trinervia, sessilia. Pedunculi axillares et terminales, erecti, simplices vel biflori, tetragoni. Calyx femiquinquefidus, laciniis acutis, uniformibus, ferè æqualibus. Corolla calyce duplò longior, tubo cylindrico, albido, limbo quinquefido, patente, purpureo, fauce ciliis erectis, purpureis, clausâ. Stamina inclusa. Styli brèves, ad basin usque divisi. Herba intensè amara. Variat floribus quadrisfidis, calycum laciniis semper uniformibus.

5. GEN-

campestris.

5. GENTIANA corollâ quadrifidâ hypocrateriformi fauce barbatâ, laciniis calycinis exterioribus maximis.

Gentiana campestris. Sp. Pl. 334. Hudf. 103. With. 281. Sibth. 86. Abbot. 56. Engl. Bot. t. 237. Fl. Dan. t. 367.

G. pratensis flore lanuginoso. Raii Syn. 275 ?

Gentianella. Tillandsf. Ic. 46.

Angl. Field Gentian.

In pascuis elatis, vel collibus maritimis, solo calcareo vel glaroso.

On Costesy common near Norwich, and near Cley, Norfolk. In Westmoreland, Yorkshire, Durham, and also in Cornwall, Suffex, &c.

Annua. Fl. Septembri.

Habitus præcedentis, sed humilior, magisque ramosa; præcipuè verò discrepat calyce quadripartito, laciniis duabus interioribus oppositis lanceolatis, angustioribus; exterioribus maximis, ovatis.

UMBELLATÆ.

Flores pentapetali superi dispermi.

A. Involucro universali partialique.

121. ERYNGIUM. *G. Pl. 127. Juss. 226.*

Gærtn. t. 20.

Involucrum polyphyllum. Flores capitati. Receptaculum commune conicum, paleaceum. Semina muricata.

maritimum.

1. ERYNGIUM foliis radicalibus subrotundis plicatis spinosis, capitulis pedunculatis.

Eryngium maritimum. Sp. Pl. 337. Hudf. 109.

With. 283. Dickf. H. Sicc. fasc. 10. 3. Woodv.

Med. Bot. t. 102. Fl. Dan. t. 875.

E. marinum. Raii Syn. 222. Ger. em. 1162.

Eryngium,

Eryngium, radix. Pharmac. Lond.
Angl. Sea Eryngo. Sea Holly.

In littoribus maritimis arenosis.

Perennis. Fl. Julio, Augusto.

Radix repens, flagelliformis. Herba glauca. Caulis erectus, ramosus, teres, striatus, foliosus. Folia reniformi-subrotunda, plicata, lobata, nervosa, dentata, dentibus spinosis, rigida, glabra; radicalia petiolata. Capitula terminalia, conica. Involucra foliis conformia, patentia. Paleæ tricuspidadæ, floribus longiores, spinosæ. Dentes calycinî erecti, spinosi. Flores cærulei.

2. *ERYNGIUM* foliis amplexicaulibus pinnato-laciniatis. *campestre.*

Eryngium campestre. Sp. Pl. 337. Hudsf. 110.

With. 284. Engl. Bot. t. 57. Fl. Dan. t. 554.

Jacq. Austr. t. 155.

E. vulgare. Raii Syn. 222.

E. mediterraneum. Ger. em. 1162.

Eryngium. Fuchsf. Hist. 296.

Angl. Field Eryngo.

In pascuis maritimis, rariùs.

By the ferry from Plymouth over to Cornwall, also on the shore called Friar-Goose near Newcastle upon Tyne. *Ray.* By the old Roman way called Watling-street, opposite Brookhall near Daventry, Northamptonshire, as mentioned by *Ray. Rev. Mr. Wood.*

Perennis. Fl. Julio, Augusto.

Herba præcedente ramosior, tenuior, magisque virens. Folia pinnatifida, laciniis lobatis, et sæpè pinnatifidis, spinosis, decurrentibus. Involucra lanceolata. Flores albido-virides. Paleæ indivisæ.

122. HYDROCOTYLE. *G. Pl.* 127. *Juss.* 226.
Gærtn. t. 22.

Umbella simplex. *Invol.* tetraphyllum. *Petala*
integra. *Fructus* orbiculatus, compressus.

vulga- 1. HYDROCOTYLE foliis peltatis, umbellis
ris. quinquefloris.

Hydrocotyle vulgaris. *Sp. Pl.* 338. *Huds.* 110.
With. 284. *Relb.* 107. *Sibth.* 91. *Curt.*
Lond. fasc. 6. *t.* 19. *Fl. Dan.* *t.* 90. *Raii*
Syn. 222.

Cotyledon palustris. *Ger. em.* 529.

Angl. Marsh Pennywort. White-rot.

In palustribus et inundatis frequentissima.

Perennis. *Fl.* Maio, Junio.

Caules repentes, filiformes. Folia opposita, petiolata,
peltata, horizontalia, subrotunda, crenata, pallidè viri-
dia, glabra, lucida. Pedunculi axillares, solitarii vel
gemini. Bractæ binæ, scariosæ. Umbella parva,
densa, subquinqueflora, e centro sæpè prolifera. Flores
albi, vel rubicundi, petalis acutis.

inunda- 2. HYDROCOTYLE foliis pinnatis incis: sub-
ta. mergis multifido-capillaceis, umbellis quinque-
floris binatis.

Sison inundatum. *Sp. Pl.* 363. *Huds.* 120.
With. 301. *Relb.* 117. *Sibth.* 98. *Engl. Bot.*
t. 227. *Dicks. H. Sicc. fasc.* 3. 3. *Fl. Dan.*
t. 89.

Sium pusillum, foliis variis. *Raii Syn.* 212.

S. minimum umbellatum, foliis variis. *Pluk.*
Phyt. t. 61. *f.* 3.

Least Water Parsnep. *Pet. H. Brit. t.* 26. *f.* 4.

Angl. Floating White-rot.

In fossis, stagnis, et inundatis.

Biennis?

Biennis? Fl. Maio.

Caulēs repentes, natantes, teretes. Folia alterna, petiolis dilatatis, amplexicaulibus, impari-pinnata: foliolis oblongis, trifidis, sæpè pinnatifidis; submersis multifido-capillaceis et decompositis. Pedunculi oppositifolii, bifidi, divaricati. Umbellæ solitariae, quinquefloræ. Involucrum plerumque tetraphyllum. Flores albi, petalis acutis, subæqualibus. Fructus ellipticus, striatus.

123. SANICULA. *G. Pl.* 128. *Juss.* 225.

Gærtn. t. 20.

Umbellæ confertæ, subcapitatae. *Petala* inflexa.

Fructus setis uncinatis muricatus. *Flores* disci abortientes.

1. SANICULA foliis radicalibus simplicibus, floribus omnibus sessilibus. *europæa:*

Sanicula europæa. Sp. Pl. 339. *Huds.* 110.

With. 284. *Relb.* 108. *Sibth.* 91. *Engl. Bot.* t. 98. *Fl. Dan. t.* 293.

S. five Diapensia. Raii Syn. 221. *Ger. em.* 948.

Diapensia. Brunf. Herb. v. 1. 80.

Angl. Wood Sanicle.

In nemorosis passim.

Perennis. Fl. Maio.

Radix e fibris subcarnosis. Caulis sesquipedalis, erectus, teres, sulcatus, nudiusculus, parum ramosus, glaber. Folia petiolata, glabra, profundè septemloba, lobis cuneatis, trifidis, incis, ferratis, ferraturis setiferis. Umbella universalis inæqualis, radiis sæpè ramosis; partiales capitatae, multifloræ. Involucra universalia diphylla, foliacea, pinnatifida; partialia polyphylla, foliis lanceolatis, acutis, integris. Flores sessiles, congesti, albi vel rubicundi; centrales masculi, cæteri foeminei, dentibus calycinis elongatis, acutis.

Petala in floribus foemineis decidua. *D. Wood:*

124. BUPLEURUM. *G. Pl.* 129. *Juss.* 224.
Gærtn. t. 22.

Involucella umbellulis majora, pentaphylla. *Petalata* involuta. *Calyx* obsoletus. *Fruetus* subrotundus, compressus, striatus.

rotundifolium. 1. BUPLEURUM involucris universalibus nullis, foliis perfoliatis.

Bupleurum rotundifolium. *Sp. Pl.* 340. *Huds.* 111. *With.* 285. *Relb.* 108. *Sibth.* 92. *Abbot.* 57. *Engl. Bot. t.* 99. *Dicks. H. Sicc. fasc.* 11. 6.

B. perfoliatum rotundifolium annuum. *Raii Syn.* 221.

Perfoliata vulgaris. *Ger. em.* 536.

Angl. Thorow-wax.

Inter segetes, imprimis solo cretaceo.

Annua. Fl. Julio.

Herba rigens, glabra. Caulis erectus, alternatim ramifusus, teres, foliosus. Folia alterna, perfoliata, ovata, acuta, integerrima, glaucescentia, nervosa. Umbellæ axillares et terminales, solitariae, longius pedunculatae, radiis 6 vel 7, involucri universalis planè destitutæ. Umbellulæ suboctofloræ, involucellis pentaphyllis, ovatis, acutis, inæqualibus. Flores flavi, brevius pedicellati.

tenuissimum. 2. BUPLEURUM umbellis simplicibus, alternis subtrifloris, involucris pentaphyllis subulatis.

Bupleurum tenuissimum. *Sp. Pl.* 343. *Huds.* 111. *With.* 285. *Relb.* 109. *Engl. Bot. t.* 478. *Dicks. H. Sicc. fasc.* 12. 15.

B. minimum. *Raii Syn.* 221. *Column. Ecphr.* 247. f. 2.

Agostana tenuissima. *Bute v.* 8. 299.

Angl. Slender Thorow-wax. Hare's-ear.

In inundatis maritimis cœnosis.

Salt-water ditches at Lynn, Wisbeach, &c.

On the shore near Worthing, Suffex, plentifully.

Mr. T. F. Forster jun.

Annua. Fl. Augusto, Septembri.

Herba rigida, acris, glabra, gracilis, a basi ramosa, ramis erecto-patentibus, flexuosis, foliosis, simplicibus. Folia alterna, lanceolata, angusta, acuta, basi attenuata. Umbellulæ axillares, subsessiles, solitariae, plerumque trifloræ, involucellis subulatis, flores superantibus. Flores flavi. Fructus angulatus.

125. ECHINOPHORA. *G. Pl.* 129. *Juss.* 225.

Involucellum turbinatum, monophyllum, sexfidum. *Flores marginales* masculi, pedicellati; *centralis* foemineus. *Semina* involucello immersa.

† 1. ECHINOPHORA foliolis subulato-spinosis *spinosa*. trifidis integrisve integerrimis.

Echinophora spinosa. *Sp. Pl.* 344. *Huds.* 112.

With. 285. *Turr. Farset.* 7. *Cavan. Ic.* t. 127.

E. maritima spinosa. *Raii Syn.* 220.

Crithmum spinosum. *Ger. em.* 533.

C. secundum. *Camer. Epit.* 273. *Matth. Valgr.* v. 1. 445. *ed. Baubini* 381.

Pastinaca marina. *Lobel. Ic.* 710. *Baub. Hist.* v. 3. 196. *Dalech. Hist.* 1396.

Prickly Sampire. *Pet. H. Brit. t.* 28. f. 6.

Angl. Prickly Sampire, or Sea Parsnep.

In littoribus maritimis arenosis, hodie dubia civis.

Observed by Mr. Lawson at Roosebeck in Low Furness, Lancashire. *Ray.* Between Feversham and Sea Salter. *Blackstone.* Between Whitstable and the Isle of Thanet, by Sandwich, and near

West-chester. *Gerarde*. Not found by any botanist of the present day.

Perennis. Fl. Julio.

Radix fusiformis, longa, carnosa. Caulis erectus, ramosissimus, foliosus, sulcatus. Folia opposita, impari-pinnata, foliolis oppositis, subulatis, canaliculatis, integerimis, scabripusculis, mucronato-pungentibus, subinde trifidis. Umbellæ terminales, magnæ. Involucra foliis conformia. Umbellulæ numerosæ, involucellis etiam spinosis. Flosculi radiantes, albi, rubicundi. Seminum rudimenta duo, quorum alterum plerumque abortit. E speciminibus Italicis descripsi.

Radices esculentæ, diureticiæ, aphrodisiacæ. *Turra*. Gustu Pastinacæ sativæ similes, quamquam nonnihil falsedine affectæ, *Cavanilles*,

126. TORDYLIUM. *G. Pl.* 130. *Juss.* 224.

Gærtn. t. 21.

Involucra longa, indivisa. *Corollæ* radiatæ; *Flores* omnes hermaphroditici. *Fructus* sub-orbiculatus, compresso-planus, margine crenatus.

officina-
le.

† 1. TORDYLIUM involucellis longitudine florum, foliolis ovatis incisissimis crenatis, caule pubescente.

Tordylium officinale. *Sp. Pl.* 345. *Huds.* 112? *Willd.* 286?

T. five Sefeli Creticum minus. *Raii Syn.* ed. 2. 102.

T. Narbonense minus. *Dill. in Raii Syn.* 206, loco tamen erroneo.

T. Apulum minimum. *Column. Ecphr.* 124. f. 1?

Sefeli Creticum minus. *Ger. em.* 1050.

Caucalis minor pulchro semine, sive Bellopii. *Bayh. Hist.* v. 3. p. 2. 84.

Augl.

Angl. Small Hart-wort. *Pet. H. Brit. t.* 24.
f. 6.

In arvis rariùs, dubia civis.

Found by Mr. Doody about Isleworth. *Ray.*
About London. *Petiver.*

Annua. Fl. Junio, Julio.

Caulis erectiusculus, ramosus, foliosus, teres, sulcatus, pilis mollibus, brevibus, deflexis, villosus. Folia impari-pinnata, pilosa, scabra; radicalia longissimè petiolata, foliolis ovatis, crenatis, sublobatis, terminali maximo; caulina alterna, subsessilia, foliolis oblongis, argutè serratis et incis. Umbellæ longiùs pedunculatæ, terminales. Involucra et involucella fetacea, scabra. Flores carnèi, petalis exterioribus maximis, radiantibus.

Distincta species a Tordyllo Apulo Linn. et Jacq. Hort.
t. 53.

2. **TORDYLIUM** umbellis confertis radiatis, *maxi-*
foliolis lanceolatis inciso-ferratis, caule fetis. *num.*
deflexis aspero.

Tordylium maximum. *Sp. Pl.* 345. *With.* 286.

Sibth. 94. *Jacq. Austr. t.* 142. *Dill. in Raii*
Syn. 206.

Scfeli Creticum majus. *Ger. em.* 1050.

Caucalis major. *Chyf. Hist. v.* 2. 201.

C. major femine minùs pulchro hirsuto. *Bauh.*
Hist. v. 3. *p.* 2. 85.

Angl. Great Hart-wort. *Pet. H. Brit. t.* 24.
f. 5.

In arvis et aggeribus rariùs.

Under the hedge on the north side of the Parks,
Oxford. *Dr. Sibthorp* and *Mr. Woodward.*
I have a wild specimen gathered in Oxfordshire,
but in what precise part I know not. Morison
(*Pl. Umbellif.* 40.) mentions it as growing about
London.

Annua. Fl. Junio, Julio.

Caules erecti, tri- vel quadri-pedales, ramosi, foliosi, flexuosi, sulcati, fetis arcuè deflexis undique scabri, intus cavi. Folia impari-pinnata, fetis scabra; caulina ferè omnia foliolis lanceolatis, elongatis, ferrato-incisis; radicalia foliolis ovato-oblongis, lobatis, crenatis. Pedunculi oppositifolii, elongati, asperi. Involucra et involucella setacea, aspera. Umbellulæ centrales subfessiles. Flores albi vel rubicundi, petalis exterioribus paulò majoribus, radiantibus. Fructus fetis undique muricatus, margine crenatus, incrassatus, rubicundus.

127. CAUCALIS. *G. Pl.* 130. *Juss.* 224.

Gærtn. t. 20.

Involucra indivisa, *Corollæ* radiatæ, *Flores* disci masculi, *Fructus* subovatus, striatus, fetis rigidis muricatus.

daucoides. 1. CAUCALIS umbellis trifidis nudis, umbellulis tricarpis, involucellis triphyllis, foliis supradecompositis.

Caucalis daucoides. *Syst. Nat. ed.* 12. 205. *ed.* 14. 276. *Mant.* 351. *Huds.* 112. *With.* 287. *Relb.* 109. *Sibth.* 92. *Abbot.* 58. *Engl. Bot. t.* 197. *Jacq. Austr. t.* 157.

C. leptophylla, *Huds. ed.* 1. 99. nec Linn.

C. tenuifolia, flosculis subrubentibus. *Raii Syn.* 219.

C. albis floribus. *Ger. em.* 1021.

Conium Royeni. *Sp. Pl.* 350.

Echinophora tertia leptophyllum purpurea. *Column. Ecphr.* 96. *t.* 97. *f.* 2.

Fine-leaved Bur-parisley. *Pet. H. Brit. t.* 27. *f.* 8.

Angl. Small Bur-parisley.

In arvis solo cretaceo.

At Marham, Norfolk. *Mr. Crowe.* Carlby, between

tween Stamford and Bourn. *Mr. Woodward.*
Thorp-arch, Yorkshire; *Mr. Aikin. With.* In
Cambridgeshire frequent, also in Oxfordshire.

Annua. Fl. Junio.

Radix fusiformis, parva. Caulis ramosus, flexuosus, divaricatus, foliosus, angulatus, angulis subinde scabriusculis. Folia alterna triplicato-pinnata, foliolis pinnatifidis, glabris, nervis subtus aculeatis. Umbellæ oppositifoliæ, pedunculo foliis longiore, divaricato, sulcato. Radii generales plerumque tres, sæpe cum uno alterove minore et imperfectiore. Involucrum vix ullum. Umbellulæ e floribus pluribus, incarnatis, parum radiantibus, quorum tres plerumque fructum perficiunt. Involucellæ pentaphyllæ, umbellulis breviora. Fructus elliptico-teres, nec compressus, costis utrinque quatuor, fetis porrectis, uncinatis, per singulam costam digestis, minoribus interstinctis, muricatus.

2. CAUCALIS umbellis trifidis involucratis, umbellulis pentacarpis, foliis pinnatis ferratis. *latifolia.*

Caucalis latifolia. *Syst. Veg. ed. 12. 205. ed. 14. 276. Hudf. 113. With. 288. Relb. 110. Abbot. 58. Engl. Bot. t. 198. Jacq. Hort. t. 128.*

C. arvensis echinata latifolia. *Raii Syn. 219. Mill. Ic. t. 85.*

C. secunda. *Tabern. Kreuterb. 290.*

Tordylium latifolium. *Sp. Pl. 345. Hudf. ed. 1. 98.*

Echinophora quarta major platyphyllon purpurea. *Column. Ecphr. 98. t. 97. f. 1.*

Broad Bur-parsley. *Pet. H. Brit. t. 27. f. 6.*

β. *Echinophora laciniata.* *Dill. in Raii Syn. 219. Pet. H. Brit. t. 27. f. 7.*

Angl. Great Bur-parsley.

In arvis solo cretaceo rariùs.

In Cambridgeshire not uncommon.

Annua.

Annua. Fl. Julio.

Caulis erectus, tripedalis, ramosus, angulatus, fetis sparsis porrectis vel adscendentibus asper. Folia alterna, impari-pinnata, pinnis lanceolatis, decurrentibus, oppositis, acutè ferratis, glaucescentibus, scabris. Pedunculi oppositifolii vel terminales, longissimi, scabri. Radii generales tres, rarius quatuor, validi, angulati, asperi. Involuerum e foliis tribus vel quatuor, ovatis, brevibus, margine membranaceis. Umbellulæ e floribus pluribus, rubris, radiantibus, quorum quinque plerumque fertiles sunt. Involucella involucro similia, carinâ subinde muricatâ. Fructus ovatus, fetis purpureis, retrorsum scabris, muricatus.

Anthriscus. 3. CAUCALIS umbellis multifidis confertis, involucro polyphylllo, foliolis pinnatifidis, ramis erectiusculis.

Caucalis Anthriscus. *Huds.* 114. *Wilh.* 288. *Relb.* 111. *Sibth.* 93. *Curt. Lond. fasc.* 6. t. 22. *Fl. Dan.* t. 919. *Lightf.* 155.

C. minor, flosculis rubentibus. *Ger. em.* 1022. *Raii Syn.* 219.

Tordylium Anthriscus. *Sp. Pl.* 346. *Jacq. Austr.* t. 261.

Angl. Upright Hedge Parsley.

In sepibus, et versuris agrorum.

Annua. Fl. Julio.

Radix fusiformis, flavicans. Caulis erectus, ferè tripedalis, foliosus, sulcatus, pilis arctè deflexis scaber, ramosus, ramis alternis, erectiusculis. Folia alterna, bipinnata, pinnis oppositis, pinnatifidis et incis, scabris, terminali elongato. Petiolus communis dilatatus, canaliculatus. Umbellæ terminales, erectæ, multiradiatæ. Umbellulæ congestæ, planæ. Involucrum polyphyllum, foliolis subulatis, scabris, radio longè brevioribus. Involucella longitudine ferè pedicellorum. Flores albi vel rubicundi, parvi, parùm radiantes. Antheræ violaceæ. Styli divaricato-reflexi. Fructus ovatus, parvus, fetis incurvis muricatus, apice violaceus.

4. CAU.

4. CAUCALIS umbellis multifidis confertis, in-*infesta*,
volucro univertali subnullo, foliolis pinnati-
fidis, ramis divaricatis.

Caucalis infesta. *Curt. Fl. Lond. fasc. 6. t. 23.*

C. arvensis. *Huds. 113. With. 288. Relb. 110.*

Sibth. 92. Lightf. 155. Ait. Hort. Kew. v. 1.

334. Abbot. 58.

C. helvetica. *Jacq. Hort. Vind. v. 3. t. 16.*

C. segetum minor, Anthriscus hispidus similis.

Raii Syn. 220.

Scandix infesta. *Linn. Syst. Nat. ed. 12. v. 3.*

732. Herb. Linn.

Angl. Spreading Hedge Parsley.

In arvis et ad vias.

Annua. Fl. Julio.

Radix fusiformis, albida. Caulis subsesquipedalis, erectus,
foliosus, teres, striatus, scabriusculus, ramosissimus, ra-
mis divaricatis, alternis. Folia ferè præcedentis. In-
volucrum universale sæpiùs nullum, subinde mono-
phyllum. Involucella polyphylla. Flores radiantes,
albi, flavescentes, rariùs carnei. Antheræ flaveolæ,
subindè purpurascens. Fructus præcedentis, sed ma-
jor, neque apice violaceus,

5. CAUCALIS umbellis lateralibus simplicibus *nodosa*.
subsessilibus, caule prostrato.

Caucalis nodosa. *Huds. 114. With. 289.*

Relb. 111. Sibth. 93. Engl. Bot. t. 199.

C. nodosa echinato semine. *Raii Syn. 220.*

Tordylium nodosum, *Sp. Pl. 346. Jacq. Austr.*

app. t. 24.

Knotted rough parsley. *Pet. II. Brit. t. 27.*

f. 11.

Angl. Knotted Stone Parsley.

Ad margines agrorum, imprimis solo glareoso vel
calcarco,

Annua.

Annua. Fl. Maio, Junio.

Radix parva. Caules prostrati, ramosi, teretes, foliosi, retrorsum scabri. Folia glaucescentia, bipinnata, pinnulis pinnatifidis et incis. Umbellæ oppositifoliæ, solitariæ, subsessiles, simplices, congestæ, albæ vel rubicundæ. Semina exteriora tantum muricata.

128. DAUCUS. *G. Pl.* 131. *Juss.* 224.

Gærtn. t. 20.

Involucra pinnatifida. *Corollæ* subradiatæ. *Fructus* muricatus.

Carota. 1. DAUCUS fructibus hispidis, petiolis subtus nervosis.

Daucus Carota. *Sp. Pl.* 348. *With.* 289.

Relb. 112. *Sibth.* 93. *Woodv. Med. Bot.*

t. 161. *Fl. Dan. t.* 723. *Mart. Fl. Rust. t.* 82.

D. polygamus. *Gouan. Obs.* 9. *Jacq. Hort. Vind. v.* 3. *t.* 78.

D. vulgaris. *Raii Syn.* 218.

Caucalis carota. *Huds.* 114.

β. *Daucus maritimus lucidus.* *Dill. in Raii Syn.* 218.

γ. *D. maritimus.* *With.* 290.

Daucus sylvestris, semen. *Pharmac. Lond.*

Angl. Wild Carrot.

In pascuis, et marginibus agrorum passim.

Biennis. Fl. Junio, Julio.

Radix fusiformis, lutescens, aromatica. Caulis ramosus, teres, sulcatus, sæpius pilosus. Rami floriferi elongati, erecti, nudi. Folia tripinnata, incisa, pilosa; in maritimis caruosa, lucida, subinde hirsutissima, vel lanata. Umbella e radiis numerosissimis, patentibus, post florescentiam incurvato-coarctatis, antrorsum scabris. Umbellulæ planæ. Involucra pinnatifida. Involucella simplicia, vel trifida. Flores radiantés, albi vel flavescentes;

centes ; centralis plerumque solitarius, atro-sanguineus, abortivus.

129. BUNIUM. *G. Pl.* 132. *Juss.* 223.

Corolla uniformis. *Umbella* conferta. *Fructus* ovatus.

1. BUNIUM involucro polyphyлло, caule basi *Bulbo-*
recto folioso. *casta-*

Bunium Bulbocastanum. *Sp. Pl.* 349. *With.* 291. *num.*
Fl. Dan. t. 220.

Bulbocastanum alterum, caule firmiore. *Barr.*
lc. t. 244.

Bulbocastanon. *Dod. Pempt.* 334.

B. majus. *Ger. em.* 1064.

Angl. Great Earth-nut.

In pascuis rariùs.

In Kensington gardens. *Mr. Wm. Wood, F. L. S.*

Perennis. *Fl. Maio.*

Radix tuberosa, subrotunda. Caulis e basi rectus, teres, alternatim ramosus, foliosus, striatus, glaber. Folia tripinnata, laciniis linearibus, sæpè trifidis. Umbellæ terminales, erectæ, multiradiatæ. Umbellulæ congestæ, albæ. Involucrum polyphyllum, radiis triplò brevius.

2. BUNIUM involucro subtriphyлло, caule basi *flexuos-*
aphyllo attenuato flexuoso. *sum.*

Bunium flexuosum. *With.* 291. *Sym. Syn.* 70.
Sibth. 94. *Abbot.* 60.

B. Bulbocastanum. *Huds.* 122. *Relb.* 118.
Curt. Lond. fasc. 4. *t.* 24.

Bulbocastanum. *Raii Syn.* 209.

Bulbocastanon minus. *Ger. em.* 1064.

Angl. Common Earth nut. Kipper or Pig-nut.

In pascuis solo arenoso vel glareoso vulgaris.

Perennis.

Perennis. Fl. Maio, Junio.

Radix tuberosa, subrotunda. Caulis basi sub terra elongatus, attenuatus, flexuosus, albidus; supernè erectus, ramosus, foliosus, striatus, glaber. Folia radicalia, tripinnata, laciniis linearibus, pinnatifidis et incis, petiolo basi flexuoso, attenuato; caulina subsessilia, triternata. Umbellæ ut in priore. Involucrum monophyllum vel triphyllum, sæpe nullum. Involucella polyphylla.

130. CONIUM. *G. Pl.* 132.

Cicuta. *Juss.* 223. *Gærtn. t.* 22.

Involucella dimidiata, subtriphylla. *Fructus* ovatus, gibbus, utrinque quinque-costatus; costis ante maturitatem crispis. *Petala* æqualia.

maculatum.

1. CONIUM feminibus inermibus, caule ramossissimo nitido maculato.

Conium maculatum. *Sp. Pl.* 349. *Huds.* 115. *With.* 292. *Relb.* 112. *Sibth.* 94. *Curt. Lond. fasc.* 1. *t.* 17. *Woodv. Med. Bot. t.* 22. *Jacq. Austr. t.* 156.

Cicuta. *Rau Syn.* 215. *Ger. em.* 1061. *Lob. Ic.* 732.

Cicuta, herba, flos, semen. *Pharmac. Lond. Angl.* Common Hemlock.

In sepibus, ruderalis, et fimetis frequens.

Biennis. Fl. Junio, Julio.

Radix fusiformis, sæpe ramosa, albida, carnosa. Caulis tri- vel quadri-pedalis, erectus, teres, cavus, sulcatus, glaber, nitidus, purpureo maculatus, supernè præcipuè ramossissimus, foliosus. Folia supradecomposita, incisa, nitida, glabra, deflexo-patentia. Umbellæ et umbellulæ multiradiatæ. Involucrum plerumque polyphyllum, breve, deflexum. Involucella tri- vel tetraphylla, secunda. Petala cordata, inflexa, subæqualia, alba. Fructus ovatus, sulcatus, glaber. Herba narcotica.

131. SELINUM. *G. Pl.* 133. *Juss.* 223.

Gærtn. t. 21.

Involucra reflexa. *Fructus* ovali-oblongus, compresso-planus, in medio striatus. *Petala* inflexo-cordata, æqualia. *Calyx* integer.

1. SELINUM lactescens, radice subsimplici, caule *palustre*. solitario, stylis post florentiam divaricatis, petalis involutis.

Selinum palustre. *Sp. Pl.* 350. *Huds.* 115.

With. 293. *Engl. Bot. t.* 229. *Relb. Suppl.*

1. 11.

S. sylvestre. *Jacq. Austr. t.* 152.

Angl. Marsh Milky Parsley.

In palustribus.

Near Doncaster. *Huds.* In Lancashire. *With.*

Near Whitgift, Yorkshire. *Rev. Mr. Wood.* In

alder swamps near Yarmouth. *Mr. Wigg.* Be-

tween Norwich and Heigham towards the river.

Mr. Pitchford. At Weel Carr and other wet

places near Beverley. *Sir T. Frankland.* Near

Prickwillow bank, Isle of Ely. *Rev. Dr. Good-*

enough.

Perennis. Fl. Julio.

Radix subfusiformis. Caulis solitarius, erectus, quadri-pedalis, cavus, fuleatus, glaber, foliosus, basi purpureus, supernè ramosus. Folia bi- vel tripinnata, glabra, foliolis oppositis, pinnatifidis. Umbellæ terminales, erectæ, planæ, magnæ, multiradiatæ. Involucra et involucella polyphylla, radiis breviora, deflexa, margine membranacea. Petala alba, involuto-cordata, æqualia. Fructus ellipticus, anceps. Herba succo albo, tenaci, amaro, foetido feætet.

132. ATHAMANTA. *G. Pl.* 133. *Juss.* 223.

Fructus ovato-oblongus, convexus, striatus. *Pe-*
8 *cula*

tala inflexo-cordata, subæqualia. *Calyx* integer.

Libano- 1. **ATHAMANTA** foliis bipinnatis planis, um-
tis. bellâ hemisphæricâ, feminibus hirsutis.

Athamanta Libanotis. Sp. Pl. 351. With. 294.

Relh. 113. t. 2. Hudf. ed. 1. 100. Engl. Bot. t. 138.

A. Oreoselinum. Hudf. 115.

Apium petræum seu montanum album. Raii

Syn. 218. Bauh. Hist. v. 3. p. 2. 105.

Angl. Mountain Stone Parsley.

In pascuis elatis calcareis rariùs.

On Gogmagog hills, Cambridgehire. *Ray. Relhan.*

Perennis. Fl. Augusto.

Radix fusiformis, recta, lignosa, acris et amaricans, apice
fibris petiolorum persistentibus coronata. Caulis sub-
bipedalis, erectus, parùm ramosus, glaber, angulatus
atque sulcatus, foliosus. Folia alterna, rariùs opposita,
bipinnata; foliolis sessilibus, oppositis, pinnatifidis, pla-
nis, acutis, venosis, subtùs pallidioribus, venis margi-
neque subhirsutis. Petioli basi dilatati, margine meni-
branacei. Umbellæ erectæ, hemisphæricæ, conden-
satæ, quandoque proliferæ. Involucra et involucella
patenti-deflexa, polyphylla, pilosa. Flores albi, parvi,
uniformes, et regulares. Fructus pubescens. Styli
persistentes, purpurei, divaricati.

133. **PEUCEDANUM.** *G. Pl. 134. Juss. 223.*

Gertn. t. 21.

Fructus ovatus, utrinque striatus, alâ cinctus.

Calyx quinquedentatus. *Involucra* brevissima.

Flores disci abortivi.

officina- 1. **PEUCEDANUM** foliis quinquies tripartitis;
le. foliolis linearibus indivisis.

Peucedanum officinale. Sp. Pl. 353. Hudf. 116.

With. 294.

Peuce-

Peucedanum: *Raii Syn.* 206. *Ger. em.* 1054.

P. minus germanicum. *Baub. Hist.* v. 3. p. 2. 36.

Peucedanus. *Fuchf. Hist.* 599.

Hog's Fennel: *Pet. H. Brit.* t. 24. f. 7.

Angl. Sea Sulphur-wort.

In palustribus maritimis.

In ditches near Shoreham, Suffex, and at Walton near Harwich, Essex. *Ray.* About a quarter of a mile below Feversham, by the river side. *Sheppard.* Never found at Yarmouth nor Cley; *Œnanthe pimpinelloides* was mistaken for it. *Mr. Rose.*

Perennis. Fl. Junio, Julio.

Radix fusiformis, resinosa, odore sulphureo. Herba glabra. Caulis erectus, teres, striatus, ramosus, foliosus. Folia strigosa, quintuplicato-tripartita, foliolis linearibus, acutis, indivisis, integerrimis. Umbellæ magnæ, radiis numerosissimis, exterioribus elongatis. Involucrum e foliolis paucis, brevibus, setaceis. Involucella pedicellis capillaribus duplò breviora. Flores lutei. Dentes calycini acuti, inflexi. Petala æqualia, incurva. Styli deflexi. Fructus ellipticus, utrinque sulcatus.

2. PEUCEDANUM foliolis pinnatifidis: laciniis *Silaus.* oppositis decurrentibus, involucro universali subdiphylo.

Peucedanum *Silaus.* *Sp. Pl.* 354. *Huds.* 116.

With. 294. *Relb.* 114. *Sibth.* 95. *Jacq.*

Austr. t. 15. *Mart. Fl. Rust.* t. 128.

Seseli pratense nostras. *Raii Syn.* 216.

Saxifraga Anglicana, facie *Seseli pratensis.* *Ger. em.* 1047.

Meadow Saxifrage. *Pet. H. Brit.* t. 28. f. 5.

Angl. Meadow Sulphur-wort.

In pratis et pascuis humidioribus.

Perennis. Fl. Augusto.

Radix fusiformis. Caulis bipedalis, erectus, ramosus, teretiusculus, sulcatus, glaber, foliosus. Folia glabra, compositione ac divisione varia, plerumque supradecomposita, foliolis oppositis, vel integris, vel pinnatifidis laciniis decurrentibus, acutis. Umbellæ radiis exterioribus elongatis. Involucrum diphyllum, subinde monophyllum, exiguum. Involucella polyphylla. Flores luteoli. Dentes calycini obsoleti. Fructus oblongus, undique sulcatus. Dubii generis planta.

134. CRITHMUM. *G. Pl.* 134. *Juss.* 223.

Fructus ovalis, compressus, striatus. *Flores* æquales. *Calyx* integer.

maritimum.

1. CRITHMUM foliolis lanceolatis carnosiss.

Crithmum maritimum. *Sp. Pl.* 354. *Huds.* 117.

With. 295. *Jacq. Hort. Vind. t.* 187. *Dicks. H.*

Sicc. fasc. 12. 16.

C. maritimum. *Raii Syn.* 217. *Ger. em.* 533.

Crethamus. *Cord. Hist.* 201. 1.

Sampire. *Pet. H. Brit. t.* 24. *f.* 8.

Angl. Sea Sampire.

In clivis maritimis.

Perennis. Fl. Augusto.

Radix ramosa, tortuosa, multiceps. Caules subpedales, erectiusculi, teretes, foliosi, parum ramosi. Folia triterinata; foliolis lanceolatis, acutis, carnosiss, integerimis, glabris, glaucescentibus, recurvato-adscendentibus. Umbellæ hemisphæricæ, congestæ, flavescentes. Involucra et involucella ovato-lanceolata, concava, patentia. Petala inflexa, basi lata. Fructus ovali-oblongus, subspongiosus, glaber.

135. HERACLEUM. *G. Pl.* 137. *Juss.* 222.

Sphondylium. *Gærtn. t.* 21.

Fructus ellipticus, emarginatus, compressus, striatus, margine dilatatus. *Flores* radiant.

Petala

Petala inflexo-emarginata. *Involucrum* caducum.

1. HERACLEUM foliis pinnatis : foliolis pinnatifidis incisiferratis. *Sphondylium*.

Heracleum Sphondylium. *Sp. Pl.* 358. *Huds.*

117. *With.* 294. *Relb.* 114. *Sibth.* 95.

Sphondylium. *Raii Syn.* 205. *Ger. em.* 1009.

S. vulgare hirsutum. *Morif. sect.* 9. t. 16. f. 1.

Acanthus germanica. *Fuchsf. Hist.* 53.

Cow Parsnep. *Pet. H. Brit.* t. 24. f. 1.

- β. *Huds.* 117. *With.* 296. var. 2.

Heracleum longifolium. *Jacq. Austr.* t. 174?

Sphondylium majus aliud, laciniatis foliis. *Raii Syn.* 205.

S. hirsutum minus, foliis tenuiter laciniatis, semine lato. *Pluk. Phyt.* t. 63. f. 3.

Jagged Cow Parsnep. *Pet. H. Brit.* t. 24. f. 2.

Angl. Common Cow Parsnep.

In sepibus, agrorum marginibus, et pratis subhumidis.

Biennis. Fl. Julio.

Radix fusiformis. Caulis quadripedalis, erectus, ramosus, foliosus, angulatus, sulcatus, scaber. Folia ternata vel pinnata, variè pinnatifida et incisa, subtùs pubescentia. Stipula petiolo adnata, ventricosa, extùs hirsuta. Umbellæ planiusculæ, albæ, interdum rubicundæ, radiatæ. Involucra et involucella lanceolata, decidua. Petala inflexo-cordata. Fructus latissimus, glaber, utrinque trinervis.

Foliola in β elongata, angustata, subtùs scabra minùsque tomentosa. Diverfa ab *H. angustifolio* Linnæi.

2. HERACLEUM foliis pinnatis, foliolis linearibus basi cruciatis margine scabris. *angustifolium*.

Heracleum angustifolium. *Linn. Mant.* 57.

Angl. Narrow-leaved Cow Parsnep.

In nemorosis.

In a wood at Castle Howard, Yorkshire. *Mr. Teesdale.*

Biennis? Fl. Julio.

Foliola linearia, longissima, decurrentia, subintegerrima, supra lævia, margine scabra; inferiora basi lobata et quasi cruciata, lobis divaricatis. Flores haud radiati, virescentes.

Sine dubio a præcedente distincta.

136. MEUM. *Tourn. t. 165. Gærtn. t. 23.*

Fructus elliptico-oblongus, tricoatus utrinque.

Involucra incisa: universale sæpè abortivum:

partialia dimidiata. *Petala* inflexa, integra.

Calyx obsoletus.

athamant- 1. MEUM foliolis omnibus setaceo-multipartitis.
ticum. Meum athamanticum. *Jacq. Austr. t. 303.*
Gærtn. v. 1. 105.

Meum. *Raii Syn. 207. Ger. em. 1052. Camer. Epit. 7. Matth. Valgr. v. 1. 22.*

Athamanta Meum. *Sp. Pl. 353. Hudf. 116.*

Dicks. H. Sicc. fasc. 11. 7.

Æthusa Meum. *Syst. Veg. ed. 13. 237. ed. 14. 287.*

Ligusticum Meum. *Crantz. Fasc. 3. 82.*

Daucus creticus. *Fuchs. Hist. 231.*

Angl. Spiguel. Meum. Bald-money.

In pascuis montosis.

In Westmoreland, the north of Yorkshire, and Merionethshire. *Ray.*

Perennis. Fl. Maio.

Radix lignosa, crassa, aromatica, apice fibroso-capillata.

Caules erecti, subsimplices, nudiusculi, striati. Folia bipinnata, foliolis setaceo-multifidis, tenuissimis, acuminatis, glabris. Stipulæ dilatatæ, membranacæ.

Umbellæ

Umbellæ erectæ, multiradiatæ. Umbellulæ patulæ, multifloræ. Involuerum paucifolium, foliis linearibus, apice plerumque tripartitis, sæpe monophyllum, vel omnino nullum. Involucella polyphylla, linearia, integra vel apice pinnatifido-incisa. Flores albi vel rubicundi, uniformes. Calyx obsoletus. Petala lanceolata, apice inflexa. Fructus parum compressus, costis acutis.

Genus Ligustico proximum, & forsitan vix distinctum.

137. **LIGUSTICUM.** *G. Pl.* 137. *Juss.* 222.
Gærtn. t. 85.

Fructus oblongus, tricosatus utrinque. *Corollæ* æquales. *Petala* involuta, integra. *Calyx* quinque-dentatus.

1. **LIGUSTICUM** foliis biternatis.

scoticum.

Ligusticum scoticum. *Sp. Pl.* 359. *Huds.* 117.

With. 296. *Lightf.* 159. *Fl. Dan.* t. 207.

L. scoticum Apii folio. *Raii Syn.* 214.

L. humilius scoticum a maritimis. *Pluk. Phyt.*
t. 96. f. 2.

Imperatoriae affinis umbellifera maritima scotica.

Sibb. Scot. p. 2. 32. t. 12. f. 3. pessima.

Seseli maritimum scoticum humile foliis *Imperatoriae*. *Herm. Parad.* 227. t. 227.

Scotch Parsley. *Pet. H. Brit.* t. 26. f. 11.

Angl. Scottish Lovage.

In maritimis Scotiæ.

About the Frith of Forth in many places, also on the western coast.

Perennis. Fl. Julio.

Caules erecti, supernè parum ramosi, teretes, striati, glabri, foliosi. Folia alterna, biternata, superiora tantum ternata; foliola subrhombea, acuta, anticè serrata vel incisa, glabra, subcarnosa. Petiolus communis basi dilatatus, membranaceus, nervosus, amplexicaulis. Um-

bellæ terminales, solitariae, erectæ, multiradiatæ, glaberr. Radii angulati. Involucrum e foliolis quatuor vel quinque, inæqualibus. Involucella polyphylla, linearia, subæqualia. Flores regulares, æquales, parvi, albi, vel extus rubicundi. Calycis foliola deltoidea, acuta. Petala involuta. Fructus oblongus, costis membranaceis. Herba virosè aromatica.

*cornubi-
ense.*

2. **LIGUSTICUM** foliis radicalibus decompositis incisis: caulinis ternatis lanceolatis integerrimis, feminum sulcis obsoletis.

Ligusticum cornubiense. Sp. Pl. 359. Hudf. 118.

With. 297. Sm. Ic. pict. plant. rar. t. 11.

Smyrnum tenuifolium nostras. Dill. in Raii Syn. 209. t. 8.

Cornwal Saxifrage. Pet. H. Brit. t. 26. f. 9.

Angl. Cornish Lovage.

In dumetis et ad sepes Cornubiæ.

Perennis. Fl. Julio.

Radix fusiformis, resinâ flavâ scatens, apice contracta. Caulis bi- vel tri-pedalis, erectus, ramosus, teres, striatus, scabriusculus, basi purpurascens, parùm foliosus. Folia radicalia petiolata, ternato-supradecomposita, latiora quam longa, foliolis variè incisis, subcuneatis, acutis, reticulato-venosis, glabris, margine nervisque scabris. Petioli scabri, canaliculati, foliis longiores. Folia caulina solitaria ad omnem caulis divisionem, parva, ternata, petiolata; foliolis sessilibus, lanceolatis, acutis, vix lobatis, integerrimis; summa simplicia. Umbellæ terminales, erectæ, multiradiatæ. Radii teretes, glabri, striati. Involucrum suboctophyllum, patens, persistens, foliolis lanceolatis, acutis, integris, nervosis, umbellâ longè brevioribus. Involucella involucro conformia, sed minora. Flores albi, regulares et uniformes. Foliola calycina deltoidea, parva. Petala oblonga, acuta, involuta, posticè canaliculata. Fructus ovato-oblongus, utrinque striatus, costis parùm exstantibus. Semina ovata, nigra,

138. ANGELICA. *G. Pl.* 138. *Juss.* 222.
Gærtn. t. 85.

Fructus subrotundus, solidus, utrinque trialatus.
Calyx quinquedentatus. *Corollæ* æquales.
Petala incurva. *Styli* reflexi.

- * 1. ANGELICA foliorum impari lobato. *Archangelica.*
Angelica Archangelica. Sp. Pl. 360. *With.* 297. *gelica.*
Fl. Dan. t. 206. *Woodv. Med. Bot. t.* 50.
A. fativa. Ger. em. 999. *Dill. in Raii Syn.* 208.
Angelica. Pet. H. Brit. t. 24. *f.* 9.
Angelica, radix, caulis, folium, semen. Pharmac.
Lond.
Angl. Garden Angelica.

In aquosis rariùs, an reverà indigena ?
 At Broadmoore, about seven miles north-west from
 Birmingham. *With.*

Biennis. Fl. Septembri.

Radix crassa, carnosa, resinosa. *Caulis* erectus, quinque-
 pedalis, ramosus, cavus, teres, glaber, sulcatus. *Folia*
 ternata, pinnata; foliolis ovato-lanceolatis, acutis, in-
 cisis et acutè ferratis, glabris, subdecurrentibus, impari
 tripartito. *Petiolus* ad basin membranaceus, nervosus,
 maximè dilatatus, ventricosus. *Umbellæ* globosæ,
 multiradiatæ. *Umbellulæ* densæ, hemisphæricæ. In-
 volucrum e foliolis paucis, linearibus, deciduis. In-
 volucella suboctophylla, lineari-lanceolata, brevissima.
Calyx exiguus. *Petala* ovata, inflexa, albido-viridia.
Fructus elliptico-subrotundus, compressus, acutè cos-
 tatus.

2. ANGELICA foliolis æqualibus ovatis ferratis. *sylvestris.*
Angelica sylvestris. Sp. Pl. 361. *Huds.* 118. *tris.*
With. 298. *Relb.* 115. *Sibth.* 96. *Raii Syn.*
 208. *Ger. em.* 999. *Woodv. Suppl. t.* 265.
Water Angelica. Pet. H. Brit. t. 24. *f.* 10.
Angl. Wild Angelica.

In aquosis et ad rivulos passim.

Perennis. Fl. Julio.

Caulis erectus, quadripedalis, teres, lævis, rore cæsius conspicuus, foliosus. Folia bipinnata, subglaucæ; foliolis ovatis, distinctis, acutè ferratis, ferraturis mucronatis. Umbellæ hemisphæricæ, multiradiatæ. Umbellulæ densæ. Involucrum subdiphyllum, angustissimum, sæpe nullum. Involucella polyphylla, fetacea. Calyx vix conspicuus. Petala ovata, inflexa, carneæ. Fructus parvus.

139. SIUM. *G. Pl.* 138. *Juss.* 222. *Gærtn. t.* 23.

Fructus subovatus, compressus, striatus. *Involucra* polyphylla. *Petala* cordata, uniformia.

latifolium. 1. SIUM foliis pinnatis; foliolis oblongo-lanceolatis æqualiter ferratis.

Sium latifolium. *Sp. Pl.* 361. *Huds.* 118.

With. 298. *Relb.* 115. *Sibth.* 96. *Engl.*

Bot. t. 204. *Fl. Dan. t.* 246. *Jacq.*

Austr. t. 66.

S. latifolium foliis variis. *Raii Syn.* 211.

S. majus latifolium. *Ger. em.* 256.

Angl. Broad-leaved Water Parsnep,

In rivulis, fossis, et paludosis.

Perennis. Fl. Julio, Augusto.

Radix repens. Caules erecti, angulati, cavi, glabri, parum ramosi, foliosi. Folia alterna, impari-pinnata; foliolis plerumque septenis, oppositis, sessilibus, ovatis vel lanceolatis, æqualiter ferratis, venosis, glabris; immerfis sæpe laciniatis. Umbellæ solitariae, terminales, et laterales, erectæ, magnæ, planiusculæ. Involucrum e foliolis plurimis, lanceolatis, inæqualibus, quandoque lobatis vel ferratis. Involucella ovata, brevissima. Calyx quinquedentatus, parvus. Petala acuminate, inflexo-cordata, alba, uniformia, sed magnitudine parum inæqualia. Fructus parvus, striatus. Planta acris; radices imprimis venenatæ æstimantur.

2. SIUM foliis pinnatis; foliolis inæqualiter lobatis ferratisque, umbellis pedunculatis oppositifoliis, caule erecto. *angustifolium.*

Sium angustifolium. *Sp. Pl.* 1672. *Huds.* 119.

With. 299. *Relb.* 116. *Sibth.* 96. *Engl.*

Bot. t. 139. *Jacq. Austr. t.* 67.

S. nodiflorum. *Fl. Dan. t.* 247.

S. erectum. *Huds. ed. 1.* 103.

S. five Apium palustre, foliis oblongis. *Raii Syn.* 211.

S. majus angustifolium. *Ger. em.* 256.

Angl. Narrow leaved Water Parsnep.

In fossis et rivulis haud infrequens.

Perennis. *Fl.* Julio, Augusto.

Præcedente minor et debilior. Radix repens. Caulis teres, striatus, ramosus, foliosus. Folia impari-pinnata; foliolis numerosis, glaberrimis, variè incisis et inæqualiter ferratis; impari sæpiùs trifido. Umbellæ oppositifoliæ, solitariae, pedunculatae, erectæ, hemisphæricæ. Involucrum polyphyllum, deflexum, quandoque incisum. Involucella ovata, plerumque integra. Calyx vix conspicuus. Petala alba, uniformia et ferè æqualia. Fructus parvus.

3. SIUM foliis pinnatis; foliolis ovatis æqualiter ferratis, umbellis sessilibus oppositifoliis, caule procumbente. *nodiflorum.*

Sium nodiflorum. *Sp. Pl.* 361. *Huds.* 119.

With. 299. *Relb.* 116. *Sibth.* 96. *Woodv.*

Med. Bot. t. 182. *Abbot.* 62. *Eng. Bot. t.* 639.

S. umbellatum repens. *Raii Syn.* 211.

S. aquaticum repens et procumbens ad alas floridum. *Morif. sect. 9. t. 5. f. 3.*

Creeping Water Parsnep. *Pet. H. Brit. t.* 26. *f. 3.*

Sium, herba. *Pharmac. Lond.*

Angl. Procumbent Water Parsnep.

In fossis et rivulis frequens.

Perennis. Fl. Julio, Augusto.

Radix repens. Caules procumbentes vel natantes, ramosi, teretes, substriati, foliosi. Folia impari-pinnata; foliolis quinque ad novem, sessilibus, ovatis, concinnè et æqualiter serratis. Umbellæ oppositifoliæ, solitariae, subsessiles. Umbellulæ quinque ad septem, divaricatæ, subduodecimfloræ. Involucrum monophyllum, sæpe nullum. Involucella ovata, concava, longitudine pedicellorum. Calyx obsoletus. Fructus ovatus.

- repens.* 4. *SIUM* foliis pinnatis; foliolis subrotundis dentato-incisis, umbellis pedunculatis oppositifoliis, caule repente.

Sium repens. *Suppl.* 181. *Jacq. Austr. t.* 260.

With. 300. *Sibth.* 97. *Abbot.* 63.

Angl. Creeping Water Parsnep.

In paludosis et inundatis turfosis.

On Bullington green, and Cowley bottom near Oxford. *Sibth.* On the bank of a pond at Fisherrow, five miles from Edinburgh, and in abundance on the moist parts of Guillon Links, East Lothian; Dr. Hope. *With.*

Perennis. Fl. Junio—Augusto.

Caules prostrati, repentes, teretes, striati, foliosi. Folia impari-pinnata; foliolis subrotundis, subinde cuneatis, dentato-incisis et variè lobatis, venosis, terminali tripartito. Umbellæ oppositifoliæ, pedunculatæ. Umbellulæ paucae, divaricatæ, multifloræ. Involucra et involucella polyphylla, ovata, nervosa, deflexa. Calyx obsoletus. Fructus subrotundus, compressus. *H. Linn. fil.*

- verticillatum.* 5. *SIUM* foliolis multifido-capillaribus verticillatis.

Sison verticillatum. *Sp. Pl.* 363. *Huds.* 120.

With. 301. *Lichtf.* 1096. *t.* 35. *Engl. Bot.*

t. 395.

Ænanthe

*Oenanthe millefolii palustris folio. Moris. Umbel-
lif. t. 6. f. 3. Hist. sec. 9. t. 7. f. 10.*

Daucus pratensis. Dalech. Hist. 718.

Angl. Whorled Water Parsnep.

In pratis humidis salis.

Plentiful in the western parts of Scotland and
Wales. *Huds. Lightf.*

Perennis. Fl. Julio, Augusto.

Radix e tuberibus pluribus, napiformibus. Caulis ses-
quipedalis, erectus, parum ramosus, teres, striatus,
glaber, nudiusculus. Folia plerumque radicalia, pin-
nata, multijuga; foliolis oppositis, digitato-multifidis,
laciniis linearibus, subinde dichotomis, acutis, divari-
cato-patentibus, et inde verticillum formantibus. Um-
bellæ terminales, erectæ, multiradiatæ. Umbellulæ
multifloræ, planiusculæ. Involucrum e foliolis quin-
que vel sex, ovatis, deflexis, brevibus. Involucella po-
lyphylla, deflexa. Calyx obsoletus. Petala alba, in-
flexo-cordata, uniformia et ferè æqualia. Fructus
ovato-subrotundus, compressus.

Ex Umbellis multiradiatis, involucro et involucello poly-
phylo, petalisque cordatis, Sium potius quam Sison
mihi videtur.

140. SISON. *G. Pl. 139. Juss. 221.*

Fructus ovatus, striatus. *Involucra* subtetra-
phylla. *Petala* lanceolata, inflexa.

1. SISON foliis pinnatis, umbellis erectis subqua- *Amo-*
driradiatis. *num.*

Sison Amomum. Sp. Pl. 362. Huds. 119.

With. 300. Relb. 116. Sibth. 97. Abbot. 63.

Jacq. Hort. Vind. v. 3. t. 17.

S. five officinarum Amomum. Bauh. Hist. v. 3.

p. 2. 107.

Sium aromaticum Sison Off. Raii Syn. 211.

Petroselinum. Fuchs. Hist. 655.

P. ma-

P. macedonicum Fuchsi. *Ger. em.* 1016.

Angl. Hedge Honewort. Bastard Stone Parsley.

In lutosis vel calcareis humidiusculis.

Annua. Fl. Augusto.

Radix ramosa. Caulis erectus, tripedalis, ramosissimus, teres, striatus, glaber, foliosus, subflexuosus. Folia impari-pinnata; inferiora foliolis 7 vel 9, ovatis, incisis, serratis, subinde compositis; superiora foliolis angustioribus, profundius incisis et sæpe pinnatifidis. Umbellæ numerosæ, terminales, solitariae, florescentes erectæ, radiis plerumque quatuor, divaricatis, centrali brevioribus. Involucrum e foliolis duobus ad quatuor, lanceolatis brevibus. Umbellulæ subquadrifloræ, pedicellis inæqualibus. Involucella plerumque tetraphylla, parva. Flores ferè regulares, albi. Calyx obsoletus. Fructus ovatus, costatus, aromaticus. Herba et semina immatura gaudent odore acri, viroso, mihi odiosissimo.

segetum. 2. *SISON* foliis pinnatis; foliolis subrotundis numerosis, umbellis cernuis amorphis.

Sison segetum. *Sp. Pl.* 362. *Huds.* 120. *With.*

300. *Relb.* 117. *Sibth.* 97. *Abbot.* 63.

Jacq. Hort. Vind. v. 2. t. 134. *Engl. Bot.*

t. 228. *Dicks. H. Sicc. fasc.* 12. 17.

Sium arvense sive *segetum.* *Raii Syn.* 211.

Selinum Sii foliis. *Goodyer in Ger. em.* 1018.

Angl. Corn Honewort.

In arvis, solo lutofo vel calcareo, subhumido.

About Mapledurham, Hants. *Goodyer.* Binham, Norfolk. *Mr. Crowe.* Near Kelmarsh, Northamptonshire. *Mr. Hanbury.* At Walthamstow. *Mr. Forster.*

Annua. Fl. Augusto.

Radix parva, sublignosa, quandoque biennis. Caules erectiusculi, ramosi, teretes, striati, juncei, foliosi. Folia impari-pinnata, foliolis numerosis, ovato-subrotundis,

dis, ferratis, vix incisfis, terminali trifido. Umbellæ terminales, radiis plerumque binis vel ternis, valdè inæqualibus. Umbellulæ nutantes, paucifloræ, pedicellis maximè irregularibus. Involucrum diphyllum, lanceolatum. Involucella subpentaphylla, ovata, acuta, concava. Flores regulares, albi vel carnei. Calyx quinquedentatus. Antheræ purpurascens. Fructus oblongo-ovatus, duplicato-costatus, aromaticus.

141. CENANTHE. *G. Pl.* 140. *Juss.* 221.
Geertn. t. 22.

Flosculi diffformes; *disci* sessiles, steriles. *Fructus* calyce styliqûe coronatus, cortice suberoso.

1. CEnanthe stolonifera, foliis caulinis pinnatis fili-*fistulosa*.
formibus fistulosis, involucro universali sub-
nullo.

CEnanthe fistulosa. *Sp. Pl.* 365. *Huds.* 121.
With. 301. *Relb.* 117. *Sibth.* 98. *Engl.*
Bot. t. 363. *Fl. Dan. t.* 846.

CEn. aquatica. *Raii Syn.* 210.

Filipendula aquatica. *Ger. em.* 1060.

Water Dropwort. *Pet. H. Brit. t.* 25. *f.* 5.

2. CEnanthe aquatica triflora, caulibus fistulosis.
Morif. sect. 9. t. 7. *f.* 8. *Raii Syn.* 210.

Small Water Dropwort. *Pet. H. Brit. t.* 25.
f. 6.

Angl. Common Water Dropwort.

In fossis, stagnis, et palustribus vulgaris.

Perennis. Fl. Julio, Augusto.

Radix tuberosa, stolonifera. Caules erecti, tripedales, ramosi, teretes, fistulosi, subinflati, sulcati, glabri, foliosi, glauci. Folia radicalia immersa, bipinnata; foliolis planis, cuneatis, lobatis: caulina alterna, pinnata; foliolis petiolisqûe cylindricis, cavis. Umbellæ alternæ, longius pedunculatæ, erectæ; primordiales triradiatæ; cæteræ multiradiatæ. Umbellulæ densæ, multifloræ. Involucrum monophyllum, sæpe nullum. In-
volucella

volucella polyphylla, lanceolata, membranacea. Dentes calycini subulati, acuti, patentes. Flores carnei, radiantes; centrales abortivi. Petala inflexo-cordata. Fructus ovato-cylindricus, suberosus, calyce coronatus, stylis persistentibus, elongatis, rectis, rigidis, capitatis, aristatus.

pimpinelloides. 2. **CENANTHE** foliolis radicalibus cuneatis fissis; caulinis linearibus integerrimis elongatis, involucri polyphylo lineari.

Cenante pimpinelloides. *Sp. Pl.* 366. *Huds.* 121. *With.* 302. *Engl. Bot. t.* 347. *Jacq. Austr. t.* 394.

C. staphylini folio aliquatenus accedens. *Bauh. Hist. v. 3. p. 2.* 191. *Raii Syn.* 210.

C. aquatica, pimpinellæ saxifragæ divisurâ nostras. *Pluk. Phyt. t.* 49. *f.* 4.
Angl. Parsley Water Dropwort.

In palustribus salis.

Salt marshes near Aber, North Wales. *Rev. Hugh Davies.* Cambridgeshire. *Rev. J. Hemsted.*

Perennis. Fl. Julio.

Radix e tuberibus pluribus, fusiformibus, gracilibus, carnosis, fibris intermixtis. Caulis erectiusculus, angulato-cylindricus, striatus, glaber, foliosus, parùm ramosus. Folia radicalia bipinnata; foliolis cuneatis, lobatis, subinde ellipticis, indivisis: caulina pinnata; foliolis elongatis, lineari-lanceolatis, acutis, integerrimis. Umbellæ terminales, pedunculatæ, erectæ, multiradiatæ, planiusculæ. Umbellulæ densæ, multifloræ. Involucra et involucella e foliolis magis vel minùs numerosis, linearibus, acutis. Flores albi, extùs sæpe rubicundi, radiantes, maximâ ex parte abortientes. Dentes calycini ovati, acuti, inæquales, patentes. Variat involucri monophyllo. Folia radicalia quandoque profundius incisa inveniuntur, more Dauci (*Staphylini antiquorum*), ut in icone Plukenetianâ videri est.

3. *CENANTHE* foliolis omnibus linearibus, in-*peucedan-*
volucro nullo, tuberibus radicalibus ellipticis *nifolia.*
sessilibus.

Cenante *peucedanifolia.* *Pollich. Palat. v. 1.*
289. *t. 3. Sibth. 98. Engl. Bot. t. 348.*
Sym. Syn. 73. Abbot. 64. Willden. Sp. v. 1.
1442.

Narrow Dropwort. *Pet. H. Brit. t. 25. f. 8 ?*

Filipendula angustifolia. *Ger. em. 1059.*

Bolbocastanum fœmina. *Dalech. Hist. 773.*

Angl. Sulphur-wort Water Dropwort.

In fossis et paludosis.

Near Bury. *Mr. Mathew.* About Bedford. *Rev.*

Mr. Abbot. In Oxfordshire. *Sibth.*

Perennis. Fl. Junio.

Radix e tuberibus confertis, ellipticis, sessilibus. Caulis
præcedente firmior, crassior, erectus, angulato-teres,
striatus, foliosus, parùm ramosus. Folia radicalia bi-
pinnata; caulina pinnata; omnia foliolis linearibus,
acutis. Umbellæ radii firmiores, minùsque numerosæ
quam in *Æ.* pimpinelloide. Involucrum universale
nullum. Umbellulæ multifloræ, planæ, disco sæpiùs
rubræ, radio albæ. Involucella polyphylla. Calyx
erectiusculus, valdè irregularis.

4. *CENANTHE* foliolis omnibus cuneatis multi-*crocata.*
fidis subæqualibus.

Cenante *crocata.* *Sp. Pl. 365. Hudf. 121.*

With. 302. Sibth. 99. Jacq. Hort. Vind. v. 3.

t. 55. Woodv. Suppl. t. 267.

Æ. *cicutæ* facie *Lobelii.* *Raii Syn. 210.*

Æ. *maxima,* succo viroso, *cicutæ* facie. *Morif.*

sect. 9. t. 7. f. 2.

Filipendula cicutæ facie. *Ger. em. 1059.*

Hemlock Dropwort. *Pet. H. Brit. t. 25. f. 7.*

Angl. Hemlock Water Dropwort.

In aquosis, et ad ripas fluviorum.

Common about the Thames, and found here and there in all parts of Great Britain.

Perennis. Fl. Julio.

Radix ferè præcedentis. Caules erecti, ramosi, subdichotomi, foliosi, teretes, sulcati. Folia saturatè viridia, bipinnata; foliolis oppositis alternisque, sessilibus, cuneatis, vel subdeltoideis, incisiss, venosis, glabris. Umbellæ terminales, multiradiatæ, hemisphæricæ. Umbellulæ multifloræ. Involucra et involucella polyphylla, foliacea, elongata, dilatata, obtusiuscula. Flores albi, subradiantes. Calyx incurvus. Fructus oblongus, stylis erectis aristatus.

Herba summè venenata, imprimis radix. Succo croceo foetido scatet.

142. CORIANDRUM. *G. Pl.* 142. *Juss.* 220.
Gærtn. t. 22.

Fructus sphæricus, glaber. *Corolla* radiata. *Petalata* inflexo-emarginata. *Involucrum* monophyllum. *Involucella* dimidiata.

sativum. * 1. CORIANDRUM fructibus globosis, seminibus hemisphæricis.

Coriandrum sativum. *Sp. Pl.* 367. *Huds.* 123.
With. 305. *Engl. Bot. t.* 67. *Woodv. Med.*
Bot. t. 181. *Mart. Fl. Rust. t.* 141.

Coriandrum. *Raii Syn.* 221. *Ger. em.* 1012.
Fuchsf. Hist. 345. *Camer. Epit.* 523. *Brunf.*
Herb. v. 1. 203.

Coriandrum, semen. *Pharmac. Lond.*
Angl. Common Coriander.

In arvis et fimetis, haud indigena.

About Ipswich, and in some parts of Essex.

Annua. Fl. Junio.

Herba glabra, foetens. Caulis erectus, ramosus, divaricatus, teres, obsolete striatus. Folia composita; inferiora pinnata, foliolis cuneato-subrotundis, incisiss; superiora

periora triternata, laciniis linearibus. Umbellæ et umbellulæ multiradiatæ. Involucrum monophyllum, lineare. Involucella dimidiata, lanceolata, 3—5-phylla. Flores albi vel rubicundi; exteriores maximè irregulares. Calyx pentaphyllus. Fructus globosus, obsoletè costatus, aromaticus. Semina concava.

B. Involucris partialibus, universali nullo.

143. PHELLANDRIUM. *G. Pl.* 140.
Juss. 221.

Flosculi omnes fertiles; *disci* minores. *Fructus* ovatus, lævis, calyce styliisque coronatus.

1. PHELLANDRIUM foliorum ramificationibus *aquaticum*,
divaricatis.

Phellandrium aquaticum. *Sp. Pl.* 366. *Huds.*
122. *With.* 303. *Relb.* 119. *Sibth.* 99. *Woodv.*
Suppl. t. 266.

P. n. 757. *Hall. Hist. v.* 1. 332.

P. vel Cicutaria aquatica quorundam. *Raii Syn.*
215. *Bauh. Hist. v.* 3. *p.* 2. 183.

Cicutaria palustris. *Ger. em.* 1063. *Lob. Ic.*
p. 1. 735.

Angl. Water Hemlock.

In fluviis et fossis.

Biennis. Fl. Junio, Julio.

Radix fusiformis, crassa, fibris verticillatis. Caulis teres, cavus, flexuosus, ramosus, foliosus, divaricatus, sulcatus. Folia patentissima, tripinnata, incisa. Umbellæ oppositifoliæ, patentes, multiradiatæ. Umbellulæ densissimæ. Involucella polyphylla, setacea. Flores albi; exteriores majores, irregulares. Calyx pentaphyllus. Fructus ovatus, subcompressus, lævis, calyce styliisque porrectis coronatus. Genus *Oenanthi* affine.

144. CICUTA. *G. Pl.* 141.
Cicutaria. *Juss.* 221.

Fruſtus ſubovatus, ſulcatus. *Corolla* regularis.

vioſa. 1. CICUTA umbellis oppoſitifoliis, ſtipulis petiolo
adnatis obtuſis.

Cicuta viroſa. *Sp. Pl.* 366. *Hudſ.* 122. *With.*
304. *Relb.* 119. *Engl. Bot. t.* 479. *Fl.*
Dan. t. 208. *Woodw. Suppl. t.* 268.

C. aquatica Geſneri. *Bauh. Hiſt. v.* 3. *p.* 2.
175.

Sium alterum Olufatri facie. *Raii Syn.* 212.
Ger. em. 256. *Lob. Ic. p.* 1. 208.

Angl. Water Hemlock. Water Cowbane.

In ſoſſis et ad ripas fluviorum haud frequens.

In Cheſhire, Staffordſhire and Shropſhire. *With.*

In the river above Norwich, and on St. Faith's
bogs. *Mr. Pitchford.* Near London. *Hudſ.*

Iſle of Ely. *Relb.* Tuddenham heath, Suffol.

Rev. Mr. Hemſted. Very rare in Scotland. *Lightf.*

In Newſham Carr near Thirſk. *Sir T. Frank-*
land.

Perennis. Fl. Auguſto.

Radix tuberoſa, cava, concamerata, ſibris ſubverticillatis.

Caulis tripedalis, ramoſus, ſulcatus, folioſus, ramis erec-
tiuſculis. *Folia* biternata, majora pinnata; foliola lan-
ceolata, acuta, ſerrata, glabra. *Stipulæ* lineares, pe-
tiolis adnatæ, parùmque breviores. *Umbellæ* oppoſiti-
foliæ, erectæ, multiradiatæ. *Umbellulæ* denſæ. *Flores*
albi, regulares, parvi. *Calyx* quinqueſidus. *Petala* in-
voluta. *Fruſtus* compreſſus, rotundatus, ſerè didymus,
coſtatus.

145. *ÆTHUSA*. *G. Pl.* 141. *Juss.* 220.

Gærtn. t. 22.

Fructus striatus. *Involucella* dimidiata, triphylla, pendula.

r. *ÆTHUSA* foliis conformibus.

Cynapi.

Æthusa *Cynapium*. *Sp. Pl.* 367. *Huds.* 123. *um.*

Wub. 304. *Relb.* 120. *Sibth.* 99. *Curt.*

Lond. fasc. 1. *t.* 18.

Cicutaria tenuifolia. *Raii Syn.* 215. *Ger. em.* 1063.

C. apii folio. *Baub. Hist.* v. 3. *p.* 2. 180.

Angl. Fool's Parsley. Lesser Hemlock.

In hortis et arvis vulgaris.

Annua. *Fl.* Julio, Augusto.

Radix fusiformis. *Caulis* teres, subramosus, foliosus, parum striatus. *Folia* glabra, saturatè viridia, bipinnata, foliolis pinnatifidis, decurrentibus. *Umbellæ* longius pedunculatæ, oppositifoliæ et terminales, planiusculæ, radiis horizontaliter patentibus. *Involucella* linearilanceolata, pendula. *Flores* albi, æquales, parum radiant. *Fructus* ovatus, striatus.

146. *SCANDIX*. *G. Pl.* 142. *Juss.* 220.

Gærtn. t. 85.

Myrrhis. *Gærtn. t.* 23.

Flores radiantes. *Petala* emarginata. *Fructus* subulatus. *Flores* disci sæpè masculi.

ii. *SCANDIX* seminibus sulcatis angulatis.

odorata.

Scandix odorata. *Sp. Pl.* 368. *Huds.* 124.

Wub. 306. *Sibth.* 100. *Jacq. Austr. app.* 1. 37.

Cerfolium magnum five *Myrrhis*. *Ger. em.* 1039.

Myrrhis. *Lob. Ic.* 734. *Camer. Epit.* 398.

X 2

Angl.

Angl. Sweet Cicely. Great Chervil.

In pratis montosis.

In meadows betwixt Morton and Rushforth, in the west riding of Yorkshire, certainly a native. *Dr. Richardson. Herb. Buddl.* Near Leeds pretty common, and certainly wild. *Rev. Mr. Wood.* Behind a farm-house about a mile west of Hafod, Cardiganshire. It is frequent, as Hudson remarks, in Westmoreland, Cumberland and Lancashire, but generally near houses.

Perennis? Fl. Maio.

Radix fusiformis, aromatica. Caules sesquipedales, subæquales, striati, foliosi. Folia sessilia, subtripinnata, foliolis pinnatifidis, acutis, ferratis, pilosis, pallidè virentibus. Umbellæ terminales, erectæ, multiradiatæ. Involucella lanceolata, acuminata, membranacea, albida, deflexa. Flores parvi, nivei; disci numerosiores, masculi, pistillis abortientibus. Semina maxima, erecta, ferè uncialia, lanceolata, acuta, non rostrata, sulcata, nitida, fusca, dulcia et aromatica, stylis divaricatis coronata.

Pecten-Veneris. 2. SCANDIX feminibus longissimè rostratis scabriusculis, foliolis lineari-multipartitis.

Scandix Pecten-Veneris. *Sp. Pl.* 368. *Huds.* 123. *With.* 307. *Relb.* 120. *Sibth.* 100. *Curt. Londl. fasc.* 5. *t.* 21. *Mart. Fl. Rust. t.* 38. *Fl. Dan. t.* 844. *Jacq. Austr. t.* 263. *Dicks. H. Sicc. fasc.* 15. 4.

S. femine rostrato vulgaris. *Raii Syn.* 207.

Pecten Veneris. *Camer. Epit.* 304.

P. Veneris five Scandix. *Ger. em.* 1040.

Angl. Needle Chervil. Venus's comb. Shepherd's needle.

In arvis vulgaris.

Annua. Fl. Junio, Julio,

Radix parva. Caules patuli. Folia tripinnata, foliolis multipartitis, laciniis alternis, linearibus, læviusculis. Umbellæ radii 2 vel 3. Umbellulæ multifloræ. Involucella apice bifida, trifida, vel pinnatifida, margine scabra. Flores albi. Semina cylindrica, scabriuscula, rostro communi longissimo, ancipiti, angulis scabro.

3. SCANDIX feminibus ovatis hispidis, corollis *Anthriscus.* uniformibus, caule lævi.

Scandix Anthriscus. *Sp. Pl.* 368. *Huds.* 124.

Relb. 121. *Sibth.* 100. *Curt. Lond. fasc.* 1.

t. 19. *Mart. Fl. Rust. t.* 75. *Jacq. Austr.*

t. 154. *Dickf. H. Sicc. fasc.* 15. 3.

Caucalis scandicina. *Wigg. Holsat.* 23. *With.* 289. *Fl. Dan. t.* 863.

Myrrhis sylvestris feminibus asperis. *Raii Syn.* 220.

M. æquicolorum nova. *Column. Ecphr.* 110. *t.* 112.

Angl. Rough Chervil.

In aggeribus et ruderatis suburbanis.

Annua. Fl. Maio.

Caulis 2—3-pedalis, teres, gibbosus, striatus, glaber, nitidus. Folia petiolata, lætè viridia, hirsuta, tripinnata, foliolis ovatis, lobatis. Umbellæ laterales et terminales subquinqueradiatæ. Umbellulæ quinque-floræ. Involucrum nullum. Involucella parva, ovata. Flores exigui, regulares, albi. Semina lanceolata, aculeis fursum curvatis aspera, rostro brevissimo, lævi.

Scandicis species proculdubio est, ut ex charactere, habitu, et totâ structurâ luce clariùs patet.

147. CHÆROPHYLLUM. *G. Pl.* 143. *Juss.* 220.

Myrrhis. *Gartn. t.* 23.

Petala cordata. *Fructus* oblongus, læviusculus.

Involucella reflexa, concava.

Sylvestre, 1. CHÆROPHYLLUM caule striato lævi, geniculis tumidiusculis.

Chærophyllum sylvestre. *Sp. Pl.* 369. *Huds.* 124. *Wilh.* 308. *Relb.* 121. *Sibth.* 100. *Curt. Lond. fasc.* 4. *t.* 25. *Mart. Fl. Rust.* *t.* 96. *Jacq. Austr.* *t.* 149.

Cicutaria vulgaris. *Raii Syn.* 207.

Myrrhis. *Puchf. Hist.* 525. *Dalech. Hist.* 761.

Angl. Smooth Cow parsley. Wild Chervil.

Ad sepes vulgaris, imprimis solo fertilior.

Perennis. Fl. Aprili, Maio.

Radix fusiformis, subramosa, lactescens. Caulis subtripedalis, erectus, ramosus, foliosus, teres, striatus, lævis, infernè pubescens, sub foliorum insertione tumidiusculus. Folia tripinnata, foliolis acutè incis, margine scabris; petiolis brevibus, dilatatis, costatis. Umbellæ erectæ, terminales. Involucella ovata, membranacea, albida, ciliata, pilis longis crebris. Flores albi, parùm radiantes. Petala planiuscula, emarginata vel integra. Fructus elliptico-oblongus, teretiusculus, læviusculus vel obsoletè striatus, glaberrimus.

temulentum, 2. CHÆROPHYLLUM caule scabro, geniculis tumidis.

Chærophyllum temulentum. *Sp. Pl.* 370. *Huds.* 125. *Wilh.* 308. *Relb.* 122. *Sibth.* 101. *Curt. Lond. fasc.* 6. *t.* 24. *Fl. Dan.* *t.* 918. *Jacq. Austr.* *t.* 65.

Cereseolum sylvestre. *Raii Syn.* 207. *Ger. em.* 1038.

Angl. Rough Cow-parsley. Rough Chervil,

Ad sepes, et in umbrosis, vulgaris,

Biennis. Fl. Junio, Julio.

Radix subramosa. Caulis ferè tripedalis, erectus, ramosus, foliosus, teres, obsoletè striatus, scaber, purpureo-maculatus, sub foliorum insertione tumidus. Folia bipinnata,

pinnata, pinnatifida, hirsuta, subtùs pallidiora. Umbellæ ferè præcedentis, sed ante florescentiam cernuæ. Radii scabri. Involucella basi subconnata, lanceolata, margine et carinâ scabra. Flores nivei. Petala inæqualia, profundè biloba. Fructus vix striatus, glaberimus.

148. IMPERATORIA. *G. Pl.* 143. *Juss.* 220. —
Gærtn. t. 21.

Petala inflexo-emarginata, subæqualia. *Fructus* subrotundus, compressus, marginatus, medio gibbus, tricostratus. *Umbellæ* planæ.

1. IMPERATORIA.

Ostruthi-

Imperatoria Ostruthium. Sp. Pl. 371. *Lightf.* *um.*
168. *Huds.* 649. *With.* 308. *Woodv. Med.*
Bot. t. 35.

Imperatoria. Ger. em. 1001. *Camer. Epit.* 552.

I. major. Garid. Prov. t. 55.

LasERPitium germanicum. Fuchsf. Hist. 763.

Smyrnion. Trag. Hist. 433.

Angl. Masterwort.

In pratis humidiusculis subalpinis.

On the banks of the Clyde in several places, particularly about Ardencaple, and in the Isle of Bute, near Mount Stuart; but whether indigenous or not is uncertain. *Lightf.*

Perennis. Fl. Junio.

Radix tuberosa, acris et aromatica. *Caules* erecti, sesquipedales, subsimplices, teretes, striati, glabri. *Folia* biternata, glabra, acutè serrata et incisa. *Umbellæ* terminales, paucæ, planæ, multiradiatæ, læves; umbellulæ multifloræ. *Involucella* e foliolis paucis, setaceis. *Flores* albi vel rubicundi, regulares. *Petala* elliptica, mucrone inflexo. *Fructus* dilatatus, apice basi que emarginatus, glaber.

C. Involucro nullo; vix universali, nunquam partialibus.

149. PASTINACA. *G. Pl.* 144. *Juss.* 219.
Gærtn. t. 21.

Fructus ellipticus, compresso-planus. *Petala* involuta, integra.

Fativa. 1. PASTINACA foliis simpliciter pinnatis subtus pubescentibus.

Pastinaca fativa. *Sp. Pl.* 376. *With.* 309.
Mart. Fl. Rust. t. 83. *Engl. Bot. t.* 556.

P. sylvestris. *Huds.* 125. *Relb.* 122. *Sibth.* 101.
Abbot. 67.

P. sylvestris latifolia. *Raii Syn.* 206. *Ger. em.* 1025.

Sifer sylvestre. *Fuchs. Hist.* 753.

Angl. Wild Parsnep.

In marginibus agrorum, collibusque, solo calcareo.

Biennis. *Fl.* Julio.

Radix fusiformis, aromatica, dulcis, sed acris. *Caulis* tripedalis, erectus, ramosus, angulatus, sulcatus, scabriusculus. *Folia* pinnata; foliolis 5—9, inciso-ferratis, subtus pubescentibus; impari trilobo. *Umbellæ* terminales, solitariae, multiradiatae, erectae, scabriusculæ. *Involucra* et *involucella* plerumque nulla. *Corolla* parva, flava. *Fructus* magnus.

150. SMYRNIUM. *G. Pl.* 144. *Juss.* 219.
Gærtn. t. 22.

Fructus oblongus, angulato-costatus. *Petala* acuminata, carinata, incurva.

Olusatrum. 1. SMYRNIUM foliis caulinis ternatis petiolatis ferratis.

Smyrnum Olusatrum. *Sp. Pl.* 376. *Huds.* 126.
With.

With. 310. *Relb.* 123. *Sibth.* 101. *Engl.*

Bot. t. 230. *Dicksf. H. Sicc. fasc.* 9. 2.

Smyrnum. *Raii Syn.* 208. *Matth. Valgr. v.* 2.
130.

Hippofelinum. *Ger. em.* 1019,

Angl. Alexanders.

Ad muros antiquos, interdum et in rupibus maritimis.

Common in Anglesea. *Ray. Pennant.* Very common in all the western counties. *With.* At Scarborough castle, Nottingham, York, Norwich and Bungay. *See Ray, With. &c.*

Biennis. Fl. Maio.

Radix carnosa, ramosa. Herba lucida, pallidè virens, glabra, aromatica, subdulcis. Caulis teres, fulcatus. Folia superiora ternata, inferiora triternata, foliolis latis, figurâ variis, inciso-ferratis, sub-petiolatis. Petiolus communis basi dilatatus, ventricosus, nervosus. Umbellæ terminales, globosæ, multiradiatæ. Flores virescentes. Fructus didymus, niger, utrinque tricostratus.

151. ANETHUM. *G. Pl.* 145. *Juss.* 219.

Fœniculum. *Gærtn. t.* 23.

Fructus ovatus, subcompressus, striatus. *Petala* involuta, integra.

1. ANETHUM fructibus gibbis, foliis caulinis *Fœniculum.*
numerosis deflexis.

Anethum Fœniculum. *Sp. Pl.* 377. *Huds.* 126.

With. 310. *Relb.* 123. *Mill. Illust. t.* 13.

Woodv. Med. Bot. t. 160.

Fœniculum. *Camer. Epit.* 534. *Fuchsf. Hist.*
501.

F. vulgare. *Raii Syn.* 217. *Ger. em.* 1032.

Gærtn. v. 1. 105.

Fœniculum dulce, semen. *Pharmac. Lond.*

Angl. Common Fennel.

In clivis cretaceis copiosè.

Biennis. Fl. Julio, Augusto.

Radix fusiformis. Caulis quadripedalis, erectus, ramossimus, foliosus, teres, striatus, glaber. Folia tripinnata, foliolis setaceis, longissimis, glabris, atro-viridibus, dependentibus. Umbellæ terminales, magnæ, multiradiatæ, planiusculæ. Flores aurei. Fructus ovatus, vix compressus, utrinque tricostratus, margine membranaceo orbatus.

152. CARUM. *G. Pl.* 145. *Juss.* 219.

Gærtn. t. 23.

Fructus elliptico-teres, striatus. *Petala* carinata, inflexo-emarginata. *Involucrum universale* sub-monophyllum.

Carui. * 1. CARUM.

Carum Carui. *Sp. Pl.* 378. *Huds.* 126. *With.*

311. *Relb.* 124. *Mart. Fl. Rust. t.* 55.

Woodv. Med. Bot. t. 45. *Jacq. Austr. t.* 393.

C. seu Careum. *Raii Syn.* 213. *Ger. em.* 1034.

Camer. Epit. 516.

Caros. *Fuchs. Hist.* 396.

Caruon, semen. *Pharmac. Lond.*

Angl. Common Caraway.

In pratis et pascuis.

Biennis. Fl. Junio.

Radix fusiformis. Caulis ferè bipedalis, ramosus, angulatus, sulcatus, glaber. Folia glabra, bipinnata, incisa, foliolis infimis decussantibus. Umbellæ numerosæ, terminales, erectæ. Involucrum e foliolis setaceis, folitariis, binis vel ternis, sæpe nullum. Involucella nulla. Flores albi. Petala inflexo-cordata, subæqualia. Fructus parvus, oblongus, gratè aromaticus.

153. PIMPINELLA. *G. Pl.* 145. *Juss.* 219.
Anisum. *Gærtn. t.* 23.

Fructus ovato-oblongus, striatus. *Petala* inflexa.
Stigmata subglobosa.

1. PIMPINELLA foliis pinnatis: foliolis radi-*saxifraga*
calibus subrotundis; summis linearibus. *ga.*

Pimpinella saxifraga. *Sp. Pl.* 378. *Huds.* 127.
With. 311. *Relb.* 124. *Sibth.* 102. *Engl.*
Bot. t. 407. *Mart. Fl. Rust. t.* 127. *Fl. Dan.*
t. 669. *Jacq. Austr. t.* 395.

P. saxifraga minor, foliis sanguisorbæ. *Raii*
Syn. 213.

P. minor. *Camer. Epit.* 776.

Bipinella, five *Saxifraga minor.* *Ger. em.* 1044.

- β. *Pimpinella dissecta.* *Sibth.* 102. *Abbot.* 68.
Retz. Obs. fasc. 3. 30: *t.* 2.

P. saxifraga minor. *Raii Syn.* 213?

Small Cut Saxifrage. *Pet. H. Brit. t.* 26. *f.* 8.

Angl. Common Burnet-Saxifrage.

In pascuis ficcioribus, glareosis vel calcareis.

Perennis. *Fl.* Julio, Augusto.

Radix lignosa, aromatica, pungens. Herba scabriuscula, subpubescens. Caules erecti, rigidi, subflexuosi, teretes, striati, ramis floriferis elongatis, nudiusculis. Folia pinnata, variè incisa; inferiora foliolis plerumque ovatis; superiora pinnato-multifidis; summa linearia, et quasi abortiva. Umbellæ terminales. Flores albi, ferè regulares. Petala lanceolata, incurva, canaliculata. Stigmata capitata, quod ex errore in *Engl. Bot.* non exhibetur. Semina parva, receptaculo floris carnosio coronata.

2. PIMPINELLA foliis pinnatis: foliolis omni-*magna*
bus ovatis; impari trilobo.

Pimpinella magna. *Mant.* 2. 219. *With.* 313.
Relb.

Relb. 125. *Sibth.* 102. *Engl. Bot.* t. 408.
Abbot. 68.

P. major. *Hudf.* 127. *Fuchsf. Hist.* 608. *Jacq.*
Austr. t. 396.

P. saxifraga. *Raii Syn.* 213. *Ger. em.* 1044.
Matth. Valgr. v. 2. 379. *Camer. Epit.* 775.

β. *P. saxifraga major*, degener, seu foliis dissectis.
Dill. in Raii Syn. 213.

Great Cut Saxifrage. *Pet. H. Brit.* t. 26. f. 6.
Angl. Great Burnet-Saxifrage.

In umbrosis et ad sepes solo calcareo.

Perennis. Fl. Julio, Augusto.

Radix præcedente major, viribus debilior. Caules bipedales, vel tripedales. Folia pinnata, omnia ferè conformia, foliolis ovatis, inciso-serratis, terminali trilobo, Umbellæ magnæ. Flores et fructus ferè præcedentis, Variat etiam foliis dissectis.

dioica,

3. PIMPINELLA foliolis omnibus linearibus, umbellis paniculatis, floribus dioicis.

Pimpinella dioica. *Syst. Veg. ed.* 13. 241.
Hudf. 128. *With.* 313.

P. pumila. *Jacq. En. Vind.* 52. 227. *Austr.* t. 28.
Seseli pumilum. *Sp. Pl.* 373.

Peucedanum minus. *Raii Syn.* 217. *Hudf.*
ed. 1. 101.

P. pumilum. *Ger. em.* 1054.

Selinum montanum pumilum. *Clus. Hist.* v. 2.
200. *Bauh. Hist.* v. 3. p. 2. 17.

Angl. Dwarf Burnet-Saxifrage. Rock Parsley.

In rupibus calcareis, rariùs.

On St. Vincent's rocks, Bristol. *Ray.* Above Uphill, Somersetshire. *Hudf.*

Perennis. Fl. Maio, Junio.

Radix carnosa, ramosa. Herba dioica, glauca, lævis.
Caulis

Caulis spithamæus, ramosus, angulatus, sulcatus, purpurescens. Folia bipinnata, foliolis linearibus, uniformibus, canaliculatis. Umbellæ numerosæ, paniculatæ, simplices et compositæ, erectæ. Flores albidî; masculinî germine plerumque destituti; fœminêi antheris abortivis muniti. Stigmata capitato-globosa, Fructus ovatus, sulcatus.

154. APIUM. *G. Pl.* 146. *Juss.* 219.

Gærtn. t. 22.

Fructus ovatus, costatus. *Petala* inflexa, æqualia. *Involucrum* monophyllum.

1. APIUM foliolis caulinis cuneiformibus, caule *graveolens.*
sulcato.

Apium graveolens. *Sp. Pl.* 379. *Huds.* 129.

With. 314. *Relb.* 125. *Sibth.* 103. *Fl. Dan.*
t. 790.

A. palustre et officinarum. *Raii Syn.* 214.

Apium. *Fuchs. Hist.* 744.

A. palustre. *Camer. Epit.* 527.

Eleoselinum five *Paludapium.* *Ger. em.* 1014.

Angl. Smallage. *Wild Celery.*

In fossis et paludibus, præsertim maritimis.

Biennis. *Fl.* Augusto.

Radix fusiformis. *Caules* patuli, elongati, sulcati, foliosi, glabri. *Folia* lucida, ternata, foliolis cuneatis, incisis.

Umbellæ terminales et laterales, subinde sessiles, inæquales. *Involucrum* sæpe nullum. *Flores* albidî, parvi.

Fructus ovato-suborbiculatus.

Herba acris, fœtida, venenata, culturâ subdulcis et esculenta evadit.

155. ÆGOPodium. *G. Pl.* 146. *Juss.* 219.

Gærtn. t. 140.

Fructus ovato-oblongus, costatus. *Petala* inflexo-cordata, inæqualia.

1. ÆGO-

Podagra- 1. **ÆGOPODIUM.**
ria.

Ægopodium Podagraria. *Sp. Pl.* 379. *Huds.* 129.
With. 314. *Relb.* 126. *Sibth.* 103. *Fl. Dan.*
t. 670.

Angelica sylvestris minor seu erratica. *Raii*
Syn. 208.

Herba Gerardi. *Ger. em.* 1001.

Angl. Gout-weed. *Pet. H. Brit. t.* 25. *f.* 10.

In cultis, umbrosis ruderalis, et ad sepes.

Perennis. *Fl.* Maio, Junio.

Radix repens, vix extirpanda. Caules erecti, sulcati, foliosi. Folia radicalia biternata; caulina ternata; foliolis petiolatis, ovatis, acutis, duplicato-ferratis, glabris. Umbellæ terminales, solitariae, multiradiatae. Flores nivei. Petala parum inæqualia. Fructus ferè ellipticus, utrinque tricostratus.

III. TRIGYNIA.

156. **VIBURNUM.** *G. Pl.* 147. *Juss.* 213.
Gærtn. t. 27.

Cal. superus, quinquepartitus. Cor. quinque-
fida. Bacca monosperma.

Lanta- 1. **VIBURNUM** foliis cordatis ferratis venosis
na. subtùs tomentosis.

Viburnum Lantana. *Sp. Pl.* 384. *Huds.* 129.
With. 315. *Relb.* 126. *Sibth.* 104. *Engl.*
Bot. t. 331. *Jacq. Austr. t.* 341.

Viburnum. *Raii Syn.* 460. *Camer. Epit.* 122.
8 *Lantana,*

Lantana, five Viburnum. *Ger. em.* 1490.

Angl. Mealy Guelder-rose. Way-faring Tree.

In nemorosis et sepibus, solo calcareo.

Arbuscula. Fl. Maio.

Rami teretes, viminei; juniores tomentosi, foliosi. Folia opposita, petiolata, cordata, obtusa, ferrata, venosa, subtus præcipuè villis stellatis pubescentia, demum rubicunda, decidua. Stipulæ nullæ. Cymæ terminales, solitariae, tomentosae, multifloræ. Calyx minutus. Corolla alba, semiquinquesida, patens. Stamina inter lacinias inserta. Stigmata fœcilia, brevissima, obtusa. Baccæ compressæ, farinosæ, nigrae. Semen cordatum, compressum, sulcatum.

2. VIBURNUM foliis lobatis, petiolis glandu- *Opulus.*
lofis.

Viburnum Opulus. *Sp. Pl.* 384. *Huds.* 130.

With. 315. *Relb.* 127. *Sibth.* 104. *Fl. Dan.*

t. 661. *Engl. Bot. t.* 332.

Opulus. *Raii Syn.* 460.

Sambucus aquatilis five palustris. *Ger. em.* 1424.

S. aquatica. *Camer. Epit.* 977.

Angl. Common Guelder-rose. Water Elder.

In umbrosis et sepibus humidis.

Arbuscula. Fl. Junio.

Arbuscula glabra, ramosissima, ramis oppositis, teretibus. Folia subcordata, triloba, inæqualiter ferrata, venosa, subtus pallidiora, sero autumnno saturatè rosea, decidua. Petioli basin versus foliolosi, apice glanduliferi. Cymæ terminales, solitarii, multiflori, nivei, radiantes, floribus exterioribus abortivis, dilatatis, irregularibus. Baccæ ellipticæ, rubrae, succosæ. Semen compressum, subcordatum.

157. SAMBUCUS. *G. Pl.* 147. *Juss.* 214.

Gærtn. t. 27.

Cal. superus, quinquepartitus. *Cor.* quinquesida.

Bacca trisperma.

Ebulus. 1. **SAMBUCUS** cymis tripartitis, stipulis foliaceis, caule herbaceo.

Sambucus Ebulus. *Sp. Pl.* 385. *Huds.* 130.

With. 316. *Relb.* 127. *Sibth.* 104. *Abbot.* 70.

Curt. Lond. fasc. 3. *t.* 18. *Engl. Bot. t.* 475.

Woodv. Suppl. t. 260.

S. humilis, seu *Ebulus.* *Raii Syn.* 461.

Ebulus. *Camer. Epit.* 979. *Fuchs. Hist.* 65.

Ic. 38.

E. five Sambucus humilis. *Ger. em.* 1426.

Angl. Dwarf Elder. *Danewort.*

In ruderatis et ad sepes, minùs vulgaris.

Perennis. *Fl. Julio.*

Radix repens. Caules herbacei, simplices, erecti, tripedales, fragiles, teretiusculi, fulcati, scabri, foliosi. Folia opposita, impari-pinnata, atro-viridia, læviuscula, venosa, serrata; foliolis basi inæqualibus et sæpiùs glandulosis. Stipulæ foliaceæ, serratæ, magnæ. Cyma terminalis, trifida, ramis ramulosis, pilosis, multifloris. Flores omnes pedicellati, tristè incarnati. Stamina crassa, erecta, alba. Antheræ bilobæ, rubræ. Baccæ globosæ, nigræ. Planta fœtida, cathartica, teterrima.

nigra. 2. **SAMBUCUS** cymis quinquepartitis, foliolis ovatis ferratis, caule arboreo.

Sambucus nigra. *Sp. Pl.* 385. *Huds.* 130.

With. 316. *Relb.* 128. *Sibth.* 105. *Woodv.*

Med. Bot. t. 78. *Fl. Dan. t.* 545. *Engl. Bot.*

t. 476.

Sambucus. *Raii Syn.* 461. *Ger. em.* 1422.

Camer. Epit. 975.

β. *S. fructu albo.* *Ger. em.* 1422.

S. acinis albis. *Raii Syn.* 461.

γ. *S. laciniatis foliis.* *Ger. em.* 1422.

S. laciniato folio. *Raii Syn.* 461.

Sambucus, cortex interior, flos, bacca. *Pharmac.*

Lond.

Angl.

Angl. Common Elder. γ Parsley-leaved Elder.

In sepibus et nemorosis vulgatissima.

Arbuseula. Fl. Junio.

Arbor humilis, ramosissima. Rami oppositi, teretes, cinerei, glabri. Folia opposita, impari-pinnata, foliolis plerumque quinis, ovatis, ferratis, glabris. Stipulæ vix ullæ. Cymæ terminales, quinquefidæ, multifloræ. Flores lactei, suaveolentes, plerumque pedicellati. Stamina divaricata. Baccæ globosæ, atro-purpureæ, subdulces sed nauseosæ; in β et γ albidæ, minusque ingratae.

158. STAPHYLEA. *G. Pl.* 148. *Juss.* 377.

Gærtn. t. 69.

Cal. inferus, quinquepartitus. *Pet.* quinque.

Capsulæ inflatæ, connatæ. *Semina* bina, globosa, cum cicatrice.

1. STAPHYLEA foliis pinnatis, stylis capsulisque *pinnatis*.
binis.

Staphylea pinnata. *Sp. Pl.* 386. *Huds.* 131.

With. 317.

Staphylodendron. *Raii Syn.* 468. *Baub. Hist.*

v. 1. 274. *Camer. Epit.* 171.

Nux vesicaria. *Ger. em.* 1437.

Angl. Bladder-nut tree.

In sepibus, vix indigena.

In hedges about Pontefract, Yorkshire, but scarcely in sufficient plenty to be deemed certainly wild.

Ray. About Ashford, Kent. *Parkinson.*

Frutex. Fl. Junio.

Arbuseula humilis, ramosa, Fraxini facie, glabra. Folia alterna, impari-pinnata, foliolis quinis, ovatis, ferrulatis, acutis. Racemi terminales, compositi, nutantes. Bractææ setaceæ, membranaceæ. Flores campanulati, luteoli. Capsulæ binæ, rarius ternæ, membranaceæ,

ventricosæ, plerumque dispermæ. Semina globosa, fusca, nitida, dura.

159. TAMARIX. *G. Pl.* 148. *Juss.* 313.

Gærtn. t. 61.

Cal. inferus, quinquepartitus. *Pet.* quinque.
Capsula unilocularis, trivalvis. *Semina* pap-
posa.

gallica. 1. TAMARIX floribus pentandris.

Tamarix gallica. *Sp. Pl.* 386. *With.* 318. *Sym.*

Syn. 77. *Tr. of Linn. Soc. v.* 3. 333. *Mill. Ic.*
t. 262. *f.* 1.

Tamariscus Narbonensis. *Ger. em.* 1378. *Lob.*
Ic. p. 2. 218.

Myrica. *Camer. Epit.* 74. *f.* 1.

Angl. French Tamarisk.

In fissuris rupium, aggeribusque, maritimis Angliæ
australis.

Plentiful on St. Michael's mount, Cornwall, and
every where about the Lizard point, though
chiefly on the banks of earth called hedges, first
communicated by Mr. Giddy. *With.* Also
near Hurst castle, Hants. *Mr. W. G. Maton.*
On the cliff to the east of Haslings. *Rev. Dr.*
Goodenough.

Frutex. *Fl.* Julio.

Rami viminei, nutantes, teretes, rubri, nitidi, ramulis
secundis foliosis. Folia parva, imbricata, acuta, sub-
carnosa, lævia. Flores racemosi, bracteolati, incar-
nati, elegantes. Ovibus pabulum gratissimum præbet,
confer *Tour on the Continent, v.* 2. 321.

160. CORRIGIOLA. *G. Pl.* 149. *Juss.* 313.

Gærtn. t. 75.

Cal. inferus, pentaphyllus. *Pet.* quinque. *Semen*
unicum, rotundato-triquetrum.

1. COR-

I. CORRIGIOLA.

*littoralis!**Corrigiola littoralis. Sp. Pl. 388. With. 318.**Fl. Dan. t. 334. Dickf. Dr. Pl. 61. H. Sicc. fasc. 14. 10.**Polygoni, vel Linifolia per terram sparsa, flore Scorpoides. Bauh. Hist. v. 3. p. 2. 379.**Polygonum polyspermon litoreum minus, flosculis ex spadiceo albicantibus. Moris. sect. 5. t. 29. f. 1.**Angl. Sand Strapwort.**In littoribus maritimis Angliæ australis.**Found by Mr. Hudson on Slapham sands beyond Dartmouth, and near the Star point. Mr. Martin. On the beach near the tin mine at Helston, Cornwall. F. Borone.**Annua. Fl. Julio, Augusto.**Radix gracilis. Caules numerosi, prostrati, subsimplices, teretes, glabri, foliosi. Folia alterna, lineari-lanceolata, obtusa, integerrima, subcarnosa, glabra, glauca, basi attenuata. Stipulæ membranaceæ, lunatæ. Thyrsi terminales et laterales, multiflori, bracteolati. Flores albi, exigui. Calyx corollaque persistentes.*

IV. TETRAGYNIA.

161. PARNASSIA. *G. Pl. 151. Juss. 245.**Gærtn. t. 60.**Cal. quinquepartitus. Pet. quinque. Nectaria quinque, cordata, ciliata apicibus globosis.**Capsula quadrivalvis.*

palustris. 1. PARNASSIA.

Parnassia palustris. *Sp. Pl.* 391. *Huds.* 131.
With. 319. *Relb.* 129. *Sibth.* 105. *Ab-*
bot. 71. *Engl. Bot. t.* 82. *Dicks. H. Sicc.*
fasc. 7. 10. *Fl. Dan. t.* 584. *Mill. Illustr.*
t. 15.

P. vulgaris et palustris. *Raii Syn.* 355.

Gramen Parnassi. *Ger. em.* 840.

Angl. Grass of Parnassus.

In palustribus spongiosis copiosè.

Perennis. *Fl.* Septembri, Octobri.

Radix fibrosa. *Caulis* erectus, simplex, angulatus, tortus, glaber, unifolius et uniflorus. *Folia* cordata, integerima, nervosa, glabra; radicalia plurima, longius petiolata. *Flos* terminalis, formosus, niveus. *Petala* calyce longiora, cordata, concava, venis pellucidis picta. *Nectaria* petalis opposita atque breviora, carnosa, setis apice globiferis radiato-ciliata. *Stamina* inter nectaria, calyci opposita, pistillo ordinatim incumbentia. *Antheræ* cordatæ, ochroleucæ. *Germen* ovatum, stigmatibus sessilibus. *Semina* numerosa, minuta.

V. PENTAGYNIA.

162. STATICE. *G. Pl.* 153. *Juss.* 92.
Gærtn. t. 44.

Cal. monophyllus, integer, plicatus, scariosus.

Pet. quinque. *Semen* unicum, superum.

Armeria, 1. STATICE scapo simplicei capitato, foliis linearibus.

Statice

Statice Armeria. *Sp. Pl.* 394. *Huds.* 132.
With. 319. *Relb.* 129. *Lightf.* 173. *Engl.*
Bot. t. 226. *Dicks. H. Sicc. fasc.* 2. 9.

S. montana minor. *Raii Syn.* 203.

Caryophyllus marinus minimus Lobelii. *Ger.*
em. 602.

Angl. Thrift. Sea Gilliflower.

In maritimis limosis, etiam in pratis irriguis alpinis,
 rupiumque fissuris.

Perennis. Fl. Julio, Augusto.

Radix lignosa. Folia radicalia, cæspitosa, linearia, canaliculata, glabra. Scapi erecti, simplices, pubescentes, primo ortu membranâ cylindricâ, involucri adnatâ, tecti. Flores capitati, rosei. Involucrum polyphyllum, imbricatum, scariosum. Calyx parvus, erectus. Petala obovata, unguiculata.

2. STATICE scapo paniculato tereti, foliis lævibus enervibus, sub apice aristatis. *Limonium.*

Statice Limonium. *Sp. Pl.* 394. *Huds.* 132.
With. 320. *Engl. Bot. t.* 102. *Fl. Dan.*
t. 315.

Limonium. *Raii Syn.* 201. *Camer. Epit.* 721.
Ger. em. 411.

β. *L. minus.* *Raii Syn.* 202.

L. parvum. *Ger. em.* 411.

γ. *L. Anglicum minus,* caulibus ramosioribus, floribus in spicis rariùs fitis. *Dill. in Raii Syn.* 202.

Angl. Sea Lavender.

In maritimis cœnosis vulgaris.

Perennis. Fl. Julio, Augusto.

Radix lignosa. Herba magnitudine maximè varians, lævis. Folia obovato-lanceolata, sub apice mucronulata, integerrima, avenia. Scapus alternatim ramosus, ramulis corymbosis. Flores cyanei, speciosi, bracteati.

reticulata. 3. STATICE scapo paniculato prostrato flexuoso, ramis inferioribus sterilibus, foliis cuneiformibus muticis.

Statice reticulata. *Sp. Pl.* 394. *Huds.* 133. *With.* 320. *Engl. Bot. t.* 328. *Hill. Fl. Brit. t.* 25. *f.* 2.

Angl. Matted Sea Lavender.

In littoribus maritimis cœnosis Angliæ orientilibus.

About Wells, Cley and Brancafter. *Mr. Crowe.*
At Holme by the sea, Norfolk. *Rev. Mr. Sutton.*
Below Wisbeach. *Rev. Mr. Hemsted.*

Perennis. Fl. Julio, Augusto.

Radix lignosa. Folia spatulata, acuta, nec aristata. Scapi prostrati, ramosissimi; ramulis divaricatis, teretibus; inferioribus sterilibus. Flores terminales, fasciculati, cærulei, bracteis obtusis suffulti.

163. LINUM, *G. Pl.* 153. *Juss.* 303.
Gartn. t. 112.

Cal. pentaphyllus. *Pet.* quinque. *Capsula* supera, decemvalvis, decemlocularis. *Semina* solitaria.

* *Foliis alternis.*

usitatissimum. 1. LINUM foliolis calycinis ovatis acutis trinerviis, petalis crenatis, foliis lanceolatis alternis, caule subsolitario.

Linum usitatissimum. *Sp. Pl.* 397. *Huds.* 133. *With.* 321. *Relb.* 130. *Curt. Lond. fasc.* 5. *t.* 22. *Woodv. Med. Bot. t.* 111. *Mart. Fl. Rust. t.* 133.

L. sylvestre, fativum planè referens. *Raii Syn.* 362.

Linum, semen. *Pharmac. Lond.*
Angl. Common Flax.

Inter

Inter segetes frequens.

Annua. Fl. Julio.

Radix parva, fibrosa, annua. Caulis erectus, subsolitaris, basi simplex, supernè paniculatus, teres, glaber, foliosus. Folia alterna, lanceolata, acuminata, integerrima, trinervia, glabra. Flores pedunculati, erecti, cyanei. Folia calycina ovata, acuminata, trinervia, glabra, margine membranacea et subciliata. Petala obovata, obsolete crenata vel erosa, lineata. Stamina basi connata. Stigmata clavata, obtusa. Capsula acuminata. Semina elliptica, nitidissima.

2. LINUM foliolis calycinis obovatis obtusis sub *perenne*, quinquenervibus glabris, caulibus numerosis adscendentibus.

Linum perenne. *Sp. Pl.* 397. *Huds.* 133. *Wib.* 321. *Relb.* 130. *Engl. Bot.* t. 40. *Mart. Fl. Rust.* t. 134.

L. sylvestre cæruleum perenne erectius, flore et capitulo majore. *Raii Syn.* 362.

β. L. sylvestre cæruleum perenne procumbens, flore et capitulo minore. *Raii Syn.* 362.
Angl. Perennial Blue Flax.

In montosis cretaceis.

In Cambridgeshire, Northamptonshire and Westmoreland. At Marham, Norfolk.

Perennis. Fl. Junio, Julio.

Radix lignosa, multiceps. Caules plurimi, patentes, adscendentes (in β procumbentes), teretes, glabri, foliosi, apice parùm ramosi, ramis racemosis. Folia lineari-lanceolata, acuminata, subrecurva, uninervia, glauca, glabra, integerrima. Flores erecti, cyanei, præcedente parùm minores. Foliola calycina obovata, subinde mucronulata, apice albida, basi quinquenervia. Petala emarginata. Stamina longitudine stylorum vel duplò breviora. Stigmata capitato-obtusa. Capsula obtusiuscula.

- angusti-* 3. LINUM foliolis calycinis ellipticis subtrinervi-
folium. bus capsulisque acuminatis, foliis lineari-lan-
 ceolatis trinervibus, caulibus numerosis.
 Linum angustifolium. *Huds.* 134. *With.* ed. 2.
 329. *Engl. Bot. t.* 381.
 L. tenuifolium. *With.* 323. *Sp. Pl.* 399 ζ.
 L. sylvestre angustifolium, floribus dilutè purpu-
 rascentibus vel carneis. *Raii Syn.* 362.
Angl. Narrow-leaved Pale Flax.

In pascuis arenosis vel calcareis, præcipuè mariti-
 mis.

About St. Ives and Truro in Cornwall plentifully.
Ray. In several parts of Kent. *Huds.* Near
 Walsingham, and in Gunton fields, Norfolk.
Rev. Mr. Bryant. At Darsham, Suffolk. *Mr.*
Davy.

Perennis. Fl. Julio.

Habitus præcedentis, sed caules pauciores, suberecti.
 Folia lineari-lanceolata, integerrima, glabra, trinervia,
 subinde quinquenervia. Ramuli paniculati, vix rac-
 mosi. Flores dilutè purpurei. Folia calycina ellip-
 tica, acuminata, obsoletè trinervia, nervis lateralibus
 post florescentiam præcipuè manifestis. Petala integra,
 vel subemarginata. Stamina brevia. Styli subclavati,
 vix capitati.

A L. tenuifolio vero Linnæi (*Hall. Hist. n.* 838) toto
 coelo diversum est, atque primo intuitu dignoscitur ca-
 lycibus nequaquam glandulosè ciliatis.

* * *Foliis oppositis.*

- catharti-* 4. LINUM foliis oppositis obovato-lanceolatis,
cum. caule supernè dichotomo, petalis acutis.
 Linum catharticum. *Sp. Pl.* 401. *Huds.* 134.
With. 323. *Relb.* 131. *Sibth.* 106. *Curt.*
Lond. fasc. 3. t. 19. *Mart. Fl. Rust. t.* 135.
Dicks. H. Sicc. fasc. 3. 7. *Engl. Bot. t.* 382.
 L. syl-

L. sylvestre catharticum. *Raii Syn.* 362. *Ger. em.* 560.

Angl. Purging Flax. Mill-mountain.

In pascuis elatis ficcioribus.

Annua. Fl. Junio—Augusto.

Radix exigua. Herba glabra, amara, cathartica. Caules foliosi, erecti, basi simplices, supernè dichotomi, multiflori. Folia opposita, subelliptico-lanceolata, obtusa, integerrima. Flores parvi, albi, ante expansionem nutantes. Folia calycina uninervia, ferrata. Petala acuta. Stamina monadelphæ. Stigmata capitata.

Obs. Lini genus vix minùs quam Geranium, cui affine est, staminibus monadelphis gaudet, ut etiam species plurimæ Oxalidis.

164. SIBBALDIA. *G. Pl.* 155. *Juss.* 337.
Gærtn. t. 73.

Cal. decemfidus. *Pet.* 5, calyci inserta. *Styli*
e latere germinis. *Semina* 5 in fundo calycis.

1. SIBBALDIA foliolis tridentatis cuneiformibus. *procumbens.*
Sibbaldia procumbens. *Sp. Pl.* 406. *Huds.* 136. *With.* 326. *Dickf. H. Sicc. fasc.* 10. 12. *Fl. Dan. t.* 32.

Fragariæ sylvestri affinis planta, flore luteo. *Sibb. Scot. p.* 2. 25. *t.* 6. *f.* 1.

Pentaphylloides pumila, foliis ternis ad extremitates trifidis. *Raii Syn.* 256.

Scotch Cinquefoil. *Pet. H. Brit. t.* 41. *f.* 7.
Angl. Procumbent Sibbaldia.

In alpinis Scotiæ copiosè, solo micaceo.
Near the top of Ben Lomond.

Perennis. Fl. Julio.

Radix lignosa, multiceps, extùs nigra. Caules brevissimi, foliosi. Folia longiùs petiolata, ternata; foliolis obcuneatis, pilosis, apice tridentatis. Stipulæ binæ, petiolo

petiolo adnatæ. Pedunculi corymbosi. Flores exigui, lutei. Calyx pilosus, persistens, laciniis alternis majoribus. Petala obovata, integra. Stamina brevissima. Styli quinque, rarius decem. Semina ovata, gibba. Plukenetii tab. 212. f. 3, mihi videtur ad *Potentillam subacaulem* pertinere.

VI. HEXAGYNIA.

165. DROSER. *G. Pl.* 154. *Juss.* 245.
Gærtn. t. 61.

Cal. quinquefidus. *Pet.* 5. *Capsula* unilocularis, trivalvis, supera, polysperma.

rotundi- 1. DROSER foliis orbiculatis radicalibus, scapo
folia. racemoso.

Drosera rotundifolia. *Sp. Pl.* 402. *Huds.* 135.
With. 323. *Relb.* 131. *Sibth.* 106. *Fl. Dan.*
t. 1028.

Ros Solis folio rotundo. *Raii Syn.* 356. *Ger.*
em. 1556.

β. *Rorella rotundifolia* perennis. *Raii Syn.* 356.
Angl. Round-leaved Sun-dew.

In paludibus turfosis.

Perennis. Fl. Julio, Augusto.

Radix fibrosa. *Folia* radicalia, petiolata, obovato-subrotunda, obtusissima, integerrima, concava, supra pilis rubris viscosis tecta. *Scapi* plures, erecti, palmares, teretes, glabri. *Racemus* terminalis, plerumque solitarius, recurvus, simplex. *Flores* pedicellati, erecti, albi. *Calycis* laciniæ ellipticæ, obtusæ. *Petala* ob-

ovata. Stamina petalis breviora. Styli sex. Stigmata clavata. Capsula trivalvis, valvulis medio seminiferis.

Quandoque caulescit, ex observationibus D. Williams, Bot. Prof. Reg. Oxonii.

2. DROSERA foliis obovatis radicalibus, scapo *longifolia*.
racemoso.

Drosera longifolia. *Sp. Pl.* 403. *Huds.* 135.

With. 324. *Relb.* 132. *Dicks.* *H. Sicc.*

fasc. 14. 9.

Ros Solis folio oblongo. *Raii Syn.* 356. *Ger.*

em. 1556. *Morif. sect.* 15. *t.* 4. *f.* 2.

β. *Rorella longifolia perennis.* *Raii Syn.* 356.

Angl. Long-leaved Sun-dew.

In paludibus cum priore.

Perennis. Fl. Julio, Augusto.

Habitus præcedentis, sed folia obovata, vel oblonga, obtusa. Variat stylis novem. *D. Sowerby.* Hæc quoque aliquando caulescit. *D. Williams.*

3. DROSERA foliis oblongis obtusis radicalibus, *anglica.*
floribus octogynis, capsulis quadrivalvibus.

Drosera anglica. *Huds.* 135. *With.* 324. *Dicks.*

H. Sicc. fasc. 2. 7. *Abbot.* 72.

Rorella longifolia maxima. *Raii Syn.* 356.

Ros Solis major longiore folio et erectiore. *Mo-*

ris. sect. 15. *t.* 4. *f.* 1.

Angl. Great Sun-dew.

In paludibus minùs frequens.

Perennis. Fl. Julio, Augusto.

Cum *D. longifolia* convenit formâ, sed duplò major est, foliis plerumque lineari-obovatis. Styli octo. Capsula quadrivalvis.

Hæ omnes quam maximè affines sunt.

VII. POLYGYNIA.

166. MYOSURUS. *G. Pl.* 155. *Juss.* 233.
Gærtn. t. 74.

Cal. pentaphyllus, basi appendiculatus. *Pet.* 5,
 ungue tubuloso, nectarifero. *Semina* nume-
 rosa, nuda.

minimus. 1. MYOSURUS.

Myosurus minimus. *Sp. Pl.* 407. *Huds.* 136.
With. 326. *Relb.* 132. *Sibth.* 107. *Ab-*
bot. 72. *Curt. Lond. fasc.* 4. *t.* 26. *Dicks.*
H. Sicc. fasc. 12. 18. *Engl. Bot. t.* 435. *Fl.*
Dan. t. 406.

Myosuros. *Raii Syn.* 251.

Cauda Muris. *Ger. em.* 426. *Tillands.* *lc.* 108.
Angl. Moufe-tail.

In arvis solo glareoso.

Annua. *Fl.* Maio.

Radix fibrosa, parva. *Folia* radicalia, erectiuscula, line-
 ari-spatulata, obtusiuscula, integerrima, subcarnosa,
 glabra, subinde, monente Hudsono, hirsuta. *Scapi*
 plures, erecti, simplices, foliis longiores, teretes, superne
 incrassati, uniflori. *Flos* parvus, erectus, flavo-viridis.
Receptaculum subulatum, seminibus nudis numerosis-
 simis tectum.

Classis VI.

HEXANDRIA.

Stamina 6.

MONOGYNIA.

Stylus 1.

* *Fl. calyculati, calyce corollâque instructi.*

182. FRANKENIA. *Cor.* pentapetala. *Cal.* monophyllus, inferus. *Capsf.* unilocularis, polysperma.
181. BERBERIS. *Cor.* hexapetala. *Cal.* hexaphyllus, inferus. *Bacca* disperma.
183. PEPLIS. *Cor.* hexapetala. *Cal.* decemfidus. *Capsf.* bilocularis.

Lythrum hyssopifolium.

** *Fl. spathacei sive glumacei.*

168. LEUCOJUM. *Cor.* supera, hexapetala, campanulata. *Stam.* æqualia.
167. GALANTHUS. *Cor.* supera, hexapetala; petalis tribus interioribus brevioribus, emarginatis.
169. NARCISSUS. *Cor.* supera, hexapetala. *Nectarium* campanulatum, petaliferum, stamina includens.
170. ALLIUM. *Cor.* infera, hexapetala; petalis ovatis, sessilibus.

*** *Fl.*

* * * *Fl. nudi.*

173. CONVALLARIA. *Cor.* infera, sexfida.
Bacca trilocularis. *Stigma* trigonum.
177. ASPARAGUS. *Cor.* infera, sexpartita.
Bacca trilocularis. *Stigmata* tria.
175. ANTHERICUM. *Cor.* infera, hexapetala,
 patens. *Sem.* angulata.
176. NARTHECIUM. *Cor.* infera, hexapetala,
 patens. *Sem.* appendiculata. *Stam.* hirsuta.
173. ORNITHOGALUM. *Cor.* infera, hexapetala. *Stam.* basi dilatata.
174. SCILLA. *Cor.* infera, hexapetala. *Stam.* filiformia.
171. FRITILLARIA. *Cor.* infera, ovata, hexapetala, basi foveâ nectariferâ.
172. TULIPA. *Cor.* infera, campanulata, hexapetala. *Stylus* nullus. *Sem.* plana.
179. ACORUS. *Cor.* infera, hexapetala. *Spadix* multiflorus.

* * * * *Fl. incompleti.*

180. JUNCUS. *Cal.* hexaphyllus. *Caps.* superâ, trivalvis, polysperma.
Peplis Portula.

TRIGYNIA.

Styli 3.

187. COLCHICUM. *Cal.* spatha. *Cor.* hexa-
petaloidea.
186. TRIGLOCHIN. *Cal.* triphyllus. *Pet.*
tria. *Capsf.* bafi dehiscens.
184. RUMEX. *Cal.* triphyllus. *Pet.* tria. *Sem.*
unicum, triquetrum.
185. TOFIELDIA. *Cal.* trifidus. *Pet.* sex. *Cap-*
fulæ tres, polyspermæ.

POLYGYNIA.

Styli plures.

188. ALISMA. *Cal.* triphyllus. *Pet.* tria. *Capsf.*
plures, aggregatæ, monospermæ.

VI. HEXANDRIA.

I. MONOGYNIA.

167. GALANTHUS. *G. Pl.* 160. *Juss.* 55.

Cor. supera, hexapetala; petalis tribus interioribus brevioribus, emarginatis. *Stigma* simplex.

nivalis. I. GALANTHUS.

Galanthus nivalis. *Sp. Pl.* 413. *Huds.* 140.

With. 331. *Engl. Bot. t.* 19. *Dicks. H. Sicc. fasc.* 6. 2. *Abbot.* 73. *Jacq. Austr. t.* 313.

Leucoium bulbosum præcox minus. *Ger. em.* 147.

Narcissus 6. *Camer. Epit.* 956.

Angl. Snowdrop.

In pratis, pomariis, et ad fepes.

At the foot of Malvern hills, Worcestershire. *Mr.*

Ballard. On the banks of the Tees about Blackwell and Conniscliffe, certainly wild. *Mr. Robson.* Bedfordshire. *Abbot.*

Perennis. Fl. Februario.

Bulbus ovatus. *Folia* bina, obtusa, carinata, glauca, basi membranâ integrâ vaginatâ. *Scapus* teres, nudus, uniflorus. *Spatha* cylindrica, apice sæpius bifido. *Flos* pedicellatus, nutans, niveus, inodorus. *Petala* obovata, concava, carnosâ; interiora intus viridi striata, apiceque extus maculata. *Stamina* ex apice germinis. *Antheræ* subulatæ, apice poris duobus. *Stigma* acutum. *Capsula* trilocularis, trivalvis. *Semina* globosa.

168. LEUCOJUM. *G. Pl.* 160. *Juss.* 55.

Cor. supera, hexapetala, campanulata; petalis apice incrassatis. *Stigma* simplex. *Stamina* æqualia.

I. LEU-

1. LEUCOJUM spathâ multiflorâ, stylo clavato. *æstivum*.

Leucojum æstivum. Sp. Pl. 414. With. 331.

Curt. Lond. fasc. 5. t. 23. Jacq. Austr.

t. 203. Engl. Bot. t. 621.

L. bulbosum inajus polyanthemum. Ger. em. 148.

L. polyanthemum. Rencalm. Spec. 100. f. 1.

Leuconarcissifolium pratense vernal. Lob.

ic. 122.

Angl. Summer Snowflake.

In palustribus ad ripas fluviorum, pratibusque humidis.

Close by the Thames side between Greenwich and Woolwich, undoubtedly wild; also in the Isle of Dogs. *Curtis.* In a small island in the river, about three miles south of Kendal; Mr. Gough. *With.* In pastures at Little Stonham, Suffolk, a troublesome weed. *Mrs. Cabbold.* Near Reading. *Mr. Murray.*

Perennis. Fl. Maio.

Bulbus subrotundus. Folia plurima, erecta, sesquipedalia, obtusa, carinata, lætè viridia. Scapus longitudine foliorum, anceps, multiflorus. Spatha lanceolata, erecta. Pedicelli uniflori. Flores nutantes, nivei; petalis ovatis, apice viridi maculatis. Antheræ obtusæ, poris duobus. Stylus clavatus. Stigma acutum. Capsula elliptica, trilobularis. Semina globosa, magna.

169. NARCISSUS. *G. Pl. 161. Juss. 55.*

Cor. supera, hexapetala, æqualis. Nectarium infundibuliforme, monophyllum, petaliferum. Stamina intra nectarium. Stigma trifidum.

1. NARCISSUS spathâ uniflorâ, nectario rotato *poeticus*, brevissimo membranaceo crenato, foliis obtusè carinatis margine reflexis.

Narcissus poeticus. *Sp. Pl.* 414. *Ait. Hort. Kew.* v. 1. 408. *Engl. Bot.* t. 275. *With.* 331.

N. patellaris. *Salisb. Prod.* 225.

N. medio purpureus. *Ger. em.* 123. *Camer. Epit.* 951.

Angl. Poetic Narcissus.

In ericetis vel campis elatis, solo arenoso.

At Woodbastwick, and in other parts of Norfolk.

Mr. Rose. On a rabbit-warren at Shorne, between Gravesend and Rochester. *Mr. Rayer.*

Perennis. Fl. Maio.

Bulbus ovatus. Folia semiunciam lata, obtusiuscula, canaliculata, obtusissimè carinata, margine reflexa. Scapus anceps, rectus, nervosus, uniflorus. Spatha membranacea, sæpiùs biloba. Flos erectiusculus, magnus, odoratus, niveus, nectarii margine sanguineo.

biflorus. 2. **NARCISSUS** spathâ biflorâ, nectario rotato brevissimo membranaceo crenato, foliis acutè carinatis margine inflexis.

Narcissus biflorus. *Curt. Mag.* t. 197. *Engl. Bot.* t. 276. *With.* 332.

N. poeticus. *Huds.* 141. *With. ed.* 2. 341. *Wade Dubl.* 96.

N. cothurnalis. *Salisb. Prod.* 225.

N. medio luteus vulgaris. *Raii Syn.* 371. *Ger. em.* 124.

Angl. Pale Narcissus. Primrose peerless.

In campis arenosis.

On a rabbit-warren at Shorne with the preceding.

Mr. Rayer. Near Hornsey church. *Sherard.*

Common in the meadows about Dublin. *Mr.*

De Luc junior.

Perennis. Fl. Aprili, Maio.

Bulbus ovatus. Folia acutè carinata, margine inflexa. Scapus

Scapus anceps, nervosus, biflorus, rarius uni- vel triflorus. Spatha membranacea, indivisa. Flores nutantes, præcedentè minores, minùsque gratè odori, petalis ochroleucis, nectarii margine albido:

3. NARCISSUS spathâ uniflorâ, nectario campa- *Pseudonulato erecto crispò obsoletè sexfido petalis narcissus.*
ovatis æquali.

Narcissus Pseudo-narcissus. *Sp. Pl.* 414. *Huds.*

141. *With.* 332. *Sibth.* 109. *Engl. Bot.*

t. 17. *Abbot.* 73.

N. festalis. *Salisb. Prod.* 220.

N. sylvestris pallidus calyce luteo. *Raii Syn.*

371.

Pseudonarcissus anglicus. *Ger. em.* 133.

Angl. Common Daffodil.

In sylvis et dumetis humidiusculis.

Perennis. Fl. Martio.

Bulbus subrotundus. Folia glaucescentia, obtusè carinata, margine planiuscula. Scapus anceps, compressus, uniflorus. Spatha indivisa, germiui approximata. Flos nutans, magnus, ingrâtè odoratus. Petala erectiuscula, pallidè flava. Nectarium campanulatum, aureum, longitudine petalorum, supernè plicatum, apice sexfidum, crenatum. Germen trifidum.

170. ALLIUM. *G. Pl.* 163. *Juss.* 53.

Gærtn. t. 16.

Cor. infera, hexapetala, patens. *Spatha* bifida, multiflora. *Umbella* congesta. *Stigma* simplex.

* *Foliis caulinis, planis.*

1. ALLIUM caule planifolio umbellifero, umbellâ *Ampeloglobosâ*, staminibus tricuspидatis, petalis dorso *præsum.*
scabris.

A a 2

Allium

Allium Ampeloprasum. Sp. Pl. 423. Hudf. 138.
With. 332.

A. Holmense, sphærico capite. *Raii Syn. 370.*

A. sphærico capite, folio latiore. *Rudb. El.*
v. 2. 151. f. 1.

Scorodoprasum primum Clusii. *Ger. em. 180.*

Angl. Great Round-headed Garlick.

In collibus apricis rarissimè.

On the Holms Island in the mouth of the Severn.

Mr. Newton. *Ray.*

Perennis. Fl. Julio.

Caulis erectus, teres, basi foliosus. Folia unciam ferè
 lata, plana, margine denticulato-scabra. Umbella ter-
 minalis, multiflora, globosa. Pedunculi simplices, ses-
 quiunciales, angulati, glabri, squamulis membranaceis
 lanceolatis basi interstinctis. Flores pallidi, campanu-
 lati, petalis ovatis, acutis, dorso scabris. Stamina dila-
 tata, membranacea, omnia apice tricuspidata. Antheræ
 terminales, erectæ. Germen ovatum. Stigma obtusum.
Herb. Buddl.

arenari- 2. ALLIUM caule planifolio bulbifero, vaginis
um. teretibus, spathâ muticâ, staminibus tricuspi-
 datis, petalis carinâ scabriusculis.

Allium arenarium. Sp. Pl. 426. Hudf. 138.
With. 333.

A. Scorodoprasum. *Fl. Dan. t. 290.*

A. sylvestre amphicarpon, foliis porraceis, flori-
 bus et nucleis purpureis. *Raii Syn. 370.*

Porrum planifolium, staminibus alternè trifidis,
 umbellâ bulbiferâ. *Rupp. Jen. ed. Hall. 154.*
t. 2. f. 2.

Angl. Sand Garlick.

In nemorosis et montosis Angliæ borealis.

About Lowther, Westmoreland. *Mr. Crowe and*
Mr.

Mr. Woodward. In a meadow called Santines at Castle Howard, Yorkshire. *Mr. Teesdale.* About Thorp-Arch. *Rev. Mr. Wood.* In Heslington fields near York. *Rev. Archdeacon Peirson.*

Perennis. Fl. Julio.

Bulbus ovatus, fuscus. Caulis erectus, tripedalis, teres, glaber, basi foliosus. Folia vix semiunciam lata, margine scabra, vaginâ longissimâ, tereti. Umbella parva, globosa, densa. Spatha brevis. Pedunculi incurvati, breves, bulbis interstinctis. Flores rubri, campanulati, petalis carinâ tantum scabriusculis. Stamina alterna dilatata, tricuspidata, intermediis simplicibus, subulatis. Germin triquetrum.

3. ALLIUM caule planifolio bulbifero, stamini-
bus omnibus subulatis, spathâ acutâ. carina-
tum.

Allium carinatum. *Sp. Pl.* 426. *Huds.* 139.

With. 333.

A. inodorum, umbellâ bulbiferâ, vaginâ bicorni, foliis carinatis. *Hall. Opusc.* 391. t. 2. f. 2.

A. montanum bicorni purpureum proliferum.

Raii Syn. 369.

Ampeloprasion proliferum. *Lob. Ic.* 156.

Moly montanum secundum Clusii. *Ger. em.* 187.

Angl. Mountain Garlick.

In rupibus et montosis Angliæ borealis.

Near Settle, Yorkshire. *Ray.* On the rocks about Longfleadale, Westmoreland. *Richardson.* Near Ramsgate, in the Isle of Thanet, and between Sandwich and Deal. *Huds.* In Heslington fields near York, and about Knareborough. *Rev. Archdeacon Peirson.*

Perennis. Fl. Julio.

Caulis erectus, tripedalis, teres, glaber, ad medium usque
A a 3 foliosus.

foliosus. Folia præcedente duplò angustiora, plana, succosa, nervosa, obtusa, margine scabra, vaginâ longissimâ. Umbella globosa, densa. Spatha diphylla, longissima; foliacea, patens. Pedunculi vix unciales, bulbis interstinctis. Flores ochroleuci, carinâ virente, glabrâ. Stamina omnia subulata, simplicia, æqualia. Descripsi e plantâ hortensi, cum nondum vidi spontaneam.

* * *Foliis caulinis, teretibus.*

oleraceum. 4. ALLIUM caule teretifolio bulbifero, foliis scabris canaliculatis subtùs sulcatis, staminibus simplicibus.

Allium oleraceum. *Sp. Pl.* 429. *Huds.* 140. *With.* 334. *Engl. Bot. t.* 488.

A. sylvestre bicorni, flore ex herbaceo albicante, cum triplici in singulis petalis striâ atro-purpureâ. *Raii Syn.* 370.

A. umbellâ bulbiferâ, vaginâ bicorni, foliis teretibus. *Hall. Opusc.* 387, t. 1. f. 2.

Angl. Streaked Field Garlick.

Inter segetes et ad margines agrorum.

At Black Notley, Essex. *Ray.* In several places in Westmoreland, and near Bristol. *Huds.* At Fincham, Norfolk. *Rev. Mr. Forby,* Near York. *Sir T. Frankland.*

Perennis. Fl. Julio.

Bulbus ovatus. Caulis erectus, bi- vel tripedalis, gracilis, teres, foliosus. Folia angusta, acuta, saturatè viridia, fistulosa; supernè complanata vel canaliculata; subtus convexa, nervosa, utrinque scabriuscula. Umbella multibulbosa, pedunculis paucioribus, elongatis, divaricatis, unifloris. Spatha diphylla; foliolis basi ventricosis, apice attenuatis, acutis, umbellam longè superantibus. Flores campanulati. Petala glabra, pallidè viridia, lineis tribus purpurascens. Stamina æqualia, simplicia, subulata, basi connexa. Germen apice scabrum.

5. *ALLIUM* caule teretifolio bulbifero, flammini-*vincale*.
bus tricuspидatis.

Allium vincale. *Sp. Pl.* 428. *Huds.* 139.
With. 333. *Relb.* 135. *Sibth.* 109. *Ab-*
bot. 74.

A. sylvestre. *Raii Syn.* 369. *Ger. em.* 179.

A. filvestre tenuifolium. *Lob. Ic.* 156.

Angl. Crow Garlick.

In pascuis aridis, muris antiquis, et inter segetes,
præcipuè solo calcareo.

Perennis. Fl. Julio.

Bulbus ovatus, parvus. Caulis gracilis, striatus, erectus,
foliosus. Folia fistulosa, teretia, nervosa, lævia. Um-
bella parva. Pedunculi breves, capillares, bulbis inter-
stinctis numerosis, obovatis, divaricatis, mucronatis.
Spatha decidua. Flores parvi, carnei. Stamina exserta,
alterna tricuspидata, setis lateralibus antheram super-
antibus.

* * * *Scapo nudo.*

6. *ALLIUM* scapo nudo semicylindrico, foliis *urfinum*.
lanceolatis petiolatis, umbellâ fastigiata.

Allium urfinum. *Sp. Pl.* 431. *Huds.* 140.
With. 334. *Relb.* 135. *Sibth.* 110. *Ab-*
bot. 74. *Engl. Bot. t.* 122. *Fl. Dan. t.* 757.
Fuchs. Hist. 739. *Ger. em.* 179. *Dicks. H.*
Sicc. fasc. 7. 5.

A. sylvestre latifolium. *Raii Syn.* 370.

Angl. Broad-leaved Garlick. *Ramsons.*

In nemorosis, pratisque humidis, frequens.

Perennis. Fl. Maio, Junio.

Bulbus oblongus, attenuatus. Folia omnia radicalia, pe-
tiolis vaginantibus, ovata, acuta, integerrima, glabra,
nervosa. Scapus foliis longior, erectus, simplex, gla-
ber, semicylindricus, vix triqueter. Umbella multi-
flora, planiuscula, bulbis destituta. Spatha ovata, pe-
dunculis brevior. Flores nivci. Petala ovata, paten-

tia. Stamina simplicia, subulata, æqualia. Germen turbinatum, trilobum, glabrum. Herba odore alliaceo vehementiore gaudet, at pecoribus armentisque pabulum non ingratum præbet. Hinc lac nauseosum foetidumque evadit.

Schæno- 7. ALLIUM scapo nudo tereti folia adæquante,
præsum. foliis teretibus subulato-filiformibus.

Allium Schænopræsum. Sp. Pl. 432. Hudf. 649.

With. 335. Lightf. 180. Fl. Dan. t. 971.

Schænopræson. Ger. em. 176.

Porrum sectivum. Fuchs. Hist. 635.

Angl. Chive Garlick.

In pratis et pascuis rariùs.

By Fast castle, on the borders of Berwickshire,
Dr. Parsons; also in Argylshire. *Lightf. West-*
moreland. Hudf. With.

Perennis. Fl. Junio.

Bulbi cylindrici, graciles, cæspitosi. Folia erecto-patentia, vaginantia, cylindrica, acuta, glabra. Scapus spithamæus, teres, lævis. Umbella globosa, densa, multiflora. Spatha ovata, floribus brevior. Petala patentia, lanceolata, alba vel incarnata, costâ purpurascente. Filamenta simplicia, subulata. Plantam in Britannia spontaneam nondum inveni; in hortis rusticis frequentissimè colitur. Ex exemplaribus in Helvetiâ natis descriptionem confeci.

171. FRITILLARIA. *G. Pl. 164. Juss. 48.*
Gærtn. t. 17.

Cor. infera, hexapetala, campanulata, supra ungues cavitate nectariferâ. Stamina longitudine corollæ. Cal. nullus. Semina plana.

Meleagris. 1. FRITILLARIA foliis omnibus alternis, caule
gris. unifloro, nectario lineari.

Fritillaria Meleagris. Sp. Pl. 436. Hudf. 144.

With.

With. 335. *Relb.* 137. *Sibth.* 110. *Abbot.* 75. *Curt. Lond. fasc.* 3. t. 20. *Engl. Bot.* t. 622. *Dicks. H. Sicc. fasc.* 9. 3. *Fl. Dan.* t. 972. *Jacq. Austr. app.* t. 32.

F. præcox purpurea variegata. *Rudb. Elys.* v. 2. 113. f. 1.

F. variegata. *Ger. em.* 149.

Angl. Common Fritillary. Checquered Daffodil. Snake's-head.

In pratis et pascuis humidis.

Maud Fields near Rislip Common, Middlesex; Mr. Ashby. *Blackstone.* Between Mortlake and Kew, and near Enfield. *Huds.* Near Laxfield, Suffolk. Mr. Woodward. At Little Stonham in the same county. Mrs. Cobbold. Near Reading. Mr. Murray.

Perennis. Fl. April.

Bulbi parvi, subrotundi, aggregati. Caulis erectus, pedalis, simplicissimus, teres, glaber, foliosus, uniflorus. Folia alterna, lineari-lanceolata, acutiuscula. Flos terminalis; cernuus, magnitudine nucis juglandis, campanulatus, purpureo tessellatus, quandoque albus. Nectarium foveâ oblongâ, extûs gibbâ. Antheræ erectæ, versatiles. Germen sexsulcatum. Stigma tripartitum.

172. TULIPA. *G. Pl.* 165. *Juss.* 48.

Gærtn. t. 17.

Cor. hexapetala, campanulata. *Stylus* nullus.

Capsula supera, trilocularis. *Semina* plana.

Cal. nullus.

1. TULIPA flore solitario subnutante, foliis lanceolatis, stigmate obtuso triquetro, staminibus *trifloris*.
basî hirsutis.

Tulipa sylvestris. *Sp. Pl.* 438. *Engl. Bot.*

t. 63.

t. 63. *With.* 336. *Abbot.* 75. *Fl. Dan.*

t. 375.

T. bononiensis. *Ger. em.* 138.

Angl. Wild Tulip.

In clivis cretaceis.

Old chalk-pits near Norwich. *Mr. Rose.* Near Bury. *Sir T. G. Cullum, Bart.* and *W. Mathew, Esq.* At Whipnade, on the borders of Hertfordshire and Bedfordshire; *Rev. D. Jenks. Abbot.*

Perennis. *Fl.* Aprili.

Bulbus ovatus, gibbus. Caulis simplicissimus, erectiusculus, teres, glaber, uniflorus, medio foliosus, basi attenuatus. Folia alterna, subamplexicaulia, lanceolata, acuta, carinata, glaucescentia. Flos terminalis, subnutans, suaveolens, flavus, extus viridescens. Petala elliptico-lanceolata, nectario destituta. Stamina complanata, basi barbata. Antheræ terminales, versatiles, oblongæ, flavæ, polline flavo. Germen trigonum. Stigma sessile, obtusum, trigonum, nec dilatatum.

173. ORNITHOGALUM. *G. Pl.* 166.

Juss. 53. *Gærtn. t.* 17.

Cor. hexapetala, persistens, supra medium patens. *Filamenta* basi dilatata. *Capsula* supera, trilocularis. *Semina* subrotunda. *Cal.* nullus.

luteum. 1. ORNITHOGALUM caule anguloso diphylo, pedunculis umbellatis simplicibus.

Ornithogalum luteum. *Sp. Pl.* 439. *Huds.* 143.

With. 336. *Sibth.* 111. *Engl. Bot. t.* 21,

Raii Syn. 372. *Fl. Dan. t.* 378.

O. luteum, five Cepe agraria. *Ger. em.* 165.

Ornithogalum. *Tillands. Ic.* 151.

Bulbus sylvestris. *Fuchs. Hist.* 169.

Angl. Yellow Star of Bethlehem.

In

In nemorosis rariùs.

In woods about Oxford. *Sir J. Banks, Bart.*
 Yorkshire. *Mr. Salisbury.* In a meadow near
 the copper-mills, Derby. *Mr. Whateley.*

Perennis. Fl. Aprili.

Bulbus parvus. Caulis spithamæus, erectus, triqueter.
 Folia carinata, lineari-lanceolata; radicalia plerumque
 solitaria; caulina floribus approximata, bina, inæqualia,
 ciliata. Pedunculi terminales, umbellati, simplices,
 uniflori. Petala obtusa, flava, viridi lineata, marces-
 centia. Stamina subulata, uniformia. Germen trigo-
 num. Stylus triqueter. Stigma obtusum.

2. ORNITHOGALUM racemo longissimo, fila-^{pyrenai-}
 mentis omnibus dilatatis, pedunculis floriferis ^{cum.}
 patentibus æqualibus: fructiferis scapo ap-
 proximatis.

Ornithogalum pyrenaicum. *Sp. Pl.* 440. *Huds.*
 143. *With.* 337. *Abbot.* 76. *Engl. Bot.*
 t. 499. *Jacq. Austr. t.* 103.

O. angustifolium majus, floribus ex albo vires-
 centibus. *Raii Syn.* 372.

Asphodelus bulbosus. *Ger. em.* 97.

Angl. Spiked Star of Bethlehem.

In pascuis, collibusve, rariùs.

In Suffex and Somersethire. *Ray.* Keyson Park
 Wood, Bedfordshire. *Rev. T. O. Marsh.*

Perennis. Fl. Junio, Julio.

Bulbus ovatus. Folia radicalia, longissima, canaliculata.
 Scapus teres, subtripedalis, erectus, multiflorus. Flores
 racemosi, pedicellati, albido-virescentes. Pedunculi vix
 unciales, sub florescentiâ patentes, dein erecti, bractea
 solitaria, lanceolata, acuminata, persistenti, ad basin.
 Petala patentissima, lineari-oblonga. Stamina ad me-
 dium usque dilatata, apice subulata. Antheræ versa-
 tiles, Stigma obtusè triquetrum.

umbellatum. 3. ORNITHOGALUM floribus corymbosis, pedunculis scapum superantibus, filamentis subulato-dilatatis integerrimis.

Ornithogalum umbellatum. *Sp. Pl.* 441. *Huds.* 143. *With.* 337. *Sibth.* 111. *Abbot.* 76. *Engl. Bot. t.* 130. *Jacq. Austr. t.* 343.

O. vulgare et verius, majus et minus. *Raii Syn.* 372.

Ornithogalum. *Ger. em.* 165.

Angl. Common Star of Bethlehem.

In pratis, pascuis et nemorosis.

At Babergh, Norfolk. *Mr. Wagstaffe.* Plentiful in a field at Knareborough, Yorkshire. *Mr. Robson.* On Everton heath, Bedfordshire. *Abbot.*

At Little Stonham, Suffolk. *Mrs. Cobbold.*

Perennis. Fl. Aprili, Maio.

Bulbus albus, subrotundus. Folia radicalia, linearia, canaliculata, apice emarcida. Scapus teres. Pedunculi terminales, alterni, corymbosi, bractea solitaria, elongata, ad basin. Petala patentia, elliptico-lanceolata; supra candida, nitida; subtus dorso viridia. Stamina omnia æqualia, complanato-subulata, integerrima, nec emarginata. Germen turbinatum. Stylus brevis.

174. SCILLA. *G. Pl.* 166. *Juss.* 53.

Cor. hexapetala, patens. *Filamenta* filiformia.

Capsula supera, trilocularis. *Semina* subrotunda. *Cal.* nullus.

verna. 1. SCILLA radice solidâ, corymbo hemisphærico paucifloro, bracteis lanceolatis obtusis, foliis linearibus canaliculatis.

Scilla verna. *Huds.* 142. *With.* 338. *Engl. Bot. t.* 23. *Dicks. H. Sicc. fasc.* 10. 8.

S. bifolia. *Lightf.* 181. *Fl. Dan. t.* 568.

Hya-

Hyacinthus n. 1. *Raii Syn.* 372, synonymis erroneis.

H. stellaris vernus pumilus. *Dill. in Raii Syn. Indic. Pl. dub.*

Angl. Vernal Squill.

In clivis maritimis.

In the Isle of Man, and Anglesey, also on the coasts of Ireland, Wales and Cornwall. *Dill.* and *Huds.* In the islands of Iona and Staffa abundantly. *Lightf.*

Perennis. Fl. Aprili.

Bulbus ovatus, parvus. Folia plurima, linearia, canaliculata, basi attenuata. Scapus teres, erectus. Corymbus terminalis, hæmisphæricus. Flores a tribus ad sex, rariùs plures, pedicellati, cærulei. Bractææ longitudine pedicellorum, persistentes, obtusæ. Petala ovata, carinata. Stamina teretia, antheris cæruleis. Germen subrotundum. Stylus brevis. Stigma obtusum.

2. SCILLA radice solidâ, floribus corymboso-*bifolia.* racemosis cbracteatiss erectiusculis, foliis lanceolatis binis.

Scilla bifolia. *Sp. Pl.* 443. *Engl. Bot. t.* 24.

With. 338. *Jacq. Austr. t.* 117.

Hyacinthus stellatus Fuchsi. *Ger. em.* 106.

H. cæruleus mas minor. *Fuchsf. Hist.* 837, et albicans seu fœminea. *Ibid.* 838.

H. stellaris bifolius Germanicus. *Rudb. Elys. v.* 2. 33. f. 1.

Star Hyacinth. *Pet. H. Brit. t.* 67. f. 5.

Narcissus martius. *Brunf. Herb. v.* 1. 184.

Angl. Two-leaved Squill.

In nemorosis Angliæ occidentalis.

Received from the west of England by Mr. Sims, druggist, of Norwich. It is also in Buddle's herbarium.

Perennis.

Perennis. Fl. Martio, Aprili.

Bulbus subovatus. Folia bina, rarius terna, lanceolata, obtusiuscula, subcarinata. Scapus foliis parum altior, erectus, teres. Corymbus terminalis, subracemosus. Flores a quatuor ad decem, longius pedicellati, erectiusculi, ebracteati, cærulei. Petala obtusiuscula. Antheræ subfuscae.

*autum-
nalis.*

3. *SCILLA* foliis linearibus, floribus corymboso-racemosis, pedunculis ebracteatis adscendentibus longitudine florum.

Scilla autumnalis. *Sp. Pl.* 443. *Huds.* 142. *With.* 338. *Engl. Bot. t.* 78. *Curt. Lond. fasc.* 6. *t.* 25.

Hyacinthus autumnalis minor. *Raii Syn.* 373. *Ger. em.* 110.

Angl. Autumnal Squill.

In pascuis ficcis sterilioribus.

On Black-heath, Moulsey-burft, Kew green, and several similar places near London. *Huds.* *Curt.*
On St. Vincent's rocks, Bristol.

Perennis. Fl. Septembri.

Bulbus ovato-subrotundus. Folia linearia, canaliculata, patentia, numerosa. Scapus foliis duplo longior, erectus, teres, basi attenuatus. Flores racemosi, subcorymbosi, rosei, germine flaminibusque cærulescentibus. Pedunculi breves, ebracteati, patentes, sursum curvati.

nutans.

4. *SCILLA* foliis linearibus, spicâ nutante, floribus cernuis campanulato-cylindricis apice reflexis, bracteis geminis.

Scilla nutans. *Engl. Bot. t.* 377.

S. festalis. *Salisb. Prod.* 242.

Hyacinthus non-scriptus. *Sp. Pl.* 453. *Huds.* 141. *With.* 343. *Relb.* 136. *Sibth.* 110. *Curt. Lond. fasc.* 2. *t.* 18.

H. anglicus. *Raii Syn.* 373. *Ger. em.* 111.

Angl. Hare-bell Squill. Wild Hyacinth.

In nemorosis, dumetis, pascuis, et ad sepes, copiosè.

Perennis. Fl. Maio.

Bulbus ovatus, albus. Folia plurima, linearia, carinata, flaccida. Scapus pedalis, teres, erectus. Spica terminalis, nutans, multiflora; fructifera erecta. Flores pedicellati, cernui, secundi, cærulei, quandoque albi, suaveolentes, campanulati; basi subcylindrici. Bractæ binæ, lineares, acutæ. Petala lineari-lanceolata, ad medium usque approximata, apice reflexo-patentia, persistencia. Stamina petalis adnata, filiformia, æqualia. Germen hexagonum, poris nectariferis planè destitutum. Stigma obtusum.

Dubii generis herba gaudet floribus campanulatis, tametsi non moris petalis, ut in Hyacintho; structurâ verò partium, figurâ staminum, totoque habitu Scilla est; nec obstat corolla, in hac specie, ut in *S. peruvianâ* Linn. & *S. campanulatâ* Hort. Kew. persistens.

175. **ANTHERICUM.** *G.Pl.* 167. *Gærtn. t.* 16. *Phalangium.* *Juss.* 52.

Cor. hexapetala, patens, persistens. *Filamenta* filiformia. *Capsula* supera. *Semina* angulata. *Cal.* nullus.

1. **ANTHERICUM** foliis semicylindricis; caulibus basi dilatatis, caule subunifloro. *serotinum.*

Anthericum serotinum. *Sp. Pl.* 444. *Huds.* 144. *With.* 339. *Jacq. Austr. app. t.* 38.

Bulbosa alpina juncifolia, pericarpio unico erecto in summo cauliculo dodrantali. *Raii Syn.* 374. *t.* 17. *f.* 1.

Pseudo narcissus gramineo folio. *Rudb. Elys.* v. 2. 64. *f.* 9.

Angl. Mountain Spiderwort.

In excelsis rupibus alpinis Cambrobritannicis rariùs.
On

On Snowdon and other mountains in Wales. *Ray*.
 On Crib y Ddewil, near Llanberris, rare; on
 rocks above Cwm Idwell, Caernarvonshire, in
 abundance; Mr. Griffith. *With*.

Perennis. Fl. Junio.

Bulbus oblongus, curvus. Caulis palmaris, erectus, teres,
 glaber, foliosus, plerumque uniflorus, apice substriatus.
 Folia radicalia caulem superantia, semicylindrico-fili-
 formia, acutiuscula, glabra; caulina tria vel quatuor,
 alterna, brevia, basi dilatata, nervosa. Flos plerum-
 que solitarius, terminalis, erectus; intus albus; extus
 rubicundus. Petala elliptica, æqualia, nervosa, basi
 attenuata, emarcida. Stamina filiformia, glabra, per-
 sistentia. Antheræ subrotundæ, pallidæ. Capsula sub-
 turbinata, membranacea, trilocularis, stylo persistente
 coronata. Semina incurva, triquetra, striata.

176. NARTHECIUM. *Huds.* 145. *Mæring*.
Gorter.

Anthericum. *G. Pl.* 167.

Cor. hexapetala, patens, persistens. *Filamenta*
 filiformia, hirsuta. *Capsula* supera, prisma-
 tica. *Semina* utrinque appendiculata. *Cal.*
 nullus.

*offifra-
gum.*

1. NARTHECIUM. *Mæring. in Eph. Nat.*
*Cur. v. 6. 389. t. 5. f. 1. Gorter. Fl. Gelro-
zutph. 69.*

Narthecium offifragum. *Huds.* 145. *With.* 339.
Engl. Bot. t. 535.

Anthericum offifragum. *Sp. Pl.* 446. *Fl. Dan.*
t. 42. Dickf. H. Sicc. fasc. 1. 5. Relb. 138.
Abbot. 77.

Asphodelus Lancastriæ. *Ger. em.* 95.

A. minimus luteus acorifolius palustris. *Lob.*
 Ic. 92.

Phalan-

Phalangium Anglicum palustre Iridis folio. Raii

Syn. 375.

Angl. Lancashire Asphodel.

In paludibus turfosis.

Perennis. Fl. Julio, Augusto.

Radix fibrosa. Caules adscendentes, simplices, teretes, glabri, subflexuosi, parum foliosi. Folia equitantia, ensiformia, nervosa; caulina alterna, brevia, carinata; Spica terminalis, erecta, multiflora. Pedicelli alterni, erecti, uniflori, angulati, bracteati. Petala lineari-oblonga, nervosa, flava, extus viridia, persistentia. Stamina petalis opposita, parumque breviora, persistentia, filiformia, flava, villis flavescentibus densissimis vestita. Antheræ lineares, demum tortuosæ, flavæ. Germen oblongum, triquetrum, apice subulatum. Stigma capitatum, parvum, subsessile. Capsula subulato-prismatica, saturatè fulva; nitida, rigida, trivalvis, trilocularis; dissepimentis e medio valvularum. Semina numerosa, erectiuscula, imbricata, elliptico-teretia, utrinque in appendiculum subulatum attenuata.

177. ASPARAGUS. *G. Pl.* 168. *Juss.* 41.

Gærtn. t. 16.

Cor. sexfida, erecta, æqualis. *Cal.* nullus. *Stylus* brevissimus. *Stigmata* tria. *Bacca* supera, trilocularis. *Semina* bina, extus convexa.

1. ASPARAGUS caule herbaceo inermi suberecto *officinalis*-tereti, foliis setaceis mollibus, stipulis subsolitariis.

Asparagus officinalis. Sp. Pl. 448. *Huds.* 145.

With. 340. *Engl. Bot. t.* 339. *Fl. Dan. t.* 805.

Asparagus. Raii Syn. 267.

A. sativus. Ger. em. 1110.

β. *A. maritimus*, crassiore folio. *Dill. in Raii Syn.* 267.

Angl. Common Asparagus, or Sperage.

In petrosis maritimis.

In several places on the west and south coasts of England. Near Weymouth. *Mr. Lambert.*

β. Near Llanfachlog, Anglesea. *Mr. Lhwyd* and *Rev. H. Davies.*

Perennis. Fl. Augusto.

Radix repens, fibris longissimis, carnosis, rectis. Caules annui, erecti, (in β procumbentes,) teretes, ramosissimi, foliosi. Folia fasciculata, setacea, acuta. Stipulæ solitariae, membranaceæ, quandoque ternæ, interioribus minimis. Flores subaxillares, pedunculati, cernui, albidii. Corolla sexpartita, laciniis æqualibus, rectis, basin versus sæpe cohærentibus. Stamina corollæ inserta, brevissima, inclusa. Germen subrotundum, trifidum. Stylus vix ullus. Stigmata tria, maturefcente fructu ad basin usque separata. Bacca globosa, miniata, nitida, glabra. Semina nigra.

178. CONVALLARIA. *G. Pl.* 169. *Juss.* 42. *Gærtn. t.* 16.

Cor. sexfida. *Cal.* nullus. *Stylus* staminibus longior. *Stigma* trigonum. *Bacca* supera, trilocularis, ante maturitatem maculata.

majalis. 1. CONVALLARIA scapo nudo semicylindrico, floribus spicatis pedicellatis nutantibus.

Convallaria majalis. *Sp. Pl.* 451. *Huds.* 146. *With.* 341. *Relb.* 138. *Sibth.* 111. *Curt. Lond. fasc.* 5. *t.* 24. *Fl. Dan. t.* 854. *Abbot.* 76. *t.* 2.

Lilium convallium. *Raii Syn.* 264. *Ger. em.* 410. *Camer. Epit.* 618. *Brunf. Herb. v.* 1. 211.

Ephemerum non letale. *Fuchf. Hist.* 240.

Angl. Lily of the Valley.

In nemorosis ad radices montium.

Perennis.

Perennis. Fl. Maio.

Radix repens, implexa. Folia radicalia, bina, elliptica, nervosa, lævia, petiolis equitantibus, extus squamis vestitis. Scapus extrafoliaceus, solitarius, erectus, simplex, semicylindricus. Flores racemosi, pedicellati, alterni, bracteati, nutantes, campanulati, albi, odoratissimi, laciniis recurvis. Baccæ globosæ, miniatæ, glabræ, triloculares, loculis plerumque dispermis.

2. CONVALLARIA foliis verticillatis.

verticillata.

Convallaria verticillata. *Sp. Pl.* 451. *Engl. lata.*

Bot. t. 128. *Tr. of Linn. Soc. v.* 2. 357.

With. 341. *Fl. Dan. t.* 86.

Polygonatum n. 1244. *Hall. Hist. v.* 2. 120.

P. minus. *Ger. em.* 903.

P. tenuifolium. *Cord. Hist.* 113. 1.

P. angustifolium. *Fuchsf. Hist.* 586. *Ic.* 337.

Angl. Narrow-leaved Solomon's Seal.

In nemorosis ad radices montium Scoticarum rarior.

In Den Rechip, a deep woody valley between the hills of Stormont, Perthshire, about four miles north-east of Dunkeld. *Mr. A. Bruce.*

Perennis. Fl. Junio.

Radix repens, carnosa. Caules erecti, bipedales, simplices, foliosi, angulati, glabri, basi nudi. Folia verticillata, terna, quaterna, in apice caulis quina vel etiam septena, lanceolata, acuta, subtus glauca. Pedunculi axillares, solitarii, ramosi, nutantes. Flores cylindrici, lactei, apice virides, laciniis intus sub apice barbatis. Stamina brevia, fauce inserta, inclusa. Stylus brevis. Bacca globosa, cærulea.

3. CONVALLARIA foliis alternis amplexicaulis, caule angulato, pedunculis axillaribus subunifloris. *Polygonatum.*

Convallaria Polygonatum. *Sp. Pl.* 451. *Huds.*

146. *With.* 342. *Engl. Bot.* t. 280. *Woodv. Med. Bot.* t. 44. *Fl. Dan.* t. 377.

Polygonatum floribus ex singulis pediculis. Raii Syn. 263.

P. latifolium secundum Clusii. Ger. em. 904.

β. *P. Hellebori albi folio, caule purpurascente.*

Raii Syn. 263. non vidi.

Angl. Angular Solomon's Seal.

In nemorosis et dumetis rariùs.

In the mountainous or rocky woods of the north of England, especially about Settle and Ingleton, Yorkshire. Lately found near Yarmouth by *Mr. Wigg.* β. In woods on the north side of Mendip hills, Somersetshire. *Bobart.*

Perennis. Fl. Maio, Junio.

Radix repens, carnosa. Caules erecti, apice nutantes, sesquipedales, simplices, foliosi, angulati, subancipites, flexuosi, glabri. Folia alterna, elliptica, nervosa. Pedunculi axillares, solitarii vel bini, simplices, uniflori, rarissimè biflori. Flores cernui, oblongi, fragrantés, oehroleuci, basi apiceque virides, laciniis patulis, barbatis. Stamina medio tubi inserta, antheris ultra laciniarum basin prominentibus. Stylus semiuncialis. Bacca cærulea.

multiflo- 4. CONVALLARIA foliis alternis amplexicaulibus, caule tereti, pedunculis axillaribus multifloris.

Convallaria multiflora. Sp. Pl. 452. *Huds.* 147.

With. 342. *Abbot.* 77. *Engl. Bot.* t. 279.

Fl. Dan. t. 152. *Dreves. Bilderb.* t. 52.

Polygonatum. Raii Syn. 263. *Ger. em.* 903.

β. *P. humile anglicum. Raii Syn.* 263. non vidi.

Angl. Common Solomon's Seal.

In nemorosis et dumetis.

Perennis. Fl. Maio, Junio.

Radix

Radix repens, carnosa, sæpe denudata. Caules bipedales, simplices, teretes, foliosi, apice nutantes. Folia alterna, elliptica, nervosa, præcedente parùm pallidiora. Pedunculi axillares, solitarii, ramosi, subquinqueflori. Flores cernui, cylindrici, albi, basi apiceque virides, vix barbati. Stamina apice tubi inserta, brevissima. Stylus semiuncialis. Bacca cæruleo-nigra.

179. ACORUS. *G. Pl.* 172. *Juss.* 25.
Gærtn. t. 84.

Spadix cylindricus, tectus flosculis. Cor. hexapetala, nuda. Stylus nullus. Capsula trilobularis.

1. ACORUS scapi mucrone longissimo foliaceo. *Calamus.*
Acorus Calamus. Sp. Pl. 462. *Huds.* 147. *Witb.* 343. *Relb.* 140. *Sibth.* 112. *Abbot.* 77. *Dryand. in Ait. H. Kew.* v. 1. 474. *Woodv. Med. Bot. t.* 173. *Engl. Bot. t.* 356. *Leers.* 86. *t.* 13. *f.* 12.

A. verus, sive Calamus officinarum. Raii Syn. 437.

A. verus, officinis falsò Calamus. Ger. em. 62. *Acorum. Camer. Epit.* 5.

Calamus aromaticus, radix. Pharmac. Lond. Angl. Sweet Flag.

In aquosis, minùs frequens.

In the rivers of Norfolk plentiful. On Hillingdon common, Middlesex, and other places about London.

Perennis. Fl. Junio.

Radix crassa, horizontalis, repens. Folia radicalia, ensiformia, tripedalia. Scapus anceps, ultra medium florifer, apice foliaceus. Spadix nudus, lateralis, solitarius, cylindricus, obtusiusculus, multiflorus. Flores tessellato-congesti, virides. Petala erecta, regularia, apice inflexa. Stamina petalis alterna, filiformia. Germen

B b 3

ellipticum.

ellipticum. Stigma sessile. Herba tota aromatica, tonica, præcipuè radices.

180. JUNCUS. *G. Pl.* 173. *Juss.* 44.
Gærtn. t. 15.

Cal. hexaphyllus, persistens. *Cor.* nulla. *Cap-
sula* supera, trivalvis, uni- vel tri-locularis.
Semina plurima. *Stigmata* tria.

* *Culmis nudis.*

acutus. 1. JUNCUS culmo nudo tereti, paniculâ terminali, involucro diphylllo spinoso, capsulis subrotundis mucronatis.

Juncus acutus. *Sp. Pl.* 463. *Huds.* 148.
Willd. 346.

J. acutus capitulis Sorghi. *Baub. Pin.* 11.
Prodr. 21. *f.* 2. *Theatr.* 173. *Raii Syn.* 431.

J. pungens, sive acutus, capitulis Sorghi. *Mor-
ris. sect.* 8. *t.* 10. *f.* 15.

J. maritimus, Sorghi paniculâ utriculatâ. *Bar-
rel. Ic. t.* 203. *f.* 2.

Angl. Great Sharp Sea Rush.

In arenosis maritimis.

On the coast of Merionethshire plentifully. *Ray.*
At Brancaster, Norfolk. *Mr. Crowe.* At Hol-
ker, Lancashire. *Mr. Woodward.* At Instow,
Devonshire, and Braunston Burrows, plentifully.
Rev. Dr. Gossacnough.

Perennis. Fl. Julio.

Radix fibrosa, cæspitosa. Culmi tripedales, erecti, stricti, simplices, nudi, teretes, glabri, rigidi. Folia culmo simillima sed breviora, radicalia, apice mucronato-pungentia. Paniculâ terminalis, corymbosa, erecta, multiflora. Involucrum diphyllum; foliola basi dilatata, apice foliis conformia, mucronato-pungentia, divaricata, inæqualia, longitudine varia: involucella plurima, membra-

membranacea, acuta, brevia, pedunculis interstincta. Flores congesti. Calycis foliola ovata. Capsula triquetra-subrotunda, turgida, mucronata, dura, nitida, trilocularis. Semina plurima, ovata, dissepimentis affixa, fusca, pellucida, utrinque appendiculata.

2. *JUNCUS* culmo nudo tereti, paniculâ terminali prolifera, involuero diphylo spinoso erecto, capsulis oblongis. *maritimus.*

Juncus acutus β . *Sp. Pl.* 464. *Huds.* 148. *With.* 346.

J. acutus maritimus anglicus. *Raii Syn.* 431. *Morif. sect. 8. t. 10. f. 14.*

Angl. Lesser Sharp Sea Rush.

In palustribus maritimis copiosè, præcedentis socius. Salt marshes of Essex and Wales. *Ray.* Lancashire. *Rev. Mr. Wood.* About Burnham and Holkham, Norfolk. Sea coast east from St. Andrews, Scotland. *Mr. Mackay.*

Perennis. Fl. Augusto.

Præcedente humilior et longè tenuior, colore magis glauco. Panicula erecta, decomposita, quasi prolifera, ramulis valdè inæqualibus et subcymosis. Involucris foliolum exterius erectum, paniculam superans, mucronatopungens; interius brevissimum, subulatum. Calycis foliola lanceolata. Capsula prismatica, lineari-oblonga, nec subrotunda.

3. *JUNCUS* culmo nudo stricto glauco, paniculâ laterali erectâ, capsulis ellipticis acutis. *glaucus.*

Juncus glaucus. *Sibth.* 113. *Engl. Bot. t. 665.*

J. effusus β . *Huds.* 149.

J. inflexus. *Relb.* 141. *With.* 345. *Abbot.* 78. *Leers.* 87. *t. 13. f. 3.*

J. acutus. *Raii Syn.* 432. *Ger. em.* 35.

J. acutus vulgaris. *Morif. sect. 8. t. 10. f. 13.* *Lob. Ic.* 85.

Angl. Hard Rush.

In pascuis et ad vias, solo humidior.

Perennis. Fl. Julio.

Culmi sesquipediales, tenacissimi, striati, glauci; apice attenuati, acuti; basi stipulis fuscis, nitidis, vestiti. Folia culmis simillima, parum breviora. Panicula lateralis, decomposita, erecta. Calycis foliola subulata, striata. Capsula elliptica, acuta, mucronulata, fusca, nitida.

conglomeratus.

4. **JUNCUS** culmo nudo striato, paniculâ laterali conglobatâ, capsulis retusis, floribus triandris.

Juncus conglomeratus. *Sp. Pl.* 464. *Huds.* 148.

With. 345. *Relb.* 140. *Sibth.* 113. *Leers.* 86. t. 13. f. 1.

J. lævis vulgaris, paniculâ compactiore. *Raii Syn.* 432.

J. lævis, paniculâ non sparsâ. *Morif. sect.* 8. t. 10. f. 7.

Angl. Common Rush.

In pascuis et ad vias, solo humido, vulgaris.

Perennis. Fl. Julio.

Culmi bipediales, minutè striati, acuti, nec pungentes. Panicula lateralis, multiflora, densissimè conglobata. Calycis foliola lanceolata, nervosa. Capsula obovata, apice retusa et ferè triloba. “Stamina constanter tria, nunquam sex;” *Leers.*

effusus.

5. **JUNCUS** culmo nudo striato, paniculâ laterali effusâ supradecompositâ, capsulis obtusis.

Juncus effusus. *Sp. Pl.* 464. *Huds.* 148. *With.*

345. *Relb.* 141. *Sibth.* 113. *Leers.* 87. t. 13. f. 2.

J. lævis vulgaris, paniculâ sparsâ nostras. *Raii Syn.* 432. *Lob. Ic.* 84.

J. lævis, paniculâ sparsâ, major. *Morif. sect.* 8. t. 10. f. 4.

J. lævis.

J. lævis. *Ger. em.* 35.

β. *J. lævis alter.* *Morif.* v. 3. 231. n. 5. *Herb.*
Bobart.

Angl. Soft Rush.

In pascuis humidis et ad vias passim.

Perennis. Fl. Julio.

Culmi pallidiores, molles, fragiles. Panicula lateralis, effusa, divaricata, ramosissima, multiflora. Calycis foliola angustè lanceolata, acuminata, striata. Capsula obovata, obtusa. Panicula in β minùs effusa est.

6. **JUNCUS** culmo nudo filiformi nutante, pani *filiformi* culâ laterali bracteâtâ subsimplici, capsulis *mis.* subrotundis.

Juncus filiformis. *Sp. Pl.* 465. *Huds.* 149.

With. 346. *Sm. Spicileg.* t. 3. *Dicks. Dr.*

Pl. 32. *H. Sicc. fasc.* 14. 16. *Leers.* 87.

t. 13. f. 4.

J. parvus, calamo supra paniculam compactam longiùs producto. *Raii Syn.* 432. *Pluk.*

Phyt. t. 40. f. 8.

Angl. Least Rush.

In pratis turfosis irriguis alpinis.

Near Ambleside, Westmoreland. *Newton.* At

Windermoor in Cartmel, Lancashire, Mr. Jack-

son. *With.* Near Derwentwater, Cumberland,

and on Ben Lawers in Scotland. *Mr. Dickson.*

Perennis. Fl. Augusto.

Habitus præcedentium, sed minor. Radix valdè repens.

Culmus pedalis, apice incurvus, acutus, longè supra paniculam extensus, basi squamis fuscis, obtulis, aristatis, striatis, vaginatus. Panicula parva, quinque- ad octo- flora, pedunculis vix ramosis, basi foliolo setaceo, longitudine vario, suffulta. Calycis foliola lanceolata, carinâ trinervi. Capsula subrotunda, obtusa.

trifidus. 7. **JUNCUS** culmo nudo, foliis floribusque tribus terminalibus.

Juncus trifidus. *Sp. Pl.* 465. *Huds.* 149.
With. 345. *Lightf.* 183. *t.* 9. *f.* 1. *Fl. Dan.*
t. 107. *Dicks.* *H. Sicc. fasc.* 1. 6. *Baub.*
Hist. v. 2. 521.

J. monanthos. *Jacq. Enum.* 61. 235. *t.* 4. *f.* 1.

J. acumine reflexo, trifidus. *Baub. Prod.* 22.
Theatr. 185.

Angl. Three-leaved Rush.

In paludosis alpinis Scotiæ.

Perennis. Fl. Julio.

Culmi spithamæi, stricti, filiformes, graciles, nudi, striati, basi squamis nitidis vaginati, apice foliis tribus, alternis, subulatis, incurvato-erectis, terminati. Flores plerumque tres, terminales, erecti, sessiles, congesti, ferruginei, nitidi. Calycis foliola ovata, acuta. Capsula elliptica, acuminata.

squarrosus. 8. **Juncus** culmo nudo, foliis setaceis, paniculâ terminali compositâ glomeratâ.

Juncus squarrosus. *Sp. Pl.* 465. *Huds.* 149.
With. 346. *Relb.* 142. *Sibth.* 114. *Dicks.*
H. Sicc. fasc. 13. 12. *Fl. Dan.* *t.* 430.

J. montanus palustris. *Raii Syn.* 432.

Gramen junceum maritimum. *Ger. em.* 21.

G. junceum palustre humilius, foliis et spicâ

Junci. *Morif. sect.* 8. *t.* 9. *f.* 13.

Angl. Moss Rush. Goose Corn.

In ericetis sterilibus fabulosis frequens.

Perennis. Fl. Junio, Julio.

Radix fibrosa, cæspitosa. Culmi pedales, stricti, rigidi, teretes, striati, nudi. Folia radicalia plurima, patentia, incurva, setacea, canaliculata, glabra. Panicula terminalis, erecta, ramosa, basi foliolo setacco, longitudine

dine vario, vaginata. Calyx fuscus, margine scarioso, albo. Capsula obovata, obtusiuscula.

* * *Culmis foliosis.*

9. JUNCUS foliis nodoso-articulatis, capitulis pa-
niculatis multifloris. *articulatus.*

Juncus articulatus. *Sp. Pl.* 465. *Huds.* 149.
With. 347. *Engl. Bot. t.* 238. *Leers.* 88.
t. 13. *f.* 6.

J. compressus. *Relb.* 142. *Sibth.* 114. *Abbot.* 79.

J. foliis articulosis, floribus umbellatis. *Raii Syn.* 433.

Gramen junceum articulatum palustre humilius, utriculis frequenter donatum. *Morif. sect.* 8.
t. 9. *f.* 2.

β. *Juncus articulatus.* *Relb.* 143.

J. nemorosus. *Sibth.* 114.

J. nemorosus, folio articulofo. *Raii Syn.* 433.

Gramen junceum articulatum palustre erectum et clatius. *Morif, sect.* 8. *t.* 9. *f.* 1.

G. junceum sylvaticum. *Ger. em.* 22.

Angl. Jointed Rush.

In pratis et pascuis humidis, aquosis et umbrosis.

Perennis. *Fl.* Junio.

Radix fibrosa, caespitosa, subrepens. Culmus erectiusculus, foliosus, teretiusculus, glaber, basi squamis vaginatus. Folia alterna, vaginantia, acuta, subcylindrica, compressiuscula, fistulosa, intus septis transversis numerosis concamerata. Panicula terminalis, ramosa, ramis compositis, laevibus. Involucrum monophyllum, foliaceum, ad basin paniculae; involucella alterna, membranacea, acuta. Florum fasciculi sessiles, laterales et terminales, subquinqueflori. Calycis foliola lanceolata, acuminata, spadiceo-purpurea, dorso subinde viridia. Capsula oblonga, acutè triquetra, mucronata.
β differt culmo erectiore, paniculâ ramosiori, floribus minoribus,

noribus, pallidioribus et obtusioribus, foliisque vix compressis.

uliginosus. 10. JUNCUS foliis setaceis subnodoso-articulatis, capitulis trifloris subproliferis, culmo bulboso radicante.

Juncus uliginosus. *Sibth.* 115. *With.* 348.

J. viviparus. *Relb.* 143. *Abbot.* 79.

J. subverticillatus. *Jacq. Coll.* v. 3. 52.

J. articulatus γ. *Huds.* 150. *With.* 347.

J. articulatus arenosus. *Bute.* v. 7. 206.

J. n. 1320. *Hall. Hist.* v. 2. 170.

Junci bulbosi varietas. *Fl. Dan.* t. 817.

Gramen junceum capsulis triangulis minimum.

Raii Syn. 434. *Morif. sect.* 8. t. 9. f. 3.

β. *G. junceum* minimum, paniculis foliaceis. *Marrif. sect.* 8. t. 9. f. 4.

G. junceum aquaticum, paniculis cum foliis capillaribus. *Pluk. Phyt.* t. 32. f. 3.

Juncoides calyculis paleaceis, glomeratis, folio varians. *Scheuchz. Agrost.* 330. t. 7. f. 10.

Angl. Little bulbous Rush.

In ericetis humidis arenosis, vel turfosis.

Perennis. Fl. Junio, Julio.

Radix fibrosa. Culmi filiformes, decumbentes, foliosi, sæpe geniculati, apice ramosi, basi bulbosi, geniculis radicanlibus et sæpiùs bulbosis. Folia conferta, setacea, glabra, intus minùs evidenter ac regulariter conca-merata. Capitula plerumque triflora, lateralialia et terminalialia, subinde inter culmi ramulos subverticillatos axillaria. Bractææ scariosæ, parvæ. Calycis foliola lanceolata, acuminata, trinervia, acutè carinata, dorso viridia, margine scariosa. Capsula acutè triquetra, obtusa.

Germen sæpè gemmiparum, viviparum.

11. JUNCUS foliis linearibus canaliculatis, culmo *bulbosus*,
basi folioso, paniculâ cymosâ, capsulis obtusis.

Juncus bulbosus. *Sp. Pl.* 466. *Huds.* 150.

With. 348. *Relb.* 143. *Sibth.* 115. *Ab-*
bot. 79. *Dicks. H. Sicc. fasc.* 13. 14. *Leers.* 89.
t. 13. *f.* 7. *Fl. Dan. t.* 431.

J. parvus cum pericarpis rotundis. *Raii Syn.* 433.
Baub. Hist. v. 2. 522.

Gramen junceum pericarpis rotundis, vel vul-
gare. *Morif. sect.* 8. *t.* 9. *f.* 11.

Angl. Round-fruited Rush.

In pratis et pascuis humidis, præcipuè falsis.

By the Long Walk, Windsor, plentifully. *Rev.*

Dr. Goodenough.

Perennis. *Fl.* Julio.

Radix repens, fibrosa, nequaquam bulbosa. Culmi erecti,
altitudine varii, simplices, teretiusculi, parùm compressi,
glabri, basi præcipuè foliosi. Folia patentia, linearia,
canaliculata, acuta. Panicula terminalis, erecta, cy-
mosa, multiflora, basi foliata. Foliola calycina obtu-
siuscula, striata, viridia, margine scariofo fusco. Cap-
sula calyce longior, obovata, obtusa.

Nominis trivialis ratio me planè fugit.

12. JUNCUS foliis linearibus canaliculatis, cul- *busonius*,
mo dichotomo racemoso, floribus solitariis.

Juncus busonius. *Sp. Pl.* 466. *Huds.* 150.

With. 348. *Relb.* 144. *Sibth.* 115. *Dicks.*

H. Sicc. fasc. 4. 6. *Leers.* 89. *t.* 13. *f.* 8.

Rose Bot. 452. *t.* 2. *f.* 5, 6.

J. palustris humilior erectus. *Raii Syn.* 434.

Gramen junceum. *Ger. em.* 4.

G. junceum vulgare, calyculis paleaceis. *Morif.*
sect. 8. *t.* 9. *f.* 14.

β. *G. juncoides* minimum Anglo-Britannicum.

Dill. in Raii Syn. 434.

Angl. Toad Rush.

In aquosis et palustribus.

Annua. Fl. Julio, Augusto.

Radix fibrosa. Culmi plurimi, cæspitosi, patentes, teretes, foliosi, dichotomi, apice racemosi. Folia linearia, acuta, lævia, supernè canaliculata, basi dilatata. Flores solitarii, subseffiles, erecti. Calycis foliola lanceolata, acuta, margine scariofo albo. Capsula lineari-elliptica, calyce brevior, obtusiuscula.

*biglu-
mis.*

13. JUNCUS foliis planis, capitulo bifloro terminali secundo basi foliato.

Juncus biglumis. *Sp. Pl.* 467. *Montin. in Amæn. Acad. v. 2. 266. t. 3. f. 3. Hudf.* 649. *With.* 348. *Lightf.* 1100. *Fl. Dan. t. 120. Angl.* Two-flowered Rush.

In locis irriguis alpium Scoticarum rariùs.

On Ben Lomond. *Rev. Mr. Stuart.* Ben Lawers, and other highland mountains. *Mr. Mackay.*

Perennis. Fl. Augusto.

Radix fibrosa. Culmus subtriuncialis, erectus, simplicissimus, teres, striatus, basi apiceque foliosus. Folia subulata, complanata, erecta. Flores terminales, unus super alterum, secundi, subseffiles, folio caulino superati, subtus bractea obtusâ suffulti. Calycis foliola lanceolata, carinata. Capsula obovata, obtusa, calyce longior.

*triglu-
mis.*

14. JUNCUS foliis planis, capitulo trifloro terminali erecto aphylo bracteato.

Juncus triglumis. *Sp. Pl.* 467. *Hudf.* 151. *With.* 349. *Lightf.* 186. *t. 9. f. 2. Fl. Dan. t. 132. Fl. Lapp. t. 10. f. 5. Dickf. H. Sicc. fasc. 2. 3. Rel. Rudb. 24. f. 2.*

Juncello accedens graminifolia planta capitulis Armeriæ proliferæ. *Raii Syn.* 430.

Gramen

Gramen cyperoides minus, Caryophylli proliferi capitulis. *Morif. sect. 8. t. 12. f. 40.*

Angl. Three-flowered Rush

In rivulis alpinis.

About the tops of the Scottish and Welch mountains. At Burrowdale, near Keswick, Cumberland, Mr. Robson. *With.* On Ben Lomond, very plentifully.

Perennis. Fl. Julio.

Radix fibrosa. Culmus spithamæus, erectus, simplicissimus, teres, striatus, basi tantum foliosus. Folia lineari-lanceolata, acuta, complanata, erectiuscula, basi vaginantia. Capitulum terminale, erectum, triflorum, subinde bi- vel quadri-florum. Bractææ ovatæ, concavæ, capitulo breviores, quarum duæ exteriores maximæ. Calycis foliola elliptico-lanceolata, obtusa, subcarinata. Capsula elliptica, obtusiuscula, calyce longior.—Distinctissima a præcedente.

15. JUNCUS foliis planis amplexicaulibus, capitulo terminali subgemino multifloro basi foliato, *castaneus.* bracteis acutis.

Juncus Jacquini. *Sym. Syn. 87. Hull. 76.*

Angl. Clustered Alpine Rush.

In paludosis alpinis Scotiæ, solo micaceo.

On Ben Lawers. *Mr. Dickson and Mr. Mackay.*

Perennis. Fl. Julio.

Radix laxè repens, farmentosa. Culmi erecti, spithamæi, subinde pedales, teretes, foliosi. Folia omnia caulina, alterna, erecta, plana, linearia, acuta; basi dilatata, amplexicaulia, vaginantia; supremum capitulis subjectum, reliquis brevius. Capitula terminalia, erecta, bina, unum super alterum, aliquando solitaria, atrofusca, nitida, sæpius quadriflora, subinde sex- vel octoflora, rariùs triflora. Bractææ solitariae sub singulo flore, a calyce parum remotæ, lineari-lanceolatæ, acutæ, fuscæ. Calycis foliola lanceolata, acuta, dorso trinervia.

Stamina

Stamina sex, capillaria, alba, longitudine calycis. Antheræ terminales, lineares, ochroleucae. Stigmata linearia, plana. Capsula elliptico-oblonga, obtusa, apice nigra, mucronulata, calyce duplo longior, trilocularis. Semina utrinque appendiculata.

- J. Jacquini, huic facie quodammodo similis, gaudet radice oblongâ, simpliciter repente; culmis congestis, stric-
tissimis, monophyllis, basi vaginatis, nec foliosis; brac-
teis calyci arctè approximatis, geminis, subrotundis, ob-
tusis; staminibus calyce triplo brevioribus.

pilosus. 16. JUNCUS foliis planis pilosis, paniculâ cymosâ
divaricatâ, floribus solitariis.

Juncus pilosus. *Sp. Pl.* 468. *Huds.* 151.

With. 349. *Relb.* 145. *Sibth.* 115. *Curt.*

Lond. fasc. 5. *t.* 25. *Lightf.* 186. *Dicks.*

H. Sicc. fasc. 14. 15. *Leers.* 90. *t.* 13. *f.* 10.

Gramen nemorosum hirsutum vulgare. *Raii*
Syn. 416.

G. hirsutum nemorosum. *Ger. em.* 19.

G. hirsutum latifolium majus. *Morif. sect.* 8.

t. 9. *f.* 1.

Angl. Hairy Rush.

In nemorosis et dumosis vulgaris.

Perennis. Fl. Martio, Aprili.

Radix fibrosa, stolonifera. Culmi spithamæi, erecti, sim-
plices, graciles, teretes, foliosi. Folia lineari-lanceo-
lata, acuta, plana, nervosa, graminea, margine pilis al-
bis, mollibus, ciliata; radicalia maxima; caulina al-
terna, vaginantia. Panicula terminalis, cymosa, ra-
mosa, divaricata, ramis inferioribus demum refractis.
Flores solitarii, bracteis geminis, acutis, calyci approxi-
matis. Calycis foliola lanceolata, acuminata, nigra,
albo marginata. Stamina brevissima. Antheræ mag-
næ, tetragonæ, flavæ. Capsula ovata, obtusa, pallidè
flavesceñs.

17. JUNCUS foliis planis pilosis acuminatis, paniculâ cymosâ decompositâ, floribus fasciculatis. *sylvaticus.*

Juncus sylvaticus. *Huds.* 151. *Sibth.* 116. *Curt. Lond. fasc. 5. t. 26.* *Dicks. H. Sicc. fasc. 13. 13.* *Lightf.* 186.

J. maximus. *With.* 349.

J. pilosus δ . *Sp. Pl.* 468.

J. latifolius. *Jacq. Coll. v. 3: 59.*

Gramen nemorosum hirsutum latifolium maximum. *Raii Syn.* 416.

G. hirsutum latifolium majus; junceâ paniculâ. *Morif. sect. 8. t. 9. f. 2.*

G. hirsutum latifolium minus. *Ibid.*

G. Luzulæ maximum. *Baub. Hist. v. 2. 493.*
Angl. Wood-Rush.

In sylvis, nemorosis et dumetis frequentis.

Perennis. Fl. Maio.

Habitus præcedentis, sed major: Culmus ferè tripèdalis: Folia magis acuminata, latiora, pilis longioribus. Paniculâ decomposita, ramis vix refractis. Flores congesti, plerumque terni, minores tamen quam in Juncopiloso.

18. JUNCUS foliis planis pilosis, spicis terminatis, capulis obtusis. *campestris.*

Juncus campestris. *Sp. Pl.* 468. *Huds.* 152. *With.* 350. *Relb.* 145. *Sibth.* 116. *Curt. Lond. fasc. 2. t. 19.* *Dicks. H. Sicc. fasc. 14. 17.* *Leers.* 91. *t. 13. f. 5.*

Gramen exile hirsutum. *Raii Syn.* 416. *Ger. em.* 17.

G. hirsutum capitulis psyllii. *Baub. Prod.* 15. *Morif. sect. 8. t. 9. f. 4.*

G. hirsutum capitulo globofo. *Ibid?*

β . *G. hirsutum* elatius, paniculâ junceâ compactâ.

Raii Syn. 410. synonymis mihi dubiis.

Angl. Hairy Field Ruth.

In pascuis ficcioribus sterilioribus. β in paludibus turfosis.

Perennis. Fl. Aprili, Maio.

Radix repens. Culmi spithamæi, foliosi. Folia linearia, plana, acuta, margine pilosissima. Spicæ terminales, maximâ ex parte pedunculatæ, ovatæ, obtusæ, subdecemfloræ, erectæ, demùm nutantes. Flores sessiles, basi bracteis tribus vel quatuor laceris. Calycis foliola lanceolata, acuminata. Stamina brevia. Capsula obovata, obtusa, imperfectè trilocularis, trisperma.

Varietas β altior est, magisque strigosa, foliis longioribus, spicis arctè conglomeratis; in structurâ verò floris, fructûs, atque bractearum, ne minimam quidem inveni differentiam. Figuræ a Raio citatæ non placent, ob spicas pedunculatas, nequaquam congestas.

spicatus. 19. *JUNCUS* foliis planis, spicâ racemosâ nutante basi compositâ, capsulis acutis.

Juncus spicatus. *Sp. Pl.* 469. *Fl. Lapp. t.* 10. *f.* 4. *Hudsf.* 650. *With.* 350. *Lightf.* 187. *Fl. Dan. t.* 270. *Dicksf. Dr. Pl.* 33. *H. Sicc. fasc.* 15. 11.

Angl. Spiked Rush.

In alpibus Scoticis.

On the grassy summit of Ben Lomond.

Perennis. Fl. Julio.

Radix fibrosa, cæspitosa. Folia linearia, recurva, marginibus parùm pilosa. Culmus spithamæus, foliosus, gracilis. Spica terminalis, solitaria, nutans, ovato-oblonga, obtusa, multiflora, ferrugineo-nigra, basiflobata. Bractear ciliato-laceræ, pallidæ. Calycis foliola lanceolata, acuminata. Capsula acuta.

181. BERBERIS. *G. Pl.* 175. *Juss.* 286.

Gærtn. t. 42.

Cal. hexaphyllus. *Petala* 6; ad ungues glandulis duabus. *Stylus* nullus. *Bacca* supera, disperma.

1. BERBERIS floribus racemosis, spinis triplici-vulgaribus, foliorum serraturis aristatis.

Berberis vulgaris. *Sp. Pl.* 471. *Huds.* 137.

With. 350. *Relb.* 134. *Sibth.* 108. *Ab-*

bot. 80. *Engl. Bot. t.* 49. *Fl. Dan. t.* 904.

Woodv. Suppl. t. 234. *Sm. Tracts.* 165.

B. dumetorum. *Rai Syn.* 465.

Spina acida five *Oxyacantha.* *Ger. em.* 1325.

Angl. Barberry.

In sepibus et dumetis, apricive calcareis.

Frutex. *Fl.* Maio, Junio.

Rami viminei, angulati, pallidi, glabri. Folia ovata, glabra, spinuloso-serrata, fasciculata, petiolata; fasciculis spinâ trifidâ, divaricatâ, suffultis. Racemi c gemmis foliorum penduli, solitarii, simplices, multiflori. Flores flavi, glandulis antherisque fulvis. Baccæ oblongæ, miniatæ. Stamina basin versus, c parte inferiore, stimulis maximè obnoxia sunt.

182. FRANKENIA. *G. Pl.* 176. *Juss.* 303.

Cal. quinquefidus, infundibuliformis. *Petala* 5.

Stigmata 3. *Capsula* supera, unilocularis, trivalvis.

1. FRANKENIA foliis linearibus confertis basi lævis. ciliatis.

Frankenia lævis. *Sp. Pl.* 473. *Huds.* 137.

With. 352. *Relb.* 135. *Engl. Bot. t.* 205.

Franca maritima supina saxatilis glauca ericoides

sempervirens, flore purpureo. *Mich. Gen.* 23.
t. 22. f. 1.

Lychnis supina maritima Ericæ facie. *Raii Syn.* 338.

Polygonum serpillifolium. *Ger. em.* 566.

Sea Heath. *Pet. H. Brit.* t. 10. f. 11.

Angl. Smooth Sea-Heath.

In palustribus maritimis cœnosis.

On the coasts of Suffex, Essex and Kent in many places. In the salt marshes at Yarmouth across the river.—Isle of Sheppey abundantly. *Rev. Dr. Goodenough.*

Perennis. Fl. Julio.

Radix lignosa. Caules procumbentes, ramosi, dichotomi, teretes, foliosi, subpubescentes. Folia fasciculata, revoluta, basi ciliata. Flores e dichotomiâ caulis, subinde terminales, solitarii, sessiles, carnei. Petala erosa, unguiculata, cum squamâ oblongâ ungui adnatâ. Stigmata tria, oblonga, obtusa. Capsula polysperma.

*pulveru- 2. FRANKENIA foliis obovatis retusis subtus
lenta.* villoso-pulverulentibus.

Frankenia pulverulenta. *Sp. Pl.* 474. *Huds.* 138.
With. 352.

Franca maritima quadrifolia annua supina, Chamæsyces folio et facie, flore ex albo purpurascente. *Mich. Gen.* 23.

Alfine maritima supina, foliis Chamæfices. *Dill. in Raii Syn.* 352.

Anthyllis valentina. *Clus. Hist.* p. 2. 186. *Ger. em.* 566.

Valentia Knotgrass. *Pet. H. Brit.* t. 10. f. 8.
Angl. Powdery Sea-Heath.

In maritimis Suffexiæ, rariùs.

Found by Mr. Brewer on the coast of Suffex. *Dill.*
Between Bognor and Brighthelmston. *Huds.*

Annua.

Annua. Fl. Julio.

Radix fibrosa. Caules prostrati, e basi ramosi, pubescentes, teretes. Folia obovata, retusa, subtus villosa et subpulverulenta. Calyx prismaticus. Specimina Britanniae indigena nondum vidi.

183. PEPLIS. *G. Pl.* 176. *Juss.* 333.

Gærtn. t. 51.

Cal. campanulatus, ore duodecimfido. *Petala* 6, calyci inserta. *Capfula* supera, bilocularis.

I. PEPLIS floribus subapetalis, foliis obovatis. *Portula.*

Peplis Portula. *Sp. Pl.* 474. *Huds.* 147. *With.*

352. *Relb.* 139. *Sibth.* 108. *Abbot.* 78.

Curt. Lond. fasc. 4. *t.* 27. *Dicks. II. Sicc. fasc.* 4. 8. *Fl. Dan. t.* 64.

Portula. *Dill. Giff.* 120. *append.* 133. *t.* 7.

Raii Syn. 368.

Alfine rotundifolia, five Portulaca aquatica. *Ger. em.* 614.

Glaux aquatica, folio subrotundo. *Loesel. Pruss.* 106. *t.* 20.

G. palustris, flore striato clauso, foliis portulacæ.

Vaill. Paris. t. 15. *f.* 5.

Glaucoides palustre, Portulacæ folio, flore purpureo. *Mich. Gen.* 21. *t.* 18. *f.* 1.

Angl. Water Purslane.

In palustribus et ericetis humidis frequens.

Annua. Fl. Julio, Augusto,

Caules prostrati, subnatantes, radicanes, simplices, foliosi. Folia opposita, petiolata, obovata, obtusa, integerrima, glabra. Flores axillares, solitarii, sessiles, parvi, rubicundi. Petala fugacia, sæpè nulla. Capfula globosa, pellucida, polysperma.

II. TRIGYNIA.

184. RUMEX. *G. Pl.* 178. *Juss.* 82.

Gærtn. t. 119.

Cal. triphyllus. *Petala* 3, conniventia. *Semina* unicum, triquetrum, superum, nudum. *Stigmata* multifida.

* *Flores hermaphroditici.*

1. RUMEX valvulis integerrimis oblongis, unicâ præcipuè graniferâ, foliis cordato-lanceolatis.

Rumex sanguineus. *Sp. Pl.* 476. *Huds.* 153. *With.* 353. *Sibth.* 118.

Lapathum folio acuto rubente. *Raii Syn.* 142.

L. sanguineum. *Munt. Brit. t.* 113. *Baub. Hist.* v. 2. 988.

L. rubens. *Camer. Epit.* 229.

L. fativum sanguineum. *Ger. em.* 390.

Blood-veined Dock. *Pet. H. Brit. t.* 2. f. 5.

β. viridis. *Sibth.* 118.

Rumex acutus. *Curt. Lond. fasc.* 3. t. 21.

With. 354. *Wade Dubl.* 103. *Huds.* 155 β.

Lapathum viride. *Dill. in Raii Syn.* 141.

Bloodless Dock. *Pet. H. Brit. t.* 2. f. 6.

An l. Bloody-veined Dock. β Green-veined Dock.

In nemorosis et ad vias.

In woody places about Hampstead. *Merret.* Near Maidstone. *Huds.* Headington hill, Oxfordshire. *Sibth.* At Lowestoft, Suffolk, and about Bristol.

β. In shady places frequent.

Perennis. Fl. Julio.

Radix

Radix fusiformis. Caulis erectus, ramosus, angulatus, foliosus, glaber. Folia omnia petiolata, lanceolata, glabra, venosa, margine subcrispa; radicalia maxima, basi cordata. Racemi terminales, patentes, subaphylli, floribus alternatim fasciculatis, pedicellatis, nutantibus. Calyx integerrimus. Petala oblonga, integerrima, persistentia, valvulas semen amplectentes constituentia, quarum unica (quandoque omnes) demum granifera evadit. Semen parvum, acutè triquetrum.

Venæ et petioli foliorum in æ succo sanguineo scatent.

2. RUMEX valvulis ovatis integris omnibus *gracrispus*.
niferis, foliis lanceolatis undulatis acutis.

Rumex crispus. *Sp. Pl.* 476. *Huds.* 153.

Willd. 353. *Relb.* 146. *Sibth.* 117. *Abbot.* 81. *Curt. Lond. fasc.* 2. t. 20.

Lapathum folio acuto crispo. *Raii Syn.* 141.

L. longifolium crispum. *Munt. Brit. t.* 104.

Curled Sharp Dock. *Pet. II. Brit. t.* 2. f. 2.

Angl. Curled Dock.

In ruderatis, pascuis et ad vias vulgaris.

Perennis. Fl. Junio, Julio.

Radix fusiformis, flavescens. Caulis angulatus, sulcatus, læviusculus. Folia lanceolata, acuta, margine undulato-crispa; caulina ferè linearia et subseffilia. Racemi minùs patentes, densiores, basin versus subfoliolosi. Valvulæ fructûs ovatæ, subcordatæ, reticulato-venosæ, integræ, subrepandæ nec dentatæ, omnes graniferæ, grano maximo, ovato, gibbo, rubro. Semen præcedente majus.

3. RUMEX valvulis oblongis subdentatis omnibus *acutus*.
bus graniferis, foliis cordato-oblongis acumina-
tatis, racemis foliolosis.

Rumex acutus. *Sp. Pl.* 478. *Huds.* 155.

Relb. 147.

R. paludosus. *Willd.* 354, ex descr.

Lapathum acutum. *Raii Syn.* 142.

Smooth Sharp Dock. *Pet. H. Brit. t. 2. f. 3.*
 β. *Lapathum acutum minimum. Dill. in Raii*
Syn. 141. Baub. Hist. v. 3. 985.
 Small Sharp Dock. *Pet. H. Brit. t. 2. f. 4.*
Angl. Sharp Dock.

In incultis et aquosis passim. *Raius.*
 In marshes at Surlingham, below Norwich.

Perennis. Fl. Julio.

Caulis angulatus, sulcatus, glaber. Folia cordato-lanceolata, acuminata, petiolata, repanda. Racemi divaricato-adscendentes, elongati. Flores semiverticillati, numerosi, densi, nutantes; verticillis omnibus ferè foliolo petiolato, patente, suffultis. Valvulae lanceolato-oblongae, subinde dentatae, omnes granulo maximo, gibbo, rubro, onustae. Semen exiguum.

obtusifolius. 4. RUMEX valvulis dentatis unicâ præcipuè graniferâ, foliis radicalibus cordatis obtusis, caule scabriusculo.

Rumex obtusifolius. Sp. Pl. 478, Hudf. 155. With. 357. Relb. 148. Sibth. 118. Abbot. 81. Curt. Lond. fasc. 3. t. 22.

Lapathum vulgare, folio obtuso. Raii Syn. 141. L. sylvestre, folio minùs acuto. Ger. em. 388. Lob. Ic. 285.

Broad Dock. *Pet. H. Brit. t. 2. f. 9.*
Angl. Broad leaved Dock.

In ruderatis et ad vias passim.

Perennis. Fl. Julio, Augusto.

Radix multiceps, fusca, intùs flava. Caulis sulcatus, supernè præcipuè scaber. Folia petiolata; radicalia maxima, cordata, obtusa; caulina ovato-lanceolata, acutiuscula. Racemi ferè aphylli. Flores semiverticillati, pedunculis capillaribus. Valvulae magnae, cordatae, venosae, dentatae, dentibus subulatis. Granulum in valvulâ exteriorè præcipuè conspicuum, lineari-oblongum,

longum, ratione valvularum parvum. Semen quam in præcedentibus majus.

5. RUMEX valvulis dentatis unicâ præcipuè *gr. pulcher*, niferâ, foliis radicalibus panduriformibus, caule glabro divaricato.

Rumex pulcher. *Sp. Pl.* 477. *Huds.* 153. *With.* 356. *Relb.* 148. *Abbot.* 82.

Lapathum pulchrum bononiense sinuatum. *Raii Syn.* 142. *Bauh. Hist.* v. 2. 988.

L. ramosum procumbens, pericarpio spinoso, folio fidium instar. *Morif. sect.* 5. *t.* 27. *f.* 13. *Angl.* Fiddle Dock. *Pet. H. Brit.* t. 2. *f.* 10.

In pascuis solo glareoso.

Perennis. Fl. Augusto.

Caulis subprocumbens, ramis elongatis, refractis, angulatis, glabris. Folia radicalia obtusa, medio utrinque sinuata; caulina parva, oblonga, integra. Racemi foliolosi. Valvulæ deltoideæ, dentatæ, reticulato-venosæ, quarum unica magis evidenter granifera est.

6. RUMEX valvulis deltoideis fetaceo-dentatis *mariti-* graniferis, foliis linearibus, verticillis confertis. *mus.*

Rumex maritimus. *Sp. Pl.* 478. *Huds.* 155.

R. aureus. *With.* 356. *Abbot* 81.

Lapathum folio acuto, flore aurco. *Raii Syn.* 142.

L. anthoxanthum. *Bauh. Hist.* v. 2. 987.

Bur Gold Dock. *Pet. H. Brit.* t. 2. *f.* 8.

Angl. Golden Dock.

In palustribus, præcipuè maritimis.

Perennis. Fl. Julio, Augusto.

Caulis angulatus, sulcatus, subflexuosus, scabriusculus, rubicundus. Folia linearia, obtusiuscula, plana. Racemi foliolosi. Verticilli arctè conferti, aurei. Valvulæ deltoideæ, omnes granulo oblongo notatæ, dentibus

bus fetaceis utrinque circiter quatuor, valvulâ longioribus. Semen exiguum.

palustris. 7. RUMEX valvulis lanceolatis graniferis basi dentatis, foliis lineari-lanceolatis, verticillis distantibus.

Rumex maritimus. *Huds.* 155 β. *With.* 356.
Curt. Lond. fasc. 3. t. 23.

Lapathum aureum. *Dill. in Raii Syn.* 142.

L. aquaticum, luteolæ folio. *Bocc. Mus. t.* 104.

Hydrolapathum minus. *Lob. Ic.* 286. *Ger. em.* 389.

Gold Dock. *Pet. H. Brit. t.* 2. f. 7.

Angl. Yellow Marsh Dock.

In paludosis, fossis, et ruderatis humidis.

About London in several places. *Dill. Curt.* By
Acle dam, Norfolk. *Mr. Pitchford.*

Perennis. Fl. Julio, Augusto.

Radix intus rubra. *Curt.* Caulis sulcatus, scaber, rubicundus. Folia lineari-lanceolata, acuta; radicalia maxima. Verticilli undique sparsi, foliolosi. Valvulæ omnes granulò oblongo notatæ, lanceolatæ, basi utrinque tridentatæ, dentibus fetaceis, multo brevioribus quam in præcedente.

aquaticus. 8. RUMEX valvulis ovatis integris obsoletè graniferis, foliis cordato-lanceolatis acutis.

Rumex aquaticus. *Sp. Pl.* 479. *Fl. Suec.* 117.
Lightf. 190.

R. Hydrolapathum. *Huds.* 154. *With.* 355.
Relb. 149. *Sibth.* 118. *Abbot.* 82. *Woodw. Med. Bot. t.* 178.

R. Britannica. *Huds. ed.* 1. 135.

Lapathum n. 1588. *Hall. Hist. v.* 2. 271.

L. maximum aquaticum five Hydrolapathum.
Raii Syn. 140. *Baub. Hist. v.* 2. 986.

Hydro.

Hydrolapathum magnum. *Lob. Ic.* 285. *Ger. em.* 389.

Hippolapathum. *Camer. Epit.* 232.

Britannica antiquorum vera. *Munt. Brit. t.* 1.

Angl. Great Water Dock. *Pet. H. Brit. t.* 2. f. 1.

In fossis et ad ripas fluviorum.

Perennis. Fl. Julio, Augusto.

Radix crassa. Caulis quinquepedalis, rectus, fulcatus, glaber. Folia subglaucæ, lanceolata, acuta; inferiora basi cordata. Verticilli approximati. Pedicelli capillares, apice incrassati, nutantes. Valvulæ magnæ, ovatæ, venosæ, integerrimæ, rariùs subdentatæ, omnes granulo parvo, lineari, sæpe obsoleto, notatæ. Semen magnum.

R. paludosus Hudsoni, a nemine adhuc detectus, hujusce varietas esse potest, cum granula in hæc et aliis speciebus magnitudine quandoque variant.

9. *RUMEX* valvulis ovatis integris grano destitu- *digynus*,
tis, floribus digynis.

Rumex digynus. *Sp. Pl.* 480. *Huds.* 156.

With. 357. *Lightf.* 190. *Fl. Dan. t.* 14.

Acetosa rotundifolia repens Eboracensis, folio in medio deliquium patiente. *Raii Syn.* 143.

Morif. sect. 5. t. 36. f. penult. *Pluk. Phyt.* t. 252. f. 2.

Welch Sorrel. *Pet. H. Brit. t.* 3. f. 4.

Angl. Mountain Sorrel.

Ad scaturigines alpinas.

On the mountains of Wales, Scotland, and the north of England, in very elevated spots, plentiful.

Perennis. Fl. Junio.

Folia omnia ferè radicalia, longiùs petiolata, reniformia, emarginata, repanda, venosa, pallidè viridia. Caulis nudius-

nudiufculus, spithamæus, paniculatus. Pedicelli fasciculati, capillares. Valvulæ ovatæ, emarginatæ, integerrimæ, rubicundæ, granulorum rudimentis planè destitutæ. Styli bini, distantes. Stigmata penicilli-formia.

* * *Flores diclini.*

Acetosa. 10. RUMEX floribus dioicis, foliis oblongis sagittatis, valvulis graniferis.

Rumex Acetosa. *Sp. Pl.* 481. *Huds.* 156.

With. 357. *Relb.* 149. *Sibth.* 119. *Abbot.* 82. *Engl. Bot. t.* 127. *Woodv. Med. Bot. t.* 69.

Lapathum acetosum vulgare. *Raii Syn.* 143.

Oxalis five Acetosa. *Ger. em.* 396.

Acetosa. *Brunf. Herb. v.* 2. 68.

β. Acetosa montana maxima. *Dill. in Raii Syn.* 143.

Acetosa pratensis, *folium.* *Pharmac. Lond.*

Angl. Common Sorrel.

In pratis et pascuis vulgaris.

Perennis. Fl. Junio.

Caulis teres, striatus, foliosus. Folia ovato-oblonga, sagittata; radicalia petiolata, obtusa; caulina sessilia, amplexicaulia, acuta, subrevoluta. Panicula ramosa. Flores semiverticillati, dioici. Valvulæ ovatæ, integræ, graniferæ,

Acetofella. 11. RUMEX floribus dioicis, foliis lanceolato-hastatis, valvulis grano destitutis.

Rumex Acetofella. *Sp. Pl.* 481. *Huds.* 156.

With. 358. *Relb.* 149. *Sibth.* 119. *Abbot.* 82. *Curt. Lond. fasc.* 5. *t.* 29.

Lapathum acetosum repens lanceolatum. *Raii Syn.* 143.

Oxalis tenuifolia. *Ger. em.* 397.

Angl. Sheep's Sorrel.

In pascuis sterilioribus glareosis vulgaris.

Perennis. Fl. Junio, Julio.

Præcedente duplò minor. Radix repens. Folia lanceolata; radicalia hastata; caulina sæpe integra. Racemi paniculati. Flores semiverticillati, dioici. Valvulæ ovatæ, integræ, granis destitutæ.

185. TOFIELDIA. *Huds.* 157.

Anthericum. *G. Pl.* 167.

Narthecium. *Juss.* 47. *Gerard.* *Villars.*

Cal. trifidus. *Petala* 6. *Styli* verticales, breves.

Capfulæ superæ, tres, basi junctæ, polyspermæ.

Antheræ subrotundæ.

1. TOFIELDIA petalis obovatis obtusis.

palustris.

Tofieldia palustris. *Huds.* 157. *With.* 358.

Sym. Syn. 89. *Engl. Bot. t.* 536.

Anthericum calyculatum. *Sp. Pl.* 447. *Fl.*

Lapp. t. 10. *f.* 3. *Lightf.* 181. *t.* 8. *f.* 2.

Dicks. H. Sicc. fusc. 13. 11. *Fl. Dan. t.* 36.

A. n. 1205. *Hall. Hist. v.* 2. 98. excluso syn.

Gorteri et Mœringii.

Narthecium. *Ger. Gallopr.* 142.

N. iridifolium. *Villars. Dauph. v.* 2. 225.

N. calyculatum. *Allion. Ped. v.* 2. 165. *La-*

marck Fl. Franc. v. 3. 298.

Scheuchzeria Pseudo-Asphodelus. *Scop. Carn.*

v. 1. 263. *Villars. Dauph. v.* 1. 322.

Phalangium Scoticum palustre minimum Iridis

folio. *Raii Syn.* 375.

P. alpinum palustre, Iridis folio. *Seguier. Veron.*

v. 2. 61. *t.* 14.

Asphodelus Lancastræ verus. *Ger. em.* 96.

Pseudo-asphodelus secundus. *Chus. Hist. p.* 1.

198.

Angl. Scottish Asphodel.

In turfosis irriguis alpinis.

Common in the highlands of Scotland.

Perennis. Fl. Augusto.

Radix lignosa, fibris longis, flexuosis, albis. Folia ensiformia, nervosa. Caulis spithamæus, erectus, simplex, nudiusculus, teres. Spica terminalis, solitaria, erecta, obtusa, densa. Flores albo-virides, subpedunculati. Bractææ solitariae, ovatae, concavae, parvae, ad basin pedunculorum. Calyx parvus, trifidus. Petala linearibovata, obtusa, concava, æqualia, persistentia. Stamina subulata, glabra, petalis opposita, at longitudine vix æqualia. Antherae incumbentes, cordato-subrotundæ, flavæ. Germen superum, tricoccum. Styli verticales, breves, distantes. Stigmata obtusa. Capsulæ tres, basi junctæ, gibbæ, vix inflatæ. Semina numerosa, elliptica, fulva.

Alia species ex Americâ boreali, *Gron. Virg.* 51, petalis lanceolatis acutis discrepat.

Scheuchzeria Linnæi differt a Tosfieldiâ calycis defectu, antheris linearibus, stigmatibus sessilibus, lateralibus, capsulis distinctis, inflatis, subdispermis.

186. TRIGLOCHIN. *G. Pl.* 179. *Juss.* 47.
Gærtn. t. 84.

Cal. triphyllus. *Petala* tria, calyciformia. *Styli* nulli. *Capsulæ* superæ, basi dehiscente. *Semina* solitaria.

palustre. 1. TRIGLOCHIN capsulis trilocularibus sublinearibus.

Triglochin palustre. *Sp. Pl.* 482. *Huds.* 152. *With.* 359. *Relb.* 146. *Sibth.* 119. *Abbot.* 83. *Engl. Bot. t.* 366. *Fl. Dan. t.* 490. *Leers.* 92. *t.* 12. *f.* 5. *Dicksf. H. Sicc. fasc.* 15. 10.

Juncago palustris et vulgaris. *Raii Syn.* 435.

Gramen aquaticum spicatum. *Ger. em.* 13.

G. enode

G. enode spicatum, five triglochin. *Morif.*

sect. 8. t. 2. f. 18.

Angl. Marsh Arrow-grafs.

In palustribus frequens.

Perennis. Fl. Julio.

Radix fibrosa. Folia radicalia, vaginantia, bifaria, linearia, canaliculata. Scapus pedalis, angulato-teres, simplex. Spica terminalis, multiflora. Capsula linearis, obtusa, trivalvis. Semina linearia.

2. TRIGLOCHIN capsulis sexlocularibus ovatis. *mariti-*

Triglochin maritimum. *Sp. Pl.* 483. *Huds.* 152. *num.*

With. 359. *Engl. Bot. t. 255. Fl. Dan. t. 306.*

Gramen marinum spicatum. *Raii Syn.* 435.

Lob. Ic. 16.

G. spicatum alterum. *Ger. em.* 20.

G. enode spicatum, pericarpiis parvis rotundis.

Morif. sect. 8. t. 2. f. 19.

Angl. Sea Arrow-grafs.

In maritimis cœnosis.

Perennis. Fl. Maio—Augusto.

Habitus prioris, sed folia latiora, magisque carnosæ. Capsula ovata, brevior, sexvalvis.

187. COLCHICUM. *G. Pl.* 180. *Juss.* 47.

Gærtn. t. 18.

Cal. spatha. *Cor.* sexpartita, tubo radicali. *Capsulæ* tres, superæ, connexæ, inflatæ, polyspermæ.

1. COLCHICUM foliis planis lanceolatis erectis. *autum-*

Colchicum autumnale. *Sp. Pl.* 485. *Huds.* 157. *nale.*

With. 359. *Sibth.* 119. *Abbot.* 83. *Engl.*

Bot. t. 133. Mart. Fl. Rust. t. 60. Woodv.

Med. Bot. t. 177.

C. com-

C. commune. *Raii Syn.* 373.

C. anglicum purpureum, et album. *Ger. ent.* 157.

Colchicum, radix recens. *Pharmac. Lond.*
Angl. Meadow Saffron.

In pratis solo pinguiore rariùs.

Chiefly in the north and west of England. *Ray.*
At Little Stonham, Suffolk, plentifully. *Mrs.*
Cobbold. Near Bury. *Mr. Mathew.* At Filkins
and Bradwell, Oxfordshire. *Rev. Dr. Good-*
enough.

Perennis. Fl. Septembri.

Bulbus ovatus, magnus. Folia lata, planiuscula, obtusa,
pedalia, vernalia. Flores radicales, aphylli, purpurei,
tubo longo, albo. Capsula elliptica, trifurca. Semina
globosa.

III. POLYGYNIA.

188. ALISMA. *G. Pl.* 181. *Juss.* 46.
Gærtn. t. 84.

Cal. triphyllus. *Petala* tria. *Capsulæ* plures,
aggregatæ, submonospermæ.

Plantago. 1. ALISMA foliis ovatis acutis, capsulis obtusè
trigonis.

Alisma Plantago. *Sp. Pl.* 486. *Huds.* 158.
With. 361. *Relb.* 150. *Sibth.* 120. *Ab-*
bot. 83. *Curt. Lond. fasc.* 5. t. 27. *Fl. Dan.*
t. 561.

Plantago

- Plantago aquatica.* *Raii Syn.* 257. *Camer. Epit.* 264. *Fuchf. Hist.* 42.
P. aquatica major. *Ger. em.* 417.
 Broad Water Plantain. *Pet. H. Brit. t.* 43. *f.* 6.
 β. *Alisma lanceolata.* *With.* 362. *Sym. Syn.* 90.
Plantago aquatica longifolia. *Dill. in Raii Syn.* 257.
 Narrow Water Plantain. *Pet. H. Brit. t.* 43. *f.* 7.
 γ. *Plantago aquatica leptomacrophyllos.* *Dill. in Raii Syn.* 258. *Loes. Pruff.* 199. *t.* 62.
Angl. Greater Water Plantain.

In aquosis et ad ripas fluviorum.

Perennis. Fl. Julio.

Radix fibrosa. Folia radicalia, petiolata, ovata, acuta, latitudine varia, nervosa, integerrima. Scapus trigonus, erectus, glaber. Panicula verticillato-decomposita, bracteolata. Flores pedicellati, erecti, dilutè purpurei. Fructus aggregatus, depressus; capsulis trigonis, obtusis.

Herba minax vi venenatâ ranunculorum, quibus ordine naturali affinis est.

2. *ALISMA* foliis cordato oblongis, floribus hexagynis, capsulis subulatis. *Damasconium.*

Alisma Damafonium. *Sp. Pl.* 486. *Huds.* 159. *With.* 361. *Curt. Lond. fasc.* 5. *t.* 28.

Damafonium stellatum Dalechampii. *Raii Syn.* 372.

Plantago aquatica minor stellata. *Ger. em.* 417.

P. aquatica minor altera. *Lob. Ic.* 301.

Star Water Plantain. *Pet. H. Brit. t.* 43. *f.* 5.

Angl. Star-headed Water Plantain.

In fossis et stagnis, solo glareoso, rariùs.

About London in several places. *Ray. Curtis.* On
 VOL. I. D d Hounslow

Hounslow heath, and on Winkfield plain near Windfor. *Rev. Dr. Goodenough.*

Perennis. Fl. Junio, Julio.

Præcedente longè humilior. Folia cordato-oblonga, obtusa. Scapus teres, verticillato-spicatus. Flores albi. Styli sex. Capsulæ sex, compressæ, mucronatæ, stellato-patentes, subinde dispermæ.

natans. 3. *ALISMA* foliis ellipticis obtusis, pedunculis solitariis, capsulis striatis.

Alisma natans. *Sp. Pl.* 487. *Huds.* 158. *With.* 362.

Damasonium repens, *Potamogetonis rotundifolii* folio. *Vaill. Act. Paris. Ann.* 1719. t. 4. f. 8. ex editione germanicâ.

Angl. Floating Water Plantain.

In lacubus alpinis rariss.

In the great lake below the old castle at Llanberris, North Wales. *Mr. Brewer.* In the south end of Bala lake, Merionethshire. *Rev. Mr. Wood.*

Perennis. Fl. Julio, Augusto.

Radix fibrosa, farmentosa. Herba natans. Caulis longitudine varius, ramosus, teres, lævis, foliosus. Folia alterna, petiolata, elliptica, obtusa, natantia; inferiora submersa, linearia, acuta, sessilia. Pedunculi oppositifolii, sæpe etiam terminales, uniflori, basi bracteâ vaginati. Flores albi. Styli plures. Capsulæ in circum digestæ, ellipticæ, compressæ, striatæ.

ranunculoides. 4. *ALISMA* foliis lineari-lanceolatis, capsulis pentagonis incurvis globoso-aggregatis.

Alisma ranunculoides. *Sp. Pl.* 487. *Huds.* 158.

With. 362. *Relb.* 150. *Sibth.* 120. *Abbot.* 84. *Engl. Bot. t.* 326. *Fl. Dan. t.* 122.

Plantago aquatica minor. *Raii Syn.* 257.

P. aquatica humilis. *Ger. em.* 417.

Angl. Small Water Plantain. *Pet. H. Brit.*
t. 43. *f.* 8.

In aquosis, et paludibus turfosis.

Perennis. Fl. Augusto.

Folia lineari-lanceolata. Scapus teres, floribus verticillatis, pallidè purpureis. Fructus globosus, squarrosus, e capsulis numerosissimis, obovatis, compressis, inæqualiter pentagonis, acutis, undique exstantibus.

Classis VII.

HEPTANDRIA.

Stamina 7.

MONOGYNIA.

Stylus 1.

189. TRIENTALIS. *Cal.* heptaphyllus. *Cor.*
septempartita, æqualis, plana. *Bacca* uni-
locularis, exsucca.

VII. HEPTANDRIA.

I. MONOGYNIA.

189. TRIENTALIS. *G. Pl.* 183. *Juss.* 96.
Gærtn. t. 50.

Cal. heptaphyllus. *Cor.* septempartita, æqualis,
plana. *Bacca* exsucca.

europæa. 1. TRIENTALIS.

Trientalis europæa. *Sp. Pl.* 488. *Huds.* 160.
With. 363. *Lightf.* 194. *Engl. Bot. t.* 15.
Fl. Dan. t. 84.

Alfinanthemos. *Raii Syn.* 286

Herba Trientalis. *Bauh. Hist. v.* 3. 536.

Pyrola alfinis flore major. *Bauh. Prodr.* 100.

P. alfinis flore europæa. *Morif. sect.* 12. *t.* 10.
f. 6.

Angl. Chickweed Winter-green.

In sylvis ad latera montium, ericisque turfosis.

In several parts of the north of England, but most
plentiful in Scotland.

Perennis. *Fl.* Maio, Junio.

Radix subtuberosa, repens. *Caulis* simplex, erectus, spi-
thamæus, infernè nudiusculus, apice foliosus. *Folia*
conferta, patentia, lanceolata, integerrima, glabra, ve-
nosa. *Pedunculi* terminales, aggregati, uniflori, pa-
tentes. *Flores* nivei, elegantissimi. *Calycis* foliola
subulata. *Corolla* ad basin ferè divisa, persistens. *Sta-*
mina numero varia, filiformia. *Antheræ* purpuræ.
Germen superum, globosum. *Stylus* longitudine sta-
minum. *Stigma* emarginatum. *Bacca* globosa, mem-
branacea. *Semina* plurima, gibba, punctata, recepta-
culo globoso, centrali, affixa.

Classis VIII.

Classis VIII.

OCTANDRIA.

Stamina 8.

MONOGYNIA.

Stylus 1.

** Fl. completi.*

195. ACER. *Pet.* quinque. *Cal.* quinquefidus, inferus. *Capsf.* monospermæ, alatæ.
190. EPILOBIUM. *Pet.* quatuor. *Cal.* quadrifidus, superus. *Capsf.* quadrilocularis. *Sem.* papposa.
191. CHLORA. *Cor.* octopartita. *Cal.* octophyllus, inferus. *Capsf.* unilocularis.
192. VACCINIUM. *Cor.* monopetala. *Cal.* quadridentatus, superus. *Bacca.*
193. ERICA. *Cor.* monopetala. *Cal.* tetraphyllus, inferus. *Capsula.*

Monotropa Hypopithys.

** * Fl. incompleti.*

194. DAPHNE. *Cal.* corollinus, quadrifidus, inferus. *Bacca* monosperma.

DIGYNIA.

Styli 2.

Chrysosplenium. Scleranthus.

TRIGYNIA.

Styli 3.

196. POLYGONUM. *Cal.* quinquepartitus, corollinus, inferus. *Cor.* nulla. *Sem.* unicum, nudum.

TETRAGYNIA.

Styli 4.

198. ADOXA. *Cal.* semiinferus. *Cor.* supera, quadri- vel quinque-fida. *Capsf.* calyce tecta.
197. PARIS. *Cal.* tetraphyllus. *Pet.* quatuor. *Bacca* supera,
199. ELATINE. *Cal.* tetraphyllus. *Pet.* quatuor. *Capsf.* supera, quadrilocularis.

VIII. OCTANDR

I. MONOGYNIA.

190. EPILOBIUM. *G. Pl.* 188. *Juss.* 315
Gærtn. t. 31.

Cal. quadrifidus. *Petala* quatuor. *Capsula* ob-
 longa, infera. *Semina* papposa.

1. EPILOBIUM foliis sparsis lineari-lanceolatis *angusti-*
venosis glabris, floribus inæqualibus, flami- *folium.*
 nibus declinatis.

Epilobium angustifolium. *Sp. Pl.* 493. *Huds.* 161.
With. 366. *Sibth.* 121. *Abbot.* 84. *Curt.*
Lond. fasc. 2. *t.* 24. *Fl. Dan.* *t.* 289.

Lythimachia speciosa, quibusdam *Onagra* dicta,
 siliquosa. *Ruii Syn.* 310. *Baub. Hist.* v. 2.
 906.

Chamaenerion. *Ger. em.* 477.

Angl. Rose-bay Willow-herb. *Persian*, or *French*
Willow.

In pratis umbrosis et ad sepes.

Meadows near Sheffield, and in several other places
 of the north plentifully. *Ray.* Near Alton,
 Hants *Dill.* *Curt.* At Grays near Henley,
 Oxfordshire. *Lightf. in herb.*

Perennis. *Fl.* Julio.

Radix repens. *Caules* erecti, tripedales, teretes, foliosi,
 glabri, supernè ramosi. *Folia* lineari-lanceolata, sub-
 sessilia, acuta, glabra, integerrima, subinde dentata,
 subtùs glaucescentia. *Racemi* terminales, aphylli, mul-
 tiflori, erecti, corymbosi. *Flores* pedicellati, alterni,
 speciosi. *Calyx* ruber. *Petala* purpurea, quandoque
 alba. *Germen* ineanum, supernè rubescendum. *An-*
theræ rubræ, polline cæruleo.

Formosa planta, in hortis vulgaris, difficilè quidem ex-
stirpanda.

*hirsu-
tum.*

2. *EPILOBIUM* foliis semiamplexicaulibus ovato-lanceolatis hirsutis, caule ramosissimo, radice repente.

Epilobium hirsutum. Sp. Pl. 494. With. 366. Relb. 153. Sibth. 122. Abbot. 84. Curt. Lond. fasc. 2. t. 21. Fl. Dan. t. 326.

E. ramosum. Hudf. 162.

Lyfimachia filiquosa hirsuta magno flore. Raii Syn. 311.

L. filiquosa. Ger. em. 476.

L. purpurea. Fuchf. Hist. 491.

Angl. Great hairy Willow-herb. Codlings and cream.

In aquosis et salicetis frequens.

Perennis. Fl. Julio.

Radix stolonibus repens. Caules erecti, quadripedales, teretes, pilis horizontalibus hirsuti, ramosissimi, foliosi. Folia opposita, amplexicaulia, ovato-lanceolata, acuta, denticulata, villosa; superiora alterna. Racemi foliosi. Flores rosei, magni. Petala obcordata, regularia, calyce duplò longiora. Germen villosum.

*parviflo-
rum.*

3. *EPILOBIUM* foliis sessilibus lanceolatis pubescentibus, caule subsimplici villosa, radice fibrosâ.

Epilobium parviflorum. With. 367. Sym. Syn. 92.

E. villosum. Curt. Lond. fasc. 2. t. 22. Relb. 152. Sibth. 121. Abbot. 84.

E. hirsutum. Hudf. 161. Sp. Pl. 494 β.

E. foliis longis villosis dentatis flore parvo. Hall. Enum. 410. Fl. Dan. t. 347.

Lyfimachia filiquosa hirsuta parvo flore. Raii Syn. 311. Moris. sect. 3. t. 11. f. 4.

Angl.

Angl. Small-flowered hoary Willow-herb.

In aquosis et ad ripas fluviorum.

Perennis. Fl. Julio.

Radix fibrosa. Caulis simplex, vel supernè tantum ramosus, bipedalis, teres, foliosus, mollissimè ac densissimè lanatus. Folia opposita, sessilia, denticulata, utrinque mollissimè villosa; superiora alterna. Racemi foliosi. Flores quam in præcedente quadruplò minores, purpurei.

4. **EPILOBIUM** foliis petiolatis ovatis dentatis, *montanum.*
caule tereti, stigmate quadripartito.

Epilobium montanum. *Sp. Pl.* 494. *Huds.* 162.

With. 367. *Relb.* 153. *Sibth.* 122. *Abbot.* 85. *Curt. Lond. fasc.* 3. t. 24. *Fl. Dan.* t. 922.

Lyfimachia campestris. *Raii Syn.* 311. *Ger. em.* 478.

Angl. Broad smooth-leaved Willow-herb.

In nemorosis, saxosis, vel ad muros.

Perennis. Fl. Julio.

Radix subrepens. Caulis erectus, strictus, sesquipedalis, subsimplex, teres, foliosus, parùm pubescens; rubicundus. Folia inferiora opposita, brevius petiolata, ovata, dentata, glabra, venis incanis; superiora alterna, angustiora. Racemus foliosus, pauciflorus. Flores purpurei, magnitudine præcedentis.

5. **EPILOBIUM** foliis petiolatis ovatis dentatis, *roseum.*
caule subtetragono, stigmate indiviso.

Epilobium roseum. *Sym. Syn.* 198.

Angl. Pale smooth-leaved Willow-herb.

In uliginosis.

Found in Lambeth marsh by Mr. Curtis, and at Morton near Ongar in Essex, by Mr. Edward Forster jun. *Symons.*

Perennis.

Perennis. Fl. Julio.

Caulis erectus, ramosus, sesquipedalis, supernè obsoletè quadrangulus, subincanus. Folia ferè ut in *E. montano*. Racemi foliosi. Flores parvi, carnei, vel pallidè rosei, venosi. Stigma clavatum, indivisum. Ance sequentis varietas?

tetrago- 6. **EPILOBIUM** foliis lanceolatis denticulatis,
num. caule tetragono, stigmate indiviso.

Epilobium tetragonum. *Sp. Pl.* 494. *Huds.* 162. *With.* 368. *Relb.* 154. *Sibth.* 122. *Abbot.* 85. *Curt. Lond. fasc.* 2. t. 23. *Fl. Dan.* t. 1029.

Lyfimachia filiquosa glabra media five minor.
Raii Syn. 311.

Angl. Square-stalked Willow-herb.

In palustribus et ad fossas.

Perennis. Fl. Julio.

Caulis erectus, ramosus, quadrangulus, glabriusculus, foliosus. Folia subsessilia, quandoque evidentiùs petiolata, lanceolata, inæqualiter denticulata, glabra; inferiora tantum opposita. Racemi pauciflori, foliosi. Flores purpurei. Stigma clavatum, indivisum.

palustre. 7. **EPILOBIUM** foliis sessilibus lanceolatis subdenticulatis, caule tereti, stigmate indiviso.

Epilobium palustre. *Sp. Pl.* 495. *Huds.* 163. *With.* 368. *Relb.* 154. *Sibth.* 123. *Abbot.* 85. *Dickf. H. Sicc. fasc.* 2. 15. *Engl. Bot.* t. 346.

Lyfimachia filiquosa glabra minor angustifolia.
Raii Syn. 311.

Angl. Round-stalked marsh Willow-herb.

In palustribus et ad fossas.

Perennis. Fl. Julio.

Caulis erectus, ramosus, teres, glabriusculus, foliosus.
Folia

Folia opposita, sessilia, subamplexicaulia, lineari-lanceolata, denticulata, sæpe integerrima, glabra. Racemi foliosi. Flores pallidè purpurei. Stigma lineare, indivisum.

8. *EPILOBIUM* foliis oppositis elliptico-lanceo-*alpinum*.
latis subintegerrimis, caule basi decumbente
paucifloro.

Epilobium alpinum. *Sp. Pl.* 495. *Huds.* 163.

With. 368. *Lightf.* 199. *t.* 10. *f.* 1. *Fl.*

Dan. *t.* 332. *Dicks. H. Sicc. fasc.* 2. 14.

Lyfimachia filiquosa glabra minor latifolia. *Raii*
Syn. 311.

L. prunellæ foliis. *Bocc. Mus.* *t.* 108.

Angl. Alpine Willow-herb.

Ad rivulos alpinos.

On the sides of the Cheviot hills. *Ray.* On Ben
Lomond, and other mountains of Scotland.

Perennis. *Fl.* Junio, Julio.

Radix repens. Caules palmares, basi decumbentes, simplices, subtetragoni. Folia opposita, breviter petiolata, elliptica, obsoletè denticulata, glabra; summa alterna. Racemus terminalis, subtriflorus, foliosus. Flores parvi, rubri. Germen ferè glabrum.

191. *CHLORA.* *Syst. Nat.* *v.* 2. 267. *Juss.* 142.

Cal. octophyllus. *Cor.* octopartita. *Stigma* quadrifidum. *Capsula* supera, unilocularis, bivalvis, polysperma.

1. *CHLORA* foliis perfoliatis.

Chlora perfoliata. *Syst. Nat. ed.* 12. *v.* 2. 267. *perfolia-*
ta.

Huds. 168. *With.* 369. *Relb.* 157. *Sibth.* 123.

Abbot. 86. *Engl. Bot.* *t.* 60. *Dicks. H. Sicc.*
fasc. 14. 13.

Chlora. *Rencalm.* *t.* 76.

Gentiana perfoliata. *Sp. Pl.* 335.

Blackstonia perfoliata. *Hudsf. ed.* 1. 146.

Centaureum luteum perfoliatum. *Raii Syn.* 287.

C. parvum luteum Lobelii. *Ger. em.* 547.

Angl. Yellow Centaury.

In apricis calcareis.

At Cherry Hinton, Cambridgeshire, and in Belton park, Lincolnshire. *Rev. Archdeacon Peirson.* St. Vincent's rocks, Bristol.

Annua. Fl. Julio, Augusto.

Radix parva, tortuosa. Herba glauca, intensè amara.

Caulis erectus, teres, foliosus, supernè dichotomus.

Folia perfoliata, integerrima, glabra, ovata, acuta.

Flores e dichotomiâ caulis, solitarii, pedunculati, aurei, stigmatè miniato, bipartito, laciniis bifidis.

192. VACCINIUM. *G. Pl.* 191. *Juss.* 162.

Gærtn. t. 28.

Cor. monopetala. Stamina receptaculo inserta.

Antherae apice poris duobus. Bacca infera, quadrilocularis, polysperma.

* *Foliis deciduis.*

Myrtillus. 1. VACCINIUM pedunculis unifloris, foliis ovatis ferratis deciduis, caule angulato.

Vaccinium Myrtillus. *Sp. Pl.* 498. *Hudsf.* 163.

With. 370. *Sibth.* 123. *Abbot.* 86. *Engl.*

Bot. t. 456. *Dicksf. H. Sicc. fasc.* 5. 12. *Fl.*

Dan. t. 974.

Vaccinia nigra. *Ger. em.* 1415.

Vitis Idæa angulosa. *Raii Syn.* 457.

Angl. Bilberry, or Bleaberry.

In ericetis montosis petrosis, et fylvis solo spongiofo.

Frutex. Fl. Maio.

Fruti-

Fruticulus ramosissimus, vix pedalis. Rami angulati, glabri, virentes, tortuosi. Folia alterna, petiolata, ovata, ferrata, glabra, lætè viridia, decidua, exstipulata. Flores axillares, solitarii, pedunculati, cernui, carnei, globosi, ore quinquedentati. Calyx quinquedentatus, parvus. Stamina inclusa, sæpe 10. Antheræ supernè bicornes. Germen obconicum, viride, glabrum. Stigma simplex. Bacca globosa, nigra, umbilicata, quinquelocularis, acida, edulis.

2. VACCINIUM pedunculis unifloris, foliis ob- *uligin-*
ovatis integerrimis glabris, ramis teretibus. *sum.*

Vaccinium uliginosum. *Sp. Pl.* 499. *Huds.* 164.

With. 370. *Fl. Dan. t.* 231. *Engl. Bot. t.* 581.

Vitis Idæa magna quibusdam, five Myrtillus grandis. *Raii Syn.* 457. *Baub. Hist. v.* 1.518.

V. Idæa foliis subrotundis major. *Ger. em.* 1416.

Angl. Great Bilberry.

In ericetis montosis paludosis.

At Gambleby, six miles from Penrith, Cumberland; also in Whinfield forest, Westmoreland.

Ray. In the highlands of Scotland. *Lightf.*

Frutex. Fl. Aprili, Maio.

Præcedente altior, ramis ramulisque teretibus. Folia decidua, obovata, obtusa, rariùs acutiuscula, integerrima, subrevoluta, venosa, glabra, subtus glauca. Flores carnei, sæpiùs quadrifidi, octandri. Antheræ, aristatæ. Baccæ nigrae, rore cæcio. Semina striata.

* * *Foliis sempervirentibus.*

3. VACCINIUM racemis terminalibus nutanti- *Vitis*
bus, foliis obovatis revolutis denticulatis sub- *Idæa.*
tùs punctatis.

Vaccinium Vitis Idæa. *Sp. Pl.* 500. *Huds.* 164.

With. 371. *Engl. Bot. t.* 593. *Dicks. H.*

Sicc. fasc. 5. 10. *Fl. Dan. t.* 40.

Vaccinia

Vaccinia rubra. *Ger. em.* 1415.

Vitis Idæa sempervirens, fructu rubro. *Raii*
Syn. 457.

Angl. Red Whortle-berry. Cow-berry.

In ericetis turfosis petrosis elatis.

Plentiful in Derbyshire, as well as other mountainous parts of England, Scotland, and Wales.

Frutex. Fl. Junio.

Fruticulus humilis, ferè herbaceus, at sempervirens.

Rami tortuosi, angulati. Folia obovata, retusa, glaberrima, nitida, venosa, subtus punctata pallida, apicem versus obsolete ferrata vel dentata, nequaquam integerrima. Racemi nutantes. Bractæ sub singulo flore solitariae, ellipticae, concavae, ciliatae. Flores campanulati, cernui, carnei, quadrifidi. Antherae muticae. Baccæ rubrae, amaricantes, inter Suecos saccharo conditæ usitatissimæ, ut apud nos *Ribes rubrum*.

Oxycoccus.

4. *VACCINIUM* foliis ovatis integerrimis revolutis acutis, caulibus repentibus filiformibus glabris.

Vaccinium Oxycoccus. *Sp. Pl.* 500. *Huds.* 164.

With. 372. *Relb.* 154. *Abbot.* 86. *Engl.*

Bot. t. 319. *Dicks. H. Sicc. fasc.* 5. 11. *Fl.*

Dan. t. 80.

Vaccinia palustris. *Ger. em.* 1419. *Lob. Ic.*

v. 2. 109.

Oxycoccus, seu *Vaccinia palustris.* *Raii Syn.*

267.

Angl. Cranberry.

In paludibus turfosis muscosis.

Plentiful in Scotland, Ireland, and the North of England. On Derfingham moor, Norfolk. *Mr. Humphrey.*

Perennis. Fl. Junio.

Radix fibrosa. Caules perennantes, suffruticosi, prostrati, radicantes,

radicantes, filiformes, glabri, foliosi. Folia alterna, erecta, petiolata, parva, ovata, revoluta, glaberrima, sempervirentia, subtus glauca. Pedunculi terminales, aggregati, uniflori, bracteolati. Flores cernui, pulcherrimè rosei, ad basin usque quadrifidi, laciniis reflexis. Stamina exserta. Stigma acutum. Bacca maculosa, quadrilocularis, edulis.

193. ERICA. *G. Pl.* 192. *Juss.* 160.

Gærtn. t. 63.

Cal. tetraphyllus. *Cor.* quadrifida. *Stamina* receptaculo inserta. *Antheræ* poris duobus. *Capsula* supera, quadrilocularis, polysperma.

* *Antheris aristatis.*

1. ERICA antheris aristatis inclusis, stylo exserto, *vulgaris*. corollâ quadripartitâ calyce brevior, foliis oppositis calcaratis.

Erica vulgaris. Sp. Pl. 501. *Huds.* 165. *With.* 374. *Relb.* 155. *Sibth.* 124. *Abbot.* 87. *Curt. Lond. fasc.* 5. *t.* 30. *Fl. Dan. t.* 677. *Raii Syn.* 470. *Ger. em.* 1380. *Camer. Epit.* 75.

β. *E. vulgaris hirsuta. Raii Syn.* 471.

E. ciliaris. Huds. ed. 1. 144.

Angl. Common Heath or Ling.

In ericetis et sylvis passim.

Frutex. Fl. Junio, Julio.

Caulis lignosus, ramosissimus, ramis teretibus, glabris, foliosis, supernè racemosis, deinde apice foliosis. Folia duplici serie opposita, hinc quadrifaria, parva, triquetra, obtusa, basi soluta vel calcarata, plerumque glabra; in β pubescenti-incana. Flores pedunculati, cernui, purpureo-incarnati, nitidi, emarcati. Calyx duplex; exterior e foliolis quatuor, ovatis, carinatis, ciliatis, virentibus; interior quadruplò major, coloratus, corollam amplectens. Corolla quadripartita, calyce interiori

duplò brevior, laciniis acutis, rectis. Stamina longitudine ferè corollæ. Antheræ conniventes, basî biaristatæ. Germen pubescens. Stylus calyce paulò longior. Stigma quadrilobum. Capsula depressa, quadrilocularis, quadrivalvis. Dissepimenta simplicia, columnellæ inserta, suturis opposita.

Tetralix. 2. *ERICA* antheris aristatis, stylo incluso, corollâ ovatâ, foliis quaternis ciliatis, floribus capitatis.

Erica Tetralix. *Sp. Pl.* 502. *Syst. Veg. ed.* 14. 365. *Huds.* 166. *Witb.* 373. *Relb.* 156. *Sibth.* 124. *Abbot.* 87. *Curt. Lond. fasc.* 1. t. 21. *Dickf. II. Sicc. fasc.* 9. 4. *Fl. Dan.* t. 81.

E. Brabantica, folio *Coridis* hirsuto quaterno.

Raii Syn. 471.

Angl. Cross-leaved Heath.

In ericetis humidiusculis.

Frutex. Fl. Julio, Augusto.

Caulis determinatè ramosus, ramis erectis, foliosis, sæpius hirsutis. Folia quaterna, petiolata, patentia, lineari-ovata, revoluta, hispida, pilis apice glandulosis. Flores terminales, umbellato-capitati, secundi, pulcherrimè incarnati. Calyx ciliatus, foliolis duobus calyculatus. Corolla elliptico-globosa, ore coarctata, quadridentata. Antheræ basî biaristatæ. Germen sericeum, quadrifurcatum. Stigma obtusum. Capsula quadrivalvis, dissepimentis e medio valvularum ut in *Andromedâ*.

* * *Antheris cristatis*.

cinerea. 3. *ERICA* antheris cristatis, stylo subexserto, stigmate capitato, corollâ ovatâ, foliis ternis.

Erica cinerea. *Sp. Pl.* 501. *Syst. Veg. ed.* 14. 366. *Huds.* 165. *Witb.* 374. *Relb.* 156. *Sibth.* 125. *Curt. Lond. fasc.* 2. t. 25. *Fl. Dan.* t. 38.

E. tenuifolia. *Ger. em.* 1382. *Raii Syn.* 471.
Angl. Fine-leaved Heath.

In ericetis vulgaris.

Frutex. Fl. Julio, Augusto.

Caulis ramosus, ramulis erectis, foliosis. Folia ternâ, triquetro-linearâ, glabra, dorso canaliculata. Racemi terminales, verticillati, foliosi. Flores pedicellati, nuntantes, scariosi, cæruleo-incarnati. Calyx glaber, foliolis binis, minutis, calyculatus. Corolla ellipticoglobosa, ore contracta, quadrifida. Antheræ basi cristatæ, appendiculis duobus ferratis. Germen octosulcatum. Stignia capitatum. Capsula quadrivalvis, dissepimentis e medio valvularum.

* * * *Antheris muticis.*

4. ERICA antheris muticis styloque exsertis, *co-vagans.*
 rollâ campanulatâ; foliis quaternis, floribus
 axillaribus confertis.

Erica vagans. *Mant.* 2. 230. *Engl. Bot.* t. 3.
With. 373. *Dicks. Dr. Pl.* 62.

E. multiflora. *Huds.* 166.

E. didyma. *With. ed.* 2. 400.

E. foliis Corios multiflora. *Raii Syn.* 471. nec
 tamen Bauhini.

Angl. Cornish Heath.

In ericetis Cornubiæ copiosè.

Frutex. Fl. Julio, Augusto.

Caulis determinatè ramosissimus, ramis erectis, foliosis, fulcatis, glabris. Folia oblique quaterna, petiolata, lineari-triquetra, glabra, dorso canaliculata. Flores laterales, axillares, conferti, pedunculati, patentes, rubri. Pedunculi capillares, colorati, medium versus bractæolati. Calyx simplex, glaber. Corolla campanulata, ore quadrifido, patulo. Antheræ exsertæ, basi muticæ, apice bipartitæ, obtusæ. Stylus staminibus longior. Stigma quadrifidum.

Variat flore albo.

Dabeoci. 5. *ERICA* antheris muticis styloque inclusis, corollâ ovatâ, foliis oppositis alternisve ovatis subtùs incanis.

Erica Dabeoci. *Sp. Pl.* 509. *Huds.* 166. *Engl. Bot. t.* 35. *Dicks. H. Sicc. fasc.* 9. 5. *With.* 372.

E. Cantabrica, flore maximo, foliis *Myrti* subtùs incanis. *Dill. in Raii Syn.* 472.

Andromeda Daboecia. *Syst. Veg. ed.* 13. 338. *ed.* 14. 406. *With. ed.* 2. 425.

Vaccinium Cantabricum. *Huds. ed.* 1. 143. *Angl. Irish Heath.*

In montibus Hiberniæ occidentalis, solo squalido et spongioso. *Ray.*

On Croagh Patrick in the county of Mayo. *A. B. Lambert, Esq.*

Frutex. Fl. Junio, Julio.

Caules graciles, erectiusculi, ramosi, teretes, foliosi. Folia inferiora opposita, quandoque ternâ; superiora alterna; ovata, integerrima, revoluta, scabriuscula; subtùs tomentosa, nivea. Racemi terminales, subnutantes, foliolosi, pilosi. Flores alterni, pedunculati, penduli, purpurei. Calyx piloso-glandulosus, simplex. Corolla ovata, subtetragona, ore quadrifido, reflexo. Antheræ lineari-oblongæ, muticæ, basi bilobæ. Stigma obtusum. Capsula quadrivalvis, quadrilocularis, dissepimentis e margine inflexo valvularum, quod nunquam in veris *Andromedis* conspiciendum.

194. *DAPHNE.* *G. Pl.* 192. *Juss.* 77.

Cal. corollinus, quadrifidus, marcescens, stamina includens. *Bacca* monosperma.

Mezereum. 1. *DAPHNE* floribus sessilibus ternis caulinis, foliis lanceolatis deciduis.

Daphne Mezereum. *Sp. Pl.* 509. *Huds.* 167. *With.*

With. 376. *Fl. Dan.* t. 268. *Woodv. Med. Bot.* t. 23.

Daphnoides. *Fuchsf. Hist.* 227.

Chamælea germanica, five *Mezereon.* *Ger. em.* 1402.

Mezereum, radicis cortex. *Pharmac. Lond.*
Angl. *Mezereon.* Spurge Olive.

In nemorosis.

In woods near Andover plentifully. *Miller.* At Laxfield, Suffolk. *Mr. Woodward.* In Needwood forest, Mr. Pitt. *With.*

Frutex. Fl. Martio, Aprili.

Rami viminei, medio floriferi ante foliorum expansionem, apice demum foliosi. Folia decidua, lanceolata, integerrima, glabra. Flores sparsi, sessiles, basi bracteati, rosei, odori. Baccæ ellipticæ, rubræ.

2. *DAPHNE* racemis axillaribus subquinqueflo- *Laureo-*
ris, foliis lanceolatis glabris. *la.*

Daphne Laureola. *Sp. Pl.* 510. *Huds.* 167.
With. 377. *Relb.* 157. *Sibth.* 125. *Ab-*
bot. 87. *Engl. Bot.* t. 119. *Jacq. Austr.*
t. 183.

Laureola. *Raii Syn.* 465. *Ger. em.* 1404.

L. et Daphnoides. *Lob. Ic.* 368.

Angl. Spurge Laurel.

In dumetis et ad sepes.

Frutex. Fl. Martio.

Caulis tripedalis, teres, glaber, ramosus, apice foliosus. Folia conferta, dependentia, lætè viridia, glabra, integerrima, sempervirentia. Racemi axillares, simplices, nutantes, bracteati. Flores virides. Baccæ nigræ.

195. *ACER.* *G. Pl.* 546. *Juss.* 251. *Gærtn. t.* 116.

Cal. quinquefidus. *Petala* 5. *Capsulæ* 2 vel 3,
L e 3 superæ,

superæ, monospermæ, alâ terminatæ.
Flores aliquot masculi.

Pseudo-platanus. *1. ACER foliis quinquelobis, inæqualiter ferratis, racemis pendulis.

Acer Pseudo-platanus. *Sp. Pl.* 1495. *Huds.* 445.
With. 368. *Relb.* 380. *Sibth.* 127. *Abbot.* 219. *Engl. Bot. t.* 303.

A. majus. *Raii Syn.* 470. *Ger. em.* 1484.

β. Var. 2. *With.* 369.

Angl. Greater Maple. *Sycamore.*

In sepibus, et ad domos, vix indigena.

Arbor. Fl. Maio.

Inter formosiores jure numeratur arbores, neque cœli nostri intemperiem timet. Caudex teres, rectus, cortice lævi, cinereo. Rami undique patuli, ramulis dependentibus. Folia opposita, petiolata, quinqueloba, ferrata, glabra, subtus pallidiora, exstipulata. Racemi axillares, solitarii, penduli, multiflori. Flores pedicellati, virides, pedicellis villosis. Calycis laciniae lanceolatae, æquales. Petala calyci conformia. Stamina subulata, plerumque glabra. Germen hirsutissimum. Stylus simplex. Stigmata duo vel tria, revoluta, pubescentia. Capsulae pendulae, binæ vel ternæ, alis divaricatis, securiformibus, obtusis, venosis, glabris. Semen reniforme. Flores pauciores masculi aliis immixti sunt.

Var. β gaudet foliis subtus glaucis, ferraturis remotioribus, flaminibus hirsutis, monente amicissimo D. Andr. Caldwell, Soc. Linn. Socius.

campestre. 2. ACER foliis lobatis obtusis incis, racemis erectis.

Acer campestre. *Sp. Pl.* 1497. *Huds.* 445.
With. 369. *Relb.* 380. *Sibth.* 127. *Abbot.* 220. *Engl. Bot. t.* 304.

A. minus. *Raii Syn.* 470. *Ger. em.* 1484.

β. A.

β. A. campestre et minus, fructu rubente. *Dill.*

in Raii Syn. 470.

Angl. Common Maple.

In sepibus et dumetis Angliæ australioris vulgaris.

Arbor. Fl. Maio, Junio.

Præcedente longè humilior, magisque patula, cortice sub-
 erofo, rimoso, in ramulis lævi. Folia opposita, petio-
 lata, parva, quinqueloba, lobis subincisis, sæpe inte-
 gerrimis. Racemi terminales, erecti, ramosi, basi brac-
 teati. Pedunculi ut et petioli pubescunt. Flores vi-
 rescentes, staminibus flavis, stylo brevissimo. Alæ cap-
 sularum divaricato-patentissimæ, rubicundæ. Lignum
 durum, compactum.

II. TRIGYNIA.

196. POLYGONUM. *G. Pl.* 195. *Juss.* 82.

Gærtn. t. 119.

Cal. quinquepartitus, coloratus, corollinus, per-
 sistens. *Semen* unicum, superum, angulatum,
 calyce tectum. *Stamina* et *pistilla* numero
 incerta.

* *Digyna.*

I. POLYGONUM floribus pentandris semidi-*amphibi-*
 gynis, spicâ ovatâ. *um.*

Polygonum amphibium. *Sp. Pl.* 517. *Huds.* 169.

With. 379. *Relb.* 158. *Sibth.* 128. *Curt.*

Lond. fasc. 4. *t.* 28. *Engl. Bot. t.* 436. *Fl.*

Dan. t. 282.

Perficaria salicis folio perennis, *Potamogiton angustifolium dicta*. *Raii Syn.* 145.

Potamogiton angustifolium. *Ger. em.* 821.

Amphibious Arsmart. *Pet. H. Brit. t. 3. f. 12.*

Angl. *Amphibious Perficaria*.

In fossis, stagnis, vel ad eorum margines.

Perennis. Fl. Julio, Augusto.

Radix repens. Caules radicanes, erecti vel natantes, stipulis tubulosis, membranaceis, integris, vestiti. Folia alterna, petiolata, lanceolata, acuta, ferrulata, glabra, lætè viridia. Spica terminalis, subfolitaria, pedunculata, erecta, ovata, densa, multiflora, basi bracteata. Flores pedicellati, rosei, nec inelegantes. Calyx infundibuliformis, laciniis obtusis, patulis. Stamina tantum quinque, in fauce calycis, longitudine varia. Stylus semibifidus. Stigmata globosa. Semen ovatum, compressum, angulis obtusis.

Perficaria. 2. *POLYGONUM* floribus hexandris semi-digynis, spicis ovato-oblongis, erectis, pedunculis lævibus stipulis ciliatis. *Curt.*

Polygonum Perficaria. *Sp. Pl.* 518. *Huds.* 169.

With. 380. *Relb.* 159. *Sibth.* 128. *Curt.*

Lond. fasc. 1. t. 23. *Fl. Dan. t. 702.*

Perficaria maculosa. *Raii Syn.* 145. *Ger. em.* 445.

Spotted Arsmart. *Pet. H. Brit. t. 3. f. 37.*

β. *Perficaria folio subtus incano*. *Dill. in Raii Syn.* 145.

Spotted hoary Arsmart. *Pet. H. Brit. t. 3. f. 8.*

γ. *Perficaria foliis Salicis albæ vulgaris*. *Dill. in Raii Syn.* 145.

Silver Arsmart. *Pet. H. Brit. t. 3. f. 9.*

Angl. *Spotted Perficaria*.

In fossis et palustribus vulgaris.

Annua.

Annua. Fl. Julio, Augusto.

Radix fibrosa. Caulis erectus, basi quandoque radicans, bipedalis, alternatim ramosus, teres, foliosus, glaber, supra genicula incrassatus, sæpe rubicundus. Folia subpetiolata, patenti-deflexa, lanceolata, acuta, integerrima, margine nervoque scabra, maculâ nigrâ, retrorsum lunatâ, plerumque notata. Stipulæ tubulosæ, amplexicaules, breves, margine ciliatæ. Spicæ terminales, subinde axillares, pedunculatæ, erectæ, obtusæ, pedunculis lævibus. Flores rosei. Stamina longitudine calycis. Stylus ad medium usque bifidus, subinde trifidus. Semen subtriquetrum.

3. POLYGONUM floribus hexandris digynis, *lapathi-*
pedunculis scabris, stipulis muticis, seminibus *folium.*
utrinque concavis. *Curt.*

Polygonum lapathifolium. *Sp. Pl.* 517. *Ait.*

Hort. Kew. v. 2. 30. *Sibth.* 129. *Syn.*

Syn. 96. *Abbot.* 89.

P. pensylvanicum. *Curt. Lond. fasc.* 1. t. 24 &
25. *Relb.* 160.

P. Persicaria ζ. *Huds.* 170.

P. pallidum. *With.* 381.

Persicaria mitis major, foliis pallidioribus. *Raii*
Syn. 145.

Hydropiper. *Lob. Ic.* 315. *Ger. em.* 445.

β. caule et floribus rubris. *Curt. sub t.* 25.

γ. caule maculato. *Curt. t.* 25.

Persicaria latifolia geniculata, caulibus maculatis.
Dill. in Raii Syn. 145.

δ. foliis subtus incanis. *Curt. sub t.* 25.

Persicaria maculosa procumbens, foliis subtus
incanis. *Dill. in Raii Syn.* 146.

White Branched Arsmart. *Pet. H. Brit. t.* 3.
f. 10.

Angl. Pale-flowered *Persicaria*.

In fimetis et cultis.

Annua.

Annua. Fl. Julio, Augusto.

Radix fibrosa. Caulis ramosus, patulus, quandoque decumbens, supra genicula tumidus. Folia subtus punctato-glandulosa, subinde pubescentia, supra quandoque maculata. Stipulae nervosae, muticae, rarius subciliatae. Spicae breves, obtusae, erectae, demum subnutantes. Pedunculi glanduloso-scabri. Flores albido-virescentes, vel rubicundi. Stamina vix calyci aequalia. Stylus ad basin ferè bipartitus. Semen utrinque excavatum, nec gibbum, subinde triquetrum.

Hydropiper.

4. POLYGONUM floribus hexandris semidigynis, foliis lanceolatis undulatis immaculatis, spicis filiformibus nutantibus, caule erecto.

Polygonum Hydropiper. *Sp. Pl.* 517. *Huds.* 170.

With. 379. *Relb.* 160. *Sibth.* 129. *Curt.*

Lond. fasc. 1. t. 26.

Perficaria vulgaris acris seu Hydropiper. *Raii*

Syn. 144.

Hydropiper. *Fuchs. Hist.* 843.

Common Arsmart. *Pet. H. Brit. t. 3. f. 5.*

Angl. Biting Perficaria.

In aquosis et ad fossas frequens.

Annua. Fl. Septembri.

Radix fibrosa. Caulis erectus, bipedalis, ramosus, teres, supra genicula tumidus, demum rubicundus. Folia lanceolata, undulata, pallida, immaculata. Stipulae ciliatae, obsoletè nervosae. Spicae terminales, lineares, laxae, nutantes. Flores pallidè rubri, extus punctato-glandulosi. Stamina longitudine calycis. Stylus ultra medium ferè bifidus. Stigmata obtusa. Semen compressum.

minus.

5. POLYGONUM floribus hexandris submonogynis, foliis lineari-lanceolatis planis, spicis filiformibus erectiusculis, caule basi radicante.

Polygonum minus. *Huds. ed. 1.* 148. *Curt.*

Lond. fasc. 1. t. 28. With. 380. *Relb.* 161.

Sibth.

Sibth. 130. *Ait. H. Kew.* v. 2. 31. *Abbot.* 89.

P. Hydropiper β . *Huds.* 170.

Perficaria pusilla repens. *Ger. em.* 446. *Raii Syn.* 145.

P. minor. *Morif. sect.* 5. t. 29. f. 5.

β . *P. angustifolia* ex singulis geniculis florens. *Raii Syn.* 145 ?

Angl. Small Creeping *Perficaria*.

In inundatis glareosis.

Plentiful in Tothill fields, Westminster. *Huds. Curt.* In a gravel pit on Malvern Chace, Worcestershire. *Dr. Stokes.* On Putney Common. *Rev. Dr. Goodenough.* On Costesy Common, near Norwich.

Annua. Fl. Septembri.

Caules plures, basi decumbentes et radicales, supernè erecti, spithamæi, parùm ramosi, foliosi, supra genicula incrassati. Folia lineari-lanceolata, plana, glabra, ferè avenia. Stipulæ ciliatæ. Spicæ erectiusculæ, filiformes, laxæ. Flores pallidi, extùs rubri, impunctati. Stamina longitudine calycis. Stylus apice tantum bifidus. Stigmata obtusa.

* * *Trigyna.*

6. **POLYGONUM** caule simplicissimo monostachyo, foliis ovatis undulatis in petiolum decurrentibus. *Bistorta.*

Polygonum Bistorta. *Sp. Pl.* 516. *Huds.* 168. *With.* 382. *Relb.* 158. *Sibth.* 128. *Abbot.* 88. *Curt. Lond. fasc.* 1. t. 22. *Woodv. Med. Bot.* t. 34. *Engl. Bot.* t. 509. *Dreves. Bilderb.* t. 39.

Bistorta major. *Raii Syn.* 147. *Ger. em.* 399.

Bistorta, radix. *Pharmac. Lond.*

Angl. Great Bistort, or Snakeweed.

In

In pratis et pascuis.

Most plentiful in the northern counties. At
Heigham near Norwich. *Mrs. Kett.*

Perennis. Fl. Junio.

Radix repens, lignosa, tortuosa. Caulis simplicissimus, sesquipedalis, erectus, strictus, foliosus. Folia ovata, integerrima, undulata, venosa, subtus glauca, radicalia præcipuè in petiolum decurrentia. Spica terminalis, erecta, cylindrica, obtusa, densa, multiflora. Flores rosei, octandri, trigyni, pedicellati, pedicellis basi bracteatis. Semen triquetrum.

*vivipa-
rum.*

7. POLYGONUM caule simplicissimo monostachyo, foliis lanceolatis margine revolutis.

Polygonum viviparum. *Sp. Pl.* 516. *Huds.* 169.

With. 383. *Lightf.* 206. *Fl. Dan.* t. 13.

Engl. Bot. t. 669.

Bistorta minor. *Raii Syn.* 147. *Ger. em.* 399.

B. minor nostras. *Morif. sect.* 5. t. 28. f. ult.

Small Bistort. *Pet. H. Brit.* t. 5. f. 2.

8. *Bistorta minima alpina*, foliis imis subrotundis et minutissimè serratis. *Raii Syn.* 147.

Welch Bistort. *Pet. H. Brit.* t. 5. f. 3.

Angl. Alpine Bistort.

In pratis et pascuis alpinis.

In several parts of Westmoreland, and the north of Yorkshire. *Ray.* *Huds.* In dry stony pastures by the sides of the Scottish mountains not unfrequent. *Lightf.* Near the top of Ben Lomond.

Perennis. Fl. Junio, Julio.

Habitus præcedentis, sed longè minor. Caulis palmaris, vix spithameus. Folia lineari-lanceolata, venosa, margine revoluta, subtus quandoque hirsuta; radicalia sæpe elliptico-subrotunda. Spica linearis, obtusa, floribus roseis, octandris, trigynis, inferioribus plerumque gemmiparis. Germen acutè triquetrum. Semina rariùs perfecta.

8. **POLYGONUM** floribus axillaribus, foliis elliptico-lanceolatis margine scabris, stipularum nervis remotis, caule procumbente herbaceo.

Polygonum aviculare. *Sp. Pl.* 519. *Huds.* 171. *With.* 383. *Relb.* 161. *Sibth.* 130. *Curt. Lond. fasc.* 1. t. 27. *Mart. Fl. Rust.* t. 91.

P. mas vulgare. *Raii Syn.* 146. *Ger. em.* 565.

Common Knot-grass. *Pet. H. Brit.* t. 10. f. 1.

β. *Polygonum* brevi angustoque folio. *Raii Syn.* 146.

Short-leaved Knot-grass. *Pet. H. Brit.* t. 10. f. 3.

γ. *Polygonum* oblongo angustoque folio. *Raii Syn.* 146.

Narrow Knot-grass. *Pet. H. Brit.* t. 10. f. 4.

δ. *Polygonum* folio rotundo. *Dill. in Raii Syn.* 147. Thick-set Knot-grass. *Pet. H. Brit.* t. 10. f. 2.

ε. *Polygonum* marinum. *Raii Syn.* 147. exclusis synonymis.

Angl. Knot-grass.

In ruderatis, plateis, cultis, et ad vias passim.

ε In arenosis maritimis.

Cornwall. *F. Borone.*

Annua. Fl. Aprili—Octobri.

Radix fibrosa, ramosa, sublignosa, tenacissima. Caules procumbentes, divaricati, ramosi, foliosi, teretes, striati. Folia alterna, elliptico-lanceolata, basi attenuata, latitudine varia, glabra, margine tantum scabriuscula. Stipulae acutae, scariosae, nervis paucis, remotis. Flores axillares, conferti, pedunculati, parvi nec inelegantes, albo viridi sanguineoque varii. Stamina octo, calyce breviora. Styli tres, brevissimi. Semen triquetrum, nitidum.

Polygonum maritimum Linnæi, quod Mediterraneis nascitur in oris, species distincta a *P. marino* Raii omnino est. Differt enim caule fruticoso, perenni, foliis latis, ellipticis, glaucis, margine vix scabris, at præcipuè stipularum

pularum nervis numerosis, approximatis. Huc pertinent synonyma omnia a Raio citata, & inde figura Petiveriana e J. Bauhino mutuata.

Fagopyrum. *9. **POLYGONUM** foliis cordato-sagittatis, caule erectiusculo inermi, feminum angulis æqualibus.

Polygonum Fagopyrum. Sp. Pl. 522. Hudf. 172.

With. 384. Sibth. 130. Mart. Fl. Rust. t. 46.

Fegopyrum. Raii Syn. 144.

Tragopyron. Ger. em. 89.

Angl. Buck-wheat. Pet. H. Brit. t. 2. f. 11.

In arvis.

Annua. Fl. Julio, Augusto.

Radix fibrosa. Caulis erectus, pedalis, subflexuosus, ramosus, teres, glaber, foliosus. Folia cordato-sagittata, lævia, integerrima; superiora sessilia. Stipulæ breves, integerrimæ, muticæ. Racemi axillares et terminales, paniculati. Flores incarnati, octandri, trigyni. Semen pyramidato-triquetrum, angulis rectis, nec sinuatis.

Convolvulus. 10. **POLYGONUM**-foliis cordato-sagittatis, caule volubili angulato, laciniis calycinis obtusè carinatis.

Polygonum Convolvulus. Sp. Pl. 522. Hudf. 171.

With. 385. Relb. 162. Sibth. 131. Curt.

Lond. fasc. 4. t. 29. Fl. Dan. t. 744.

Fegopyrum scandens sylvestre. Raii Syn. 144.

Volubilis nigra. Ger. em. 863.

Trailing Buck-wheat. Pet. H. Brit. t. 2. f. 12.

Angl. Climbing Buck-wheat. Black Bindweed.

In arvis et sepibus frequens.

Annua. Fl. Junio, Julio.

Radix fibrosa. Caulis volubilis, angulatus, striatus, subinde scabriusculus, supernè ramosus. Folia sagittata, subcordata, integerrima, glabra. Racemi terminales,
inter-

interrupti, foliolosi. Flores fasciculati, nutantes, albi, extus viridi-purpurei, laciniis tribus exterioribus obtusè carinatis, nec carinâ dilatatis. Stylus apice trilobus.

III. TETRAGYNIA.

197. PARIS. *G. Pl.* 198. *Juss.* 42.
Gærtn. t. 83.

Cal. tetraphyllus. *Petala* 4, angustiora. *Bacca* supera, quadrilocularis. *Antheræ* mediis filamentis adnatæ.

I. PARIS.

Paris quadrifolia. *Sp. Pl.* 526. *Huds.* 172. *quadrifolia*
With. 385. *Relb.* 163. *Sibth.* 131. *Ab-*
bot. 90. *Engl. Bot. t.* 7. *Dicks. H. Sicc.*
fasc. 13. 15. *Fl. Dan. t.* 139.

Herba Paris. *Raii Syn.* 264. *Ger. em.* 405.
Lob. Ic. 267. *Camer. Epit.* 835. *Tillandsf.*
Ic. 32.

Aconitum pardalianches. *Fuchsf. Hist.* 87. *Ic.* 48.
Angl. Herb Paris. Pet. H. Brit. t. 44. *f.* 8.

In nemorosis, haud vulgaris.

Perennis. *Fl. Maio.*

Radix repens. *Caulis* simplicissimus, pedalis, teres, glaber, apice foliosus, uniflorus. *Folia* quatuor, rarius tria vel quinque, verticillata, ovata, acuta, integerrima, tristè viridia, nervosa, glabra. *Pedunculus* terminalis, solitarius, erectus, angulatus. *Flos* viridis, inamœnus.
Calycis.

Calycis foliola ovata, acuminata, nervosa. Petala linearia, calyci alterna. Stamina petalis calycique opposita, parumque breviora, latere interiori medium versus antherifera. Antheræ lineares, filamentis adnatæ. Germen superum, subrotundum, tetracoccum, violaceum. Styli lineares, acuti, patentes. Bacca nigra, quadrilocularis. Semina numerosa, duplici ordine digesta, columellæ inserta.

198. ADOXA. *G. Pl.* 198. *Juss.* 309.
Gærtn. t. 112.

Cal. bifidus vel trifidus, semiinferus. *Cor.* quadri- vel quinque-fida, supera. *Capsula* quadri- vel quinque-locularis, calyci coalita.

Moschatellina. 1. ADOXA.

Adoxa Moschatellina. *Sp. Pl.* 527. *Huds.* 172.
With. 386. *Relb.* 163. *Sibth.* 131. *Abbot.* 90. *Curt. Lond. fasc.* 2. t. 26. *Dicks. H. Sicc. fasc.* 5. 7. *Engl. Bot. t.* 453. *Fl. Dan. t.* 94.

Moschatellina foliis *Fumariæ* bulbosæ. *Raii Syn.* 267. *Baub. Hist. v.* 3: 206.

Moscatella. *Tillands. Ic.* 53.

Radix cava minima viridi flore. *Ger. em.* 1091.
Angl. Tuberous *Moschatell.*

In nemorosis, dumetis et ad sepes umbrosas.

Perennis. Fl. Aprili, Maio.

Radix repens, carnosa, imbricata, alba. Herba tenera, lævis, pallida, moschum redolens. Caulis spithamæus, simplex, erectus, quadrangulus, diphyllus. Folia ternata vel biternata, lobata; radicalia longius petiolata; caulina opposita. Capitulum terminale, solitarium, cubicum, quinqueflorum. Flores ochroleuco-virides, sessiles; laterales quinquesidi; terminalis quadrifidus. Stamina breviora, 8 vel 10. Styli laciniis corollæ numero æquales. Stigmata simplicia.

199. ELATINE. *G. Pl.* 198. *Juss.* 300.
Gærtn. t. 112.

Cal. tetraphyllus. *Petala* 4. *Capsula* supera, quadrilocularis, quadrivalvis, polysperma; dissepimenta futuris opposita.

† 1. ELATINE foliis verticillatis.

Alfinastrum.

Elatine Alfinastrum. *Sp. Pl.* 527. *Huds.* 173. *trum.*
With. 386.

Alfinastrum gratiolæ folio. *Dill. in Raii Syn.* 346.

A. gallii folio. *Vaill. Paris. t.* 1. f. 6.

Angl. Waterwort.

In fossis et paludibus, dubia cívís.

Found by Mr. J. Sherard on boggy ground, on the common just by the road from Eltham to Chislehurst. *Dill.* Not found, I believe, in England by any one since.

Perennis. Fl. Augusto.

Radix fibrosa, verticillata. Caules erecti, teretes, cavi, spithamæi, foliosi, basi ramosi. Folia verticillata; submersa ferè duodena, linearia, angustissima; emersa quaterna, ovata, integerrima, nervosa. Flores axillares, solitarii, sessiles, albi. Capsula globoso-depressa; valvulæ stylis persistentibus aristatæ. Semina numerosa, oblonga, octangulata, transversè rugosa.

Ad exemplaria optima, Parisiis lecta, descriptionem elaboravi; hæc communicavit egregius juvenis D. Stephanus Delessert, botanicus indefessus, nuper eheu! convulsis sedibus paternis, per Americam vagans, inopinatâ morte abreptus, periit.

Classis IX.

ENNEANDRIA.

Stamina 9.

HEXAGYNIA.

Styli 6.

200. BUTOMUS. *Cal.* nullus. *Pet.* fex. *Caps.*
fex, polyspermæ.

IX. ENNEANDRIA.

I. HEXAGYNIA.

200. BUTOMUS. *G. Pl.* 201. *Juss.* 46.
Gærtn. t. 19.

Cal. nullus. *Petala* 6. *Capsulæ* 6, superæ,
polyspermæ.

umbella- 1. BUTOMUS.

tus.

Butomus umbellatus. *Sp. Pl.* 532. *Huds.* 174.
With. 393. *Relb.* 164. *Sibth.* 134. *Ab-*
bot. 91. *Curt. Lond. fasc.* 1. *t.* 29. *Fl. Dan.*
t. 604. *Engl. Bot. t.* 651.

Butomus. *Raii Syn.* 273.

Juncus floridus. *Camer. Epit.* 781.

J. cyperoides floridus paludosus. *Lob. Ic.* 86.

Gladiolus palustris Cordi. *Ger. em.* 29 †.

Water Gladiole. *Pet. II. Brit. t.* 43. *f.* 3.

Angl. Flowering Rush.

In fossis et ad ripas fluviorum.

Perennis. *Fl.* Junio, Julio.

Formosa planta. Radix horizontalis, tuberosa. Folia radicalia, linearia, triquetra, integerrima, glabra, superne plana, acuta, ferè tripedalia, erecta. Scapus foliis altior, simplex, teres. Umbella terminalis, multiflora. Pedunculi basi bracteati. Flores rosei, inodori. Petala concava, alterna minora. Stamina receptaculo inserta. Germina carinata. Styli breves. Stigmata emarginata.

END OF THE FIRST VOLUME.

RB 7.12.1979

